

Successful Electroejaculation and Other Semen Procurement Methods from four Species of Rhinoceros (*Ceratotherium simum*, *Diceros bicornis*, *Rhinoceros unicornis*, *Dicerorhinus sumatrensis*)

N.E. Schaffer, DVM, MS.; Tom P. Meehan, DVM; R.S. Jeyendram, DVM., PH.D.:
SOS Rhino, 680 N. Lakeshore Dr. #807, Chicago, IL 60611, USA (nan@sosrhino.org);
Brookfield Zoo, Brookfield, IL 60513, USA; Andrology Laboratory Services,
680 N. Lakeshore Dr., Chicago, IL 60611, USA

Several semen procurement techniques, including penile massage, rectal massage, artificial vagina, and rectal probe electroejaculation were applied to four rhinoceros species. Penile massage, rectal massage, and artificial vaginas were applied to cognizant animals; rectal probe electroejaculation was applied to anesthetized animals. In some instances, multiple procedures were applied to individual animals. Success of these procedures was based on whether or not sperm was acquired. Penile massage was the direct manual massage of the penis. Rectal massage was manual manipulation of the accessory glands through the rectal wall. Artificial vaginas were applied to the erect penis during penile massage. Rectal probe electroejaculation utilized an ultrasound probe incorporated into a custom engineered electrical probe (Schaffer et al. 1998). These procurement techniques were applied to a total of 21 animals. Penile massage was successful for all 5 black, all 5 white, 2 of 4 Greater One-horned Asian and 3 of 5 Sumatran rhinoceroses. However, only two animals, one black and one Indian, became trained to reliably produce samples with sperm. Rectal massage facilitated other methods, but was unsuccessful when used alone on 2 black, 1 white, 2 Greater One-horned Asian and 2 Sumatran rhinoceroses. An artificial vagina was applied to 2 black, 1 white, 2 Greater One-horned Asian and 1 Sumatran rhinoceroses, but was effective in only the black rhino that had been successfully trained for penile massage. Rectal probe electroejaculation was applied successfully to 2 black, 2 white and 1 Greater One-horned Asian. Seminal fluid was successfully collected in all 12 attempts to electroejaculate these 5 animals. Sperm was not present, however, in the fluids of one attempt in the Greater One-horned Asian. Rectal probe electroejaculation proved the most repeatable and reliable technique in producing seminal fluid with sperm.

References

SCHAFFER, N. E.; BRYANT, W.; AGNEW, D.; MEEHAN, T.; BEEHLER, B.
Ultrasonographic monitoring of artificially stimulated ejaculation in three rhinoceros species (*Ceratotherium simum*, *Diceros bicornis*, *Rhinoceros unicornis*). *Journal of Zoo and Wildlife Medicine* 29:386-393 1998.

An IRF/SOS Rhino Sponsored Integrated Approach for the Enhancement of Reproductive Performance of White Rhinoceroses in the EEP

F. Schwarzenberger, C. Walzer, K. Tomasova, F. Göritz, T. Hildebrandt, R. Hermes

Univ. of Vet.med., A-1210 Vienna, Austria
Salzburg Zoo Hellbrunn, A-5081 Salzburg, Austria
Zoo Dvur Kralove, C-54401 Dvur Kralove, Czech Republic
Inst. of Zoo and Wildlife Biology, D-10307 Berlin, Germany

The goal of this IRF-SOS Rhino funded multi-disciplinary, multi-institutional research is to work on possible solutions to overcome the reproductive problems occurring in the captive white rhinoceros population. Until recently these problems were summarized as 1) acyclicity and variable estrous cycle lengths of 35 or 70 days, 2) mating failure due to acyclicity, or silent estrus due to sibling relationship/mate choice problems and 3) conception - pregnancy failure due to presumed uterine pathology. Reasons for these problems are not definitively identified, but our results suggest that in addition to the problems summarized above, male infertility and ovarian pathology are severe additional problems and causes for the low reproductive rate of captive white rhinoceroses.

Our combined approach works to overcome these problems; the techniques used include: 1) endocrine monitoring; 2) transfer of animals to enhance natural breeding; 3) clinical examinations of reproductive soundness in male and female white rhinos and 4) development of artificial insemination (AI). In addition, work on conditioning 2 males for manual semen collection is currently underway.

The goals of this project are to provide management recommendations, which will help to increase the reproductive rate of the captive population. In order to achieve this goal research into the causes for the low reproductive rate is necessary and thus currently the major focus is to increase our reference data basis of male and female reproductive biology and pathology. Particular focus is being placed on the old non-reproducing animals in order to conserve their genetic potential and to be able to better manage white rhinos in the future. Generation of the presently available data was only possible by combining the techniques of endocrine monitoring with ultrasonographic examinations and the development of a reliable anesthesia protocol. Although work is still in progress, currently available results make the white rhino EEP one of the most progressive due to direct inclusion of research results into management decisions.

Females with missing ovarian cycles are a major cause for the low reproductive rate of the captive white rhinoceros population. Ultrasound examinations in combination with endocrine monitoring revealed several reasons for the state of acyclicity. The two major groups are 1) females with no luteal activity (true flatliners) and 2) females with persistent luteal activity. In adult animals the reasons for acyclicity are either small cysts, 'micro

250 JAHRE
1752-2002
TIERGARTEN
SCHÖNBRUNN



Harald M. Schwammer
Thomas J. Foose
Michael Fouraker
Deborah Olson



A Research Update on Elephants and Rhinos

Proceedings of the
International Elephant
and Rhino Research Symposium,
Vienna, June 7-11, 2001