

200
Morph
G

IN MEMORIAM.

THE COLLECTED
SCIENTIFIC PAPERS

OF THE LATE

ALFRED HENRY GARROD, M.A., F.R.S.,

FELLOW OF ST. JOHN'S COLLEGE, CAMBRIDGE; FULLERIAN PROFESSOR OF PHYSIOLOGY
AT THE ROYAL INSTITUTION; PROFESSOR OF COMPARATIVE ANATOMY IN KING'S
COLLEGE, LONDON; PROSECTOR TO THE ZOOLOGICAL SOCIETY OF LONDON.

EDITED,

WITH A BIOGRAPHICAL MEMOIR OF THE AUTHOR,

BY

W. A. FORBES, B.A.,

FELLOW OF ST. JOHN'S COLLEGE, CAMBRIDGE; PROSECTOR TO THE ZOOLOGICAL SOCIETY OF LONDON.

LONDON:

R. H. PORTER: 6, TENTERDEN STREET, W.

1881.

5890
26/11/89



Page 2. 24. ON THE DEATH OF A RHINOCEROS IN THE SOCIETY'S GARDENS, AND ON SOME POINTS IN ITS ANATOMY.*

"Mr. A. H. Garrod, in drawing attention to the death on December 14th of the female *Rhinoceros unicornis*, which had lived in the Society's Gardens for more than twenty-three years, remarked that the only pathological sign detected was the enlargement of the lymphatic glands at the base of the heart. Mr. Garrod's observations on the visceral anatomy of this Rhinoceros were quite confirmatory of those of Professor Owen. In addition, he mentioned that there was a minute *os cordis* at the attached margin of one of the aortic valves, and that in the Perissodactyla this bone is not always absent, as by some supposed, he having found a large one in a Sumatran Tapir. The remarkable difference between the arrangement of the mucous membrane of the small intestine in the Indian and Sumatran Rhinoceroses (that of the former being produced into villi nearly an inch long through its whole length, whilst in the latter these were represented by *valvulæ conniventes*) was also illustrated from specimens in spirit."

Page 707. 25. ON SOME POINTS IN THE VISCERAL ANATOMY OF THE RHINOCEROS OF THE SUNDERBUNDS (*RHINOCEROS SONDAICUS*).†

OUR present knowledge of the visceral anatomy of the Rhinocerotidæ is confined to that of the two species *Rhinoceros unicornis* and *Ceratohinus sumatrensis*. Professor Owen has given us, in the "Transactions" of this Society (vol. iv. pp. 31 *et seq.*) an exhaustive account of the former of these animals; and in the "Proceedings" (1873, pp. 92 *et seq.*)‡ it has been my endeavour to indicate most of the important features in the latter, which, as Professor Flower has kindly pointed out to me, were briefly described by Sir E. Home in the

* "Proceedings of the Zoological Society," 1874, p. 2. Read, Jan. 6, 1874.

† "Proceedings of the Zoological Society," 1877, pp. 707-11. Read, Nov. 6, 1877.

‡ *Suprà*, p. 130.

"Philosophical Transactions" (1821, p. 271). On the present occasion I bring before the Society my notes on a young female of the Sondaic Rhinoceros (*Rhinoceros sondaicus*), which died in the menagerie of Mr. C. Jamrach, after having been in this country for a little more than half a year. It was only the skinned trunk which came into my possession. It is the nature of the mucous membrane of the small intestine which was certain to be of greatest interest; and this I am able to describe in detail.

The individual under consideration measures, stuffed, 6 feet 2 inches from the tip of the nose to the base of the tail. The tail itself is a foot long, whilst the height of the animal at the shoulder is 3 feet. From the middle of the occipital crest, along the curve of the superior surface of the skull, to the tips of the nasal bones is $13\frac{1}{2}$ inches, the same measurement in adult animals being 22 inches.

The single milk-incisor on each side of each jaw is still in place, as are all the milk-molars. The first true molar has not cut the gum; but its cap is seen within the bony alveolus. No traces of the other molars are visible.

Mr. E. Gerrard has kindly lent me the skull for examination. In Page 708. its base it exhibits the characteristic peculiarities of the species so clearly enunciated by Professor Flower,* the vomer being free behind and developed into a tongue-shaped process; the mesopterygoid fossa being expanded, and the free ends of the pterygoids everted at the same time that they are broad. No second combing-plate is present on the uncut first upper molar tooth.

The animal is too young to be contrasted advantageously with Professor Peters' drawing† of *Rhinoceros inermis*, Lesson. I have, however, taken the opportunity of comparing that figure with the skulls of *R. sondaicus* in the College of Surgeons' Museum, and fail to see that there are sufficient differences to justify specific differentiation. Professor Flower had previously done the same, and had arrived at a similar conclusion, as he found that even greater differences than those pointed out by Professor Peters are to be detected in individuals which are all undoubtedly of Indo-Malay origin.

In skin-folding and surface-texture the Sunderbund and Javan Page 709. specimens agree exactly; the young Sunderbund animal presenting a most striking uniformity in the size of the epidermic tuberculation, except in the gluteal region, where the boiler-bolt-shaped tubercles are somewhat larger than elsewhere. Along the back the scattered brown hairs, which spring from the yielding linear intertubercular surfaces, are also well developed.

* "Proceedings of the Zoological Society," 1876, p. 447.

† "Monatsb. der königl. Akad. zu Berlin," 1877, p. 68, pl. ii.

Fig. 1.

Mucous surface of duodenum of *Rhinoceros sondaicus*.

The following are the lengths of the alimentary viscera:—

Small intestine, 26 feet 2 inches.

Large intestine, 9 feet 10 inches.

Cæcum, 1 foot 3 inches.

The stomach, in shape, is very much like that of *R. unicornis* as figured by Professor Owen. Its cardiac surface is lined with the smooth white squamous epithelium found in all the Perissodactyla. This occupied about one third of the total gastric area, extending along most of the lesser curvature, the rest being covered with a smooth and thick digestive coat. There is no trace of any œsophageal valve like that found in the Horse.

The small intestine is somewhat larger in the duodenal region than elsewhere. Its first three inches are destitute of the flattened papillæ found elsewhere; but here, as all along the small intestines, minute villi are present everywhere. Three inches from the pylorus the papillæ commence, and resemble those similarly situated in *Rhinoceros unicornis*,* except that they are not quite so long. They are repre-

* *Vide* Professor Owen's figure, "Transactions of the Zoological Society," IV. pl. XII. fig. 1.

sented in Fig. 1 [p. 148], where they are seen to consist of flattened, round-tipped processes of the mucous membrane, several of which are blended at their bases, in transverse lines. None are more than $\cdot 3$ of an inch in length, and most about $\cdot 6$ inch broad where they first become free. They give the impression of being incomplete valvulæ conniventes which have been cut and deeply jagged at their free edges. The opening of the bile-duct is 7 inches from the pylorus, being a nipple-like tubular projection, nearly an inch long, among the papillæ. From the spot where they commence, all the way to the ileo-cæcal valve, these papillæ are found—those near the last-named situation differing from those in the duodenum in being more scattered and freer from one another, many in the ileum springing independently from the mucous membrane. Nowhere, however, are they otherwise Page 711.
than flattened, broad, and blunt-tipped, none anywhere being circular and slender like those in the ileum of *R. unicornis*,* the existence of which I have had the opportunity of verifying. They never exceed

Fig. 2.

Mucous surface of ileum of *Rhinoceros sondaicus*.

$\cdot 3$ of an inch in length. Numerous Peyer's patches exist in the ileum, as may be inferred from Fig. 2, which is a representation of


* "Transactions of the Zoological Society," IV. pl. XII. fig. 3.

a portion of the inner surface of the small intestine quite close to the ileo-cæcal valve.

Such being the case, *R. sondaicus* differs from *R. unicornis* in that the papillæ of the ileum are short, flat, and broad, instead of long, cylindrical and narrow, "like tags of worsted" (Owen).

The cæcum coli is a short blunt cone, with the diameter at its base as great as its length (1 foot 3 inches); and comparing the disposition of the colic flexures and proportionate diameter, I found them identical with those of the Sumatran species as I have figured them.*

Fig. 3.



Liver of *Rhinoceros sondaicus*. Visceral surface.

L.L. Left lateral. L.C. Left central. R.C. Right central. R.L. Right lateral. C. Caudate. Sp. Spigelian lobe.

The liver wants the gall-bladder, and differs but little from that of the Sumatran species. Fig. 3 is an outline-sketch of its abdominal surface, which, when compared with that of *Ceratorhinus sumatrenis* ("Proceedings of the Zoological Society," 1873, p. 102),† shows that the right central lobe is larger than the right lateral, instead of smaller. The Spigelian lobe is equally long and slender.

The pancreas is of good size and fairly concentrated.

The uterus is bicorn, each cornu measuring 8 inches, at the same time that the corpus uteri is 3 inches long. Each ovary is situated in a pocket of the peritoneum.

* "Proceedings of the Zoological Society," 1873, pp. 99, 100. (*Suprà*, pp. 137, 138.)

† *Suprà*, p. 141.