15

Copy in Zoology Library
Univ. Camborn

LIST OF QUADRUPEDS

BROUGHT BY

MR. BURCHELL

FROM

SOUTHERN AFRICA,

AND PRESENTED BY HIM TO

The British Puseum

On the 30th of September, 1817.

- 8. Hystrix cristata. (1.) (Porcupine.) Called Yzer-vark by the Dutch Colonists.
- 9. ORYCTEROPUS CAPENSIS. (Cape Anteater.) The Aardtvarh of the Colonists.
- 10. Phacocherus æthiopicus. (1.) The Vlakte-varh of the Colonists. Shot in the Cisgariepine, at "Wild-boar Station," on the banks of the Nugariep, or Black River.
- 11. Rhinoceros biconnis. (9.) Rhenoster of the Dutch Colonists. Killed at "Hot Station," on the Moshowa River, in the Transgariepine, in company with its dam, on the 28th October, 1812. This species is called Killenyán by the Bachapins; who distinguish the Rhinoceros simus B. (Trav. vol. ii. p. 75.), a hitherto nondescript and much larger species, by the name of Mokōhu.
- 12. Equus Quagga. (20.) The Quahha of the Hottentots and Colonists. It was shot from a troop of more than thirty, seen near Riet Fontein, on the Kowī River, in the eastern district of the Cape Colony, 14th October, 1813.
- 13. Equus Zebra. (22.) The Zebra of the earlier writers: vide "Travels," vol. i. p. 139. Called Pitze by the Bachapins. Shot at Little Klibbolikhónni Fountain, in the Transgariepine, on 24th November, 1812. This is the Equus Burchellii of later writers.
- 14. Equus montanus B. (2.) Vide "Travels," vol. i. p. 139. Shot on Wagenpadsberg, beyond the Snow Mountains, on the 27th March, 1813. This is the Dauw of the Hottentots, the Wilde Paard (wild horse) of the Dutch Colonists, and the Zebra of later writers.

- 15. Camelopardalis (5.) A male. Shot in the Transgariepine, on 26th December, 1812, at the Koraqua Klip-Fontein. The skin was accompanied by a complete skull. This animal was nearly 18 feet in height. The Camelopardalis is called $T\bar{u}hlua$ by the Bachapins; and Kameelpaard or Kameel by the Dutch colonists.
- 16. Camelopardalis A female. Shot at the Chue Spring, by the Maadji Mountains, on the 13th October, 1812. This skin had also its entire skull.
- 17. Antilope Euchore. (49.) The *Springboh* of the Colonists, and the *Tsépi* of the Bachapins. This animal was found drowned at "Amaryllis Station," on the Nugariep, or Black River.
- 18. Antilope melampus. (20.) Shot at Litākun, where it is called *Pāala* by the Bachapins, and *Roodebok* by the Hottentots of Klaarwater. *Vide* "Travels," vol. ii. p. 300.
- 19. Antilope Pygarga. (1.) Antilope albifrons of "Trav." vol. ii. p. 335. The Blesbok of the Colonists, and sometimes Bontebok. Shot near Zwellendam, on the 17th January, 1815, and the species is now become very scarce.
- 20. Antilope lunata B. (1.) ("Crescent-horned Antelope.") Vide "Travels," vol. ii. p. 334. Shot at the Makkwarin River, in the Transgariepine, 10th July, 1812. A very rare species.
- 21. Antilope villosa B. (7.) "Trav." vol. ii. p. 302. Ant. Capreolus Licht. It is called Pēeli, in the Sichuāna language; and is the Vaal Reeboh of the Colonists. A male, with horns exceeding the ordinary length; and shot at Zoetemelks River, in the district of Zwellendam, 19th November, 1814.