

Plate 17: Mastodonts from Arrisdrift, Namibia. (Scale bars = 5 cm)  
1-2. AD 527'99, *Afromastodon coppensi* nov. gen. nov. sp., right m/2, occlusal and lingual views.  
3-5. PQAD 2748, gomphothere molar fragment, posterior, occlusal and lingual views.

## Miocene Rhinocerotidae of the Orange River Valley, Namibia

Claude Guérin\*

\* UMR 5125 "Paléoenvironnements et Paléobiosphère" and UFR des Sciences de la Terre,  
Université Claude Bernard-Lyon I, 27-43 boulevard du 11 novembre 1918  
69622 Villeurbanne Cedex, France  
e-mail : Claude.Guerin@pop.univ-lyon1.fr

Among the Miocene sites excavated since 1991 by the Namibia Paleontology Expedition, two located in the Orange River Valley have yielded rhinoceros remains: Arrisdrift and Auchas Mine. An atlas vertebra and a mandibular fragment, both encrusted with sediment and thus specifically undeterminable, have been recorded from Auchas Mine. A fine lot of generally well preserved rhino material was found at Arrisdrift among a very diverse Vertebrate fauna. All but one of the 112 rhino pieces constitute a homogeneous sample pertaining to a very large species of cursorial rhino first described in 2000, *Diceros australis* Guérin, of which Arrisdrift is the type locality. The large form from Arrisdrift seems to be the largest of the African Miocene Rhinos; the size and proportions of the metapodials and the other limb bones suggest a strong analogy with *Diceros gr. pachygnathus-neumayri* of the Upper Miocene of the Near East; the type of construction of the upper cheek teeth, in particular the fourth premolar, is of Dicerotine kind and presents, as do the dimensions, close resemblances with *Diceros douaniensis* of the Upper Miocene of North Africa and Italy; the mandible shows analogies with the Dicerotines, especially the apparently short symphysis. *Diceros australis* is thus by far the oldest known species of the subfamily; the small reduced lower tusks could represent an evolutionary stage prior to the loss of the entire anterior dentition, which is effective in the subfamily since the Upper Middle Miocene. The exception among the Arrisdrift rhino material is an isolated magnum which suggests a small to medium sized short-legged form, probably *Chilotheridium pattersoni*, a species described from Loperot in Kenya, the age of which is about the same as Arrisdrift, i.e. 17 Ma.

### Version française abrégée:

La Namibia Paleontology Expedition, dirigée par B. Senut et M. Pickford, a fouillé depuis 1991 les gisements miocènes du Sperrgebiet (Pickford *et al.*, 1995). Quatre gisements à rhinocéros d'âge miocène moyen ont été découverts (Guérin, 2000), dont deux dans la vallée du fleuve Orange: Arrisdrift et Auchas Mine.

#### 1.- Les gisements :

Auchas Mine est un site diamantifère alluvionnaire situé à 50 km en amont d'Oranjemund (Pickford *et al.*, 1995), datant d'environ 18 millions d'années. Parmi les sept espèces de Mammifères recueillies dans la carrière AM 02 se trouve un rhinocéros indéterminé représenté par un atlas et un fragment de mandibule, tous deux encroûtés de sédiment et indéterminables spécifiquement.

Arrisdrift est un gisement très riche situé près du fleuve Orange, à 35 km à l'Est d'Oranjemund. Il date d'environ 17 M.A. et a livré de nombreux restes de Vertébrés (Pickford *et al.*, 1996). A une seule exception près, les 112 restes de rhinocéros constituent un matériel homogène appartenant à un très grand rhinocéros coureur décrit pour la première fois en 2000, *Diceros australis* Guérin, dont Arrisdrift est le gisement-type. L'exception est un magnum isolé totalement différent de ceux rapportés à *D. australis*: il évoque *Chilotheridium pattersoni* dont le gisement-type, Loperot au Kenya, est d'âge comparable à Arrisdrift.

#### 2.- Matériel et méthodes:

J'ai eu l'occasion d'étudier une bonne quantité de restes de rhinocéros du Miocène d'Afrique: diverses pièces des trois espèces de *Brachypotherium*, de bons moules de crânes et de dents de *Paradiceros mukirii*, quelques restes d'*Aceratherium campbelli*, des crânes et mandibules d'*Aceratherium acutirostratum* et de *Dicerorhinus leakeyi*, le type de *Diceros douaniensis* et tout le matériel connu de *Chilotheridium pattersoni* de Loperot. Celà ne constitue pas toutefois un ensemble d'échantillons suffisant.

Pour pallier le manque d'éléments post-crâniens en *Aceratherium* africains de taille moyenne j'ai utilisé comme terme de comparaison un regroupement de variables mesurées sur deux espèces miocènes européennes de taille et proportions voisines appartenant à la même lignée, *Aceratherium tetradactylum* et *A. incisivum*; dans les

tableaux de mesures l'échantillon correspondant est baptisé Acérathères. On ne connaît pas actuellement en Afrique de grande espèce miocène de *Dicerorhinus*, j'ai donc utilisé comme comparaison *Dicerorhinus schleiermacheri* du Miocène supérieur d'Europe. Enfin, pour avoir un bon échantillon de grands *Diceros* miocènes, j'ai regroupé sous le nom de *Diceros gr. pachygnathus-neumayri* du matériel de *D. pachygnathus* et *D. neumayri*, deux espèces très proches sinon identiques du Miocène de la région méditerranéenne orientale.

#### 3. - *Diceros australis* Guérin, 2000

La partie la plus intéressante du matériel recueilli comprend les fragments d'un crâne et de neuf hémimandibules, trois incisives inférieures, 32 dents jugales isolées, un humérus, quatre radius, trois cubitus, neuf carpiens, cinq métacarpiens, un tibia, treize tarsiens, neuf métacarpiens et plusieurs phalanges.

Trois des fragments d'hémimandibules sont suffisamment importants et comprennent une bonne part de la branche horizontale et de la symphyse, qui était probablement très courte. Le bord postérieur de la symphyse se situe dans tous les cas entre le premier tiers et la moitié de la longueur de la P/2. La brièveté de la symphyse suggère un faible développement sinon une absence d'incisives inférieures, ce qui est un caractère de Dicerotiné. Or nous disposons de deux petites incisives inférieures isolées; ces défenses ne devaient pas être fonctionnelles. Elles constituent par leur faible taille un stade évolutif antérieur.

Parmi les jugales, les plus significatives sont les troisième et quatrième prémolaires supérieures. Toutes présentent un ectolophe doté d'un fort parastyle, et un pli du paracône épais mais peu saillant; il n'y a ni mésostyle ni pli du métacône. Le principal repli interne est un fort crochet, et plusieurs prémolaires possèdent une médi fossette fermée. Il existe un fort cingulum lingual continu et crénelé, et le protocône ne montre aucune trace d'étranglement. Une telle morphologie, tout particulièrement celle de la muraille externe, est très proche de celle observée chez *Diceros douaniensis* et *D. gr. pachygnathus-neumayri*, et les dimensions sont très voisines. Les rangées dentaires inférieures présentent des longueurs du segment molaire et du segment P /3-P /4 très

proches de celles relevées chez *D. douariensis* et *D. gr. pachygnathus-neumayri*.

Un humérus gauche atteint environ 500 mm de long, soit 10 % de plus que les plus grands spécimens connus de *D. gr. pachygnathus-neumayri* et *D. schleiermacheri*, et ses proportions sont différentes.

Le radius dépasse lui aussi les plus grands *D. gr. pachygnathus-neumayri* et *D. schleiermacheri*, ce dernier étant par ailleurs plus élancé. Celui de l'*Aceratherium acutirostratum-Dicerorhinus leakeyi* indifférencié de Rusinga, dont les dimensions et proportions sont identiques à celles de notre échantillon d'Acérathères miocènes européens de taille moyenne, est plus petit et présente des proportions différentes. Les radius de *Chilotheridium* et tout particulièrement de *Brachypotherium snowi* sont plus trapus et beaucoup plus courts. Les caractères morphologiques du radius du grand rhinocéros d'Arrisdrift plaident en faveur d'un rapprochement avec la sous-famille des Dicerotinae.

Le plus grand des métacarpiens médians (Mc III) est un peu plus long que les plus grands spécimens connus de *Diceros gr. pachygnathus-neumayri* et *Dicerorhinus schleiermacheri*, mais un peu plus élancé. Ceux de *Brachypotherium snowi* et *B. heinzlini* sont plus courts et plus trapus, et il en est de même pour le petit *Chilotheridium*. Celui des acérathères européens montre plus ou moins les mêmes proportions mais est bien plus petit. Un diagramme de Simpson montre que le Mc III d'Arrisdrift n'appartient ni à un Brachypothère, ni à un Acérathère, mais qu'il présente des analogies avec *Diceros gr. pachygnathus-neumayri* et *Dicerorhinus schleiermacheri*.

Les métacarpiens abaxiaux (Mc II et Mc IV) sont plus longs que ceux des plus grands *Diceros gr. pachygnathus-neumayri* et *Dicerorhinus schleiermacheri* connus, et nettement plus élancés. Ceux de *Chilotheridium* sont bien plus courts et trapus, ceux des Acérathères sont plus courts et montrent des proportions très différentes.

Le seul tibia recueilli est très endommagé dans sa partie proximale mais sa longueur peut néanmoins être mesurée. Comme les Mc III il est un peu plus long que les plus grands *Diceros gr. pachygnathus-neumayri* et *Dicerorhinus schleiermacheri*, et plus élancé. On notera qu'une épiphyse brisée de *Diceros cf. douariensis* de Baccinello V3 en Italie présente des dimensions identiques à celles d'Arrisdrift.

Les dimensions et proportions de trois astragales sont proches de *Diceros gr. pachygnathus-neumayri*, et diffèrent peu de *Dicerorhinus schleiermacheri*. L'astragale de *Brachypotherium snowi* est un peu plus gros mais nettement plus bas, et ces proportions se retrouvent pour *Chilotheridium*. Chez les Acérathères les dimensions sont plus faibles pour des proportions différentes, et il en est de même pour l'échantillon indifférencié *Aceratherium acutirostratum-Dicerorhinus leakeyi*, dont on notera au passage les similitudes avec les Acérathères.

Parmi les cinq calcanéums trois sont adultes et complets. Ici encore, les dimensions et proportions sont voisines de *Diceros gr. pachygnathus-neumayri* et *D. schleiermacheri*, et n'ont rien à voir avec celles de *Paradiceros*, *Chilotheridium* et des Acérathères.

Le métatarsien médian (Mt III) est à peu près aussi long que celui des plus grands spécimens connus de *Diceros gr. pachygnathus-neumayri* et a sensiblement les mêmes

proportions sauf que son diamètre transversal sus-articulaire distal est relativement plus faible. Il est significativement plus long que chez *Dicerorhinus schleiermacheri*. Celui de *Chilotheridium* est bien plus court et relativement plus trapu. Celui des Acérathères est plus petit, avec des proportions différentes.

Les métatarsiens abaxiaux (Mt II and Mt IV) sont plus graciles et beaucoup plus longs que ceux de *Diceros gr. pachygnathus-neumayri* et *Dicerorhinus schleiermacheri*. Par rapport à *Chilotheridium* et aux Acérathères les différences sont les mêmes que pour le métarsien médian.

Le diagramme des rapports des segments de membres montre lui aussi des similitudes avec *Dicerorhinus schleiermacheri* et *Diceros gr. pachygnathus-neumayri*, avec toutefois une particularité notable, la bien plus grande longueur relative des métapodes abaxiaux.

La grande espèce d'Arrisdrift semble être le plus grand des Rhinocéros du Miocène d'Afrique; la taille et les proportions des métapodes et des os longs montrent une grande similitude avec les *Diceros gr. pachygnathus-neumayri* du Miocène supérieur du Proche-Orient; le type de construction des jugales supérieures est celui des Dicerotinae et montre, comme les dimensions, une forte ressemblance avec *Diceros douariensis* du Miocène supérieur du Maghreb et d'Italie; la mandibule évoque celle des Dicerotinae, notamment par sa symphyse apparemment très courte. Ce sont ces caractères qui nous permettent de l'attribuer au genre *Diceros*. *Diceros australis* est actuellement le plus ancien Dicerotinae connu; les défenses inférieures très réduites pourraient représenter un stade évolutif précédent immédiatement la perte totale de la denture antérieure, perte qui est accomplie au sein de la sous-famille dès la fin du Miocène moyen.

4. cf. *Chilotheridium pattersoni* Hooijer, 1971

L'exception parmi les restes de rhinocéros d'Arrisdrift est un magnum bien conservé. Par sa morphologie, sa taille et ses proportions ce magnum est totalement différent de ceux recueillis dans le même gisement et attribués à *Diceros australis*.

Bas et très large, doté d'une face antérieure aplatie et oblique, il présente un rapport largeur/hauteur inversé par rapport à *D. australis*, ce qui montre que nous avons affaire à une espèce petite à moyenne aux pattes courtes et fortes, probablement *Chilotheridium*, genre monospécifique dont l'espèce-type a été définie dans le site kényan de Loperot, à même âge qu'Arrisdrift.

Pour 10 spécimens incomplets de *C. pattersoni* de Loperot, D.A. Hooijer (1971) donne une hauteur antérieure légèrement plus forte mais une largeur antérieure un peu plus faible; ces différences ne sont toutefois pas significatives, car elles peuvent résulter d'une variation géographique, mais surtout d'une technique de mesures sans doute quelque peu différente; quoi qu'il en soit les ordres de grandeur sont les mêmes.

*Chilotheridium* est le seul Chilotherinae connu jusqu'à présent en Afrique. Sa découverte à Arrisdrift accroît très largement son aire de répartition, limitée jusqu'à présent au Kenya et à l'Ouganda. Du point de vue de son extension stratigraphique, il était limité au Miocène moyen (on le connaît jusqu'ici entre 18 et 11 Ma) mais la toute récente découverte d'un Mc IV gauche dans le gisement namibien de Grillental (20 à 21 Ma) montre que l'espèce remonte à la fin du Miocène inférieur.

## History of study

Remains of Miocene mammals were discovered about a century ago in what was at the time German South-West Africa. They were sent to Germany where E. Stromer studied them in 1926, but did not identify the rhinocerotid remains more precisely than "Rhinocerine g. et sp. indet". Several later articles re-examined this old collection, revising certain species and describing new ones, the latest being by R. Hamilton & J.A. Van Couvering (1977), who synthesized previous work and updated the faunal list. In this list the only identified rhinoceros is *Brachypotherium heinzlini*, based on a brief note by K. Heissig (1971) on a hemi-mandible from Langental already mentioned by Stromer.

Several sporadic finds occurred later on in Miocene and Quaternary sediments. In 1933, R. Heinz briefly described a skull, mandible and footprints of rhinos discovered near Kolmannskuppe (=Kolmanskop) and Charlottenthal; these remains, preserved at Lüderitz and identified as *Diceros bicornis*, were cited by C. Guérin & G. Demathieu (1993). In 1978 G. Corvinus collected an M3/ of *Ceratotherium* (South African Museum, SAM PQ 2126) in the Upper Grillental. Finally J. Schneider found a complete but eroded metatarsal III of a large *Brachypotherium*, in 1983, 8 km to the SE of Bogenfels (most probably at Glastal 1 or 2 (Pickford & Senut, 1999)); this specimen is preserved in the South African Museum, Cape Town (SAM PQ 2517).

It was from 1991 that the Namibia Palaeontology Expedition, led by B. Senut and M. Pickford, continued excavations in the Miocene deposits of the Sperrgebiet (Pickford et al., 1995; Pickford & Senut, 1999). Five localities yielded rhinoceroses, of which two are in the Orange River Valley, first excavated by Corvinus in 1976-1978. One of these sites, Arrisdrift, is particularly rich and contains a diverse fauna. In 1998, I was able to examine more than 80 fossils from the site, most of which belong to a new species, *Diceros australis*, which is the oldest known representative of the sub-family, Dicerotinae, to which belong the two extant species of rhinoceros. The results of this study were presented at the congress of the Palaeontological Society of South Africa held at Windhoek in September, 1998, and published a short while later (Guérin, 2000). A second visit to Namibia in May 2001, enabled me to examine the new material found at Arrisdrift since 1998, and thus to complete my study.

## Localities

Two localities, of Early and Middle Miocene age, in the Orange River Valley (Arrisdrift and Auchas Mine) were excavated by the Namibia Palaeontology Expedition.

**Arrisdrift:** This is a very rich site close to the Orange River 35 km east of Oranjemund. It is about 17 Ma and yielded abundant remains of vertebrates (Pickford et al., 1996). Apart from one specimen, all the 112 rhino fossils belong to a homogeneous sample of a large species of cursorial rhinoceros, *Diceros australis*, of which Arrisdrift is the type locality. The exception is a magnum, which is completely different from those attributed to *D. australis*: it is low and wide, evoking the small to medium sized forms with short feet. It could belong to *Chilotheridium pattersoni*, defined at Loperot (Kenya), a site with the same age as Arrisdrift.

**Auchas Mine:** Auchas Mine is a diamond-bearing deposit located 50 km upstream from Oranjemund (Pickford et al., 1995); it is aged 19 to 20 Ma. Among the 7 mammal species discovered in Pit AM 02 there is an indeterminate rhinoceros, represented by an atlas vertebra and a small fragment of mandible.

## Method of study and comparison

Even though the methods used are well known and accurate, there remains a difficulty concerning the elements of comparison: most of the Miocene rhinos of Africa are still poorly known, so I have used several samples from the Miocene of Western Europe and the Near East (Greece and Turkey).

## Methods

The methods of study used in this article are the same as those explained by C. Guérin (1980b), notably for the way of measuring, for the statistical treatment of the variables and for the non-quantifiable characters which are useful for discriminating teeth and skeletal remains of Rhinocerotidae. Simpson diagrams (or ratio diagrams) were systematically used to compare proportions of the main elements of the post-cranial skeleton; the reference always being a sample of some 30 adult specimens of extant *Diceros bicornis*.

## Miocene Rhinocerotidae of Africa

In the present state of knowledge the family Rhinocerotidae is represented in the Miocene of Africa by 6 lineages corresponding to sub-families. These lineages contain 9 genera and 14 species (Hooijer, 1973, 1978; Guérin, 1980b, 1989, 2000; Prothero et al., 1989). The absolute ages of many of the localities concerned were kindly furnished by M. Pickford.

**Aceratheriinae:** The aceratheres are medium to large forms, lacking horns, are cursorial and have the aspect of a large tapir. They have 4 metapodials and four functional digits in the front feet. They have a strong pair of lower tusks (the i/2), and their cheek teeth are very brachydont; the upper cheek teeth have an ectoloph which is more or less flat, without marked folds. They were aquaphile. Two species have been reported from Africa.

- *Aceratherium acutirostratum* (Deraniyagala, 1951) is of medium size. It is known from at least 10 sites: Alengerr Beds (14 to 12 Ma), Chemeron Formation - Northern Extension (5 to 4.5 Ma), Karungu (18 Ma), Moruarot Hill near Losidok (about 17.5 - 17.2 Ma), Ngorora Formation (12 to 11 Ma), Ombo (15 Ma), and Rusinga (18 Ma) in Kenya; Napak (19.5 Ma) in Uganda; Karugamania (more than 7 Ma) and Sinda (more than 6 Ma) in Congo.

- *Aceratherium campbelli* Hamilton, 1973 is large: it is only known from Jebel Zelten (about 17 to 16 Ma) in Libya.

**Dicerorhininae:** These are two-horned rhinos of medium to large size, with cursorial legs. During the Miocene, the tusks were well developed. The cheek teeth are relatively brachydont, the uppers (particularly P3/ and P4/) have the ectoloph adorned with two strong vertical folds, corresponding to the paracone and the metacone.

The face is long. They live most often in more or less wooded or bushland zones. One or two species are known in Africa:

- *Dicerorhinus leakeyi* Hooijer, 1966 is medium sized. It is recorded from at least 7 sites: Alengerr Beds (14 to 12 Ma), Chemeron Formation - Northern Extension (about 5 Ma), Karungu (18 Ma), Maboko (15.5 Ma), Ombo (15 Ma), Rusinga (18 Ma) in Kenya and Napak (19.5 Ma) in Uganda. It is noteworthy that in most of the sites that have yielded *D. leakeyi* the species *Aceratherium acutirostratum* also occurs, suggesting that they were sympatric.

- *Dicerorhinus primaevus* Arambourg, 1959, of the Late Miocene of Algeria, the generic position of which is debated (Geraads, 1986).

Up to now, large species of *Dicerorhinus* are not known in the Miocene of Africa, but they exist in Europe, such as *D. schleiermacheri* and several others (Guérin, 1980).

**Dicerotinae:** This sub-family contains the two extant species of African rhinos, *Diceros bicornis* ("Black" Rhinoceros) and *Ceratotherium simum* ("White" Rhinoceros). The genus *Ceratotherium*, more evolved of the two, with very hypodont cheek teeth, dates from the end of the Miocene, the species which it contains are grazers living in savanna. The genus *Diceros*, much less specialised, dates from the Middle Miocene and includes large, strongly built, two horned species. The face is short, the mandibular symphysis is short and there is no anterior dentition. The upper cheek teeth are brachydont, and possess a single well defined vertical fold on the ectoloph, the paracone fold. Its preferred habitat is spiny bushland in arid regions. Miocene forms of *Diceros* existed in the Maghreb, in Southern Spain, in Italy and the Near East (Greece and Turkey). Four species of Dicerotinae are known in the Miocene of Africa:

- *Diceros douariensis* Guérin, 1966: the species is defined at Douaria (9.5 Ma), in Tunisia, it is also present at Djebel Krechem el Artsouma (Late Miocene) in the same country (Geraads, 1989), as well as at Baccinello V3 (zone MN 13) in Italy (Guérin, 1980). Several remains found at Gravitielli (Late Miocene) in Sicily (Italy), as well as at Cenes de la Vega and Los Hornillos (both sites in MN 13) in the Grenada Basin, Spain, most likely belong to *Diceros* but the available material does not permit definite specific identification

- *Diceros australis* Guérin, 2000 is at present only known from Arrisdrift (Namibia) where it was defined. A detailed description is provided below.

- *Paradiceros mukirii* Hooijer, 1968 is a small species discovered at Fort Ternan (about 13 Ma) and perhaps at Maralal, also in Kenya, it is present at Kisegi (14 to 13 Ma) in Uganda, and at Beni Mellal (12.5 Ma) in Morocco (Hooijer, 1968; Guérin, 1976, 1994).

- *Ceratotherium praecox* Hooijer & Patterson, 1972, which is very abundant in the Pliocene of East and South Africa (Guérin, 1999), is also present in the Late Miocene, notably at Lothagam (7.2 to 5.5 Ma), Lukeino (6 to 5.5 Ma) and Mpesida (6.5 to 6 Ma).

**Iranotheriinae:** These are very large rhinos of Eurasia, heavily constructed with cheek teeth characterised by very marked hypodonty and the strongly folded enamel. The sub-family is often assimilated into the Elasmotheriinae but I consider that the resemblances are convergences and not due to identity. Two species of Iranotheriinae are known in Africa, one of which remains poorly known:

- *Kenyatherium bishopi* Aguirre & Guérin, 1974 has for the present only been found in Kenya, at Nakali (9.5 Ma) which is the type site, and at Samburu Hills, of similar age (Nakaya et al., 1999).

- *Ougandatherium napakense* Guérin & Pickford, 2003, is known from partial skeletons from Napak, Uganda. It is a small, hypodont form, with much cementum on the cheek teeth, and long slim metapodials.

**Brachypotheriinae:** The brachypotheres are large rhinos with a hippo-like appearance, with barrel-shaped bodies and short thick legs. They have strong tusks. The cheek teeth tend towards hypodonty, the ectoloph becoming flat. They were aquatic, their behaviour being similar to that of hippos. Three species are known in Africa, which, as noted by M. Pickford et al. (1993, p. 109) necessitate a revision:

- *Brachypotherium snowi* (Fourtau, 1920) has been found only at Wadi Moghara in Egypt and at Jebel Zelten (17 to 16 Ma) in Libya.

- *Brachypotherium heinzelini* Hooijer, 1963 is known from a dozen sites: Arongo Uyoma (Lower Miocene), Chemeron Formation - Northern Extension (about 5 to 4.5 Ma), Karungu (18 Ma), Rusinga (18 Ma) in Kenya; Napak (19.5 Ma) in Uganda; Karugamania (more than 7 Ma) and Sinda (more than 6 Ma) in Congo; Langental (18 Ma) in Namibia.

- *Brachypotherium lewisi* Hooijer & Patterson, 1972 is known from Kanapoi (4.5 Ma), Lothagam (7.2 to 5.5 Ma), Mpesida (6.5 to 6 Ma), Ngorora (12 to 11 Ma) in Kenya and at Sahabi (6.5 Ma) in Libya.

**Chilotheriinae:** The chilotheres, even though they have some resemblances to brachypotheres, constitute a separate sub-family. They are single horned, small, almost hippopotamoid forms with short legs (of which the front ones are tetradactyl). The tusks are small, the cheek teeth hypodont, and they were aquatic. A single species is present in the Miocene of Africa:

- *Chilotheridium pattersoni* Hooijer, 1971 is known from six localities in Kenya and Uganda, listed in the penultimate paragraph of this paper.

### Comparative material

For many years, I have had occasion to study a good quantity of Miocene African rhinoceros remains, in particular those stored at the Natural History Museum, London, where there are various specimens of three species of African *Brachypotherium*, good casts of the skull and teeth of *Paradiceros mukirii*, several remains of *Aceratherium campbelli*, skulls and mandibles of *Aceratherium acutirostratum* and *Dicerorhinus leakeyi*. I also had the opportunity to examine all the fossils attributed to *Chilotheridium pattersoni* collected at Loperot when they transited the Netherlands. Finally, the type specimen of *Diceros douariensis* is at the Université Claude Bernard-Lyon I, and the fossils from Baccinello V3 are kept at the Musée de Bâle, Switzerland. My comparative sample, suffers nevertheless from a scarcity of certain limb elements.

Apart from the fact that some species are rare, another reason for this lack is the affirmation of D.A. Hooijer (1966, 1973) who stated that it was impossible to distinguish the post-cranial skeletons of *Dicerorhinus leakeyi* from those of *Aceratherium acutirostratum*, which

are generally sympatric and closely similar in size. Because of this, no-one has tried, and the leg bones of these two species are not separated in the London collections, where, in any case, they are not sufficiently numerous to permit their identification taking into account individual variation. Having been confronted with similar problems at various stages when studying different families of Neogene and Quaternary mammals with several sympatric species, and having resolved them (see for example Guérin, 1980), I am convinced that such a distinction is possible as soon as there is enough material. Thus I believe that the abundant remains preserved in the National Museums of Kenya in Nairobi will permit a resolution of this problem. In the meantime, however, I have overcome the difficulty by using the following comparative terms:

- In the tables the undetermined specimens belonging to *D. leakeyi* and/or *A. acutirostratum* are referred to as IDA;

- Not having limb bones of medium sized African aceratheres, I have used measurements of two species from Europe that I consider to belong to the same lineage, and which have quite similar proportions, even though they are not identical. They are *Aceratherium tetracactylum* from the late Middle Miocene and early Upper Miocene, and *Aceratherium incisivum* from the Upper Miocene. In the tables, the sum of this addition, which does not exceed the limits of a single lineage, is referred to as Aceratheres; in the text it is called true aceratheres in order to avoid confusion with a certain number of related genera (*Alicornops* and others), also from the Miocene of Europe, which belong to different lineages and are therefore not considered further here.

- In the absence of large *Dicerorhinus* in the Miocene of Africa, I used *Dicerorhinus schleiermacheri* from the Upper Miocene of Europe.

- Finally, in order to have a good sample of large Miocene *Diceros*, I used remains of *Diceros pachynathus* from Pikermi in Greece (which are preserved in many museums in Europe) associated with others of *D. neumayri* from Turkey, preserved at the Museum of Munich in Germany. The status of these two species is unsettled, (for some they are synonyms, for others they are distinct but close), and I call the ensemble *Diceros* gr. *pachynathus-neumayri*.

### Systematics and Taxonomy

#### Sub-family Dicerotinae Genus *Diceros* Gray, 1821 Species *D. australis* Guérin, 2000

**Diagnosis:** Very large cursorial dicerotine. Upper cheek teeth brachydont, with more or less continuous crenulated lingual cingulum, the crochet being the only or the main internal fold. Ectoloph of the premolars with strong parastyle, paracone fold thick and not very projecting, and devoid of mesostyle and metacone fold. Upper molars with the ectoloph bearing a strong paracone fold and a weak vertical median fold, with the protocone having a weak constriction on the anterior surface. Limb bones long but robust. Abaxial metapodials remarkably long in comparison with the axial metapodial.

**Locus typicus and Stratum typicum:** Detritic fluvial diamictiferous sediments of Arrisdrift, Sperrgebiet,

Southern Namibia; early Middle Miocene, ca 17 Ma.

**Holotype:** Third left metacarpal AD 52'97 (Guérin, 2000, fig. 5. 3 and 4).

**Referred specimens:** Left demi-mandible AD 300'97; 4th right upper premolar AD 578'98 (Guérin, 2000, fig. 3: 3); Left upper 4th milk molar AD 292'94 (Guérin, 2000, fig. 3: 2); Right 3rd upper molar PQ AD 339 (Guérin, 2000, fig. 3: 4); Left lower 2nd premolar AD 86'98 (Guérin, 2000, fig. 3: 6 and 9); Left lower 3rd premolar AD 200'98 (Guérin, 2000, fig. 3: 7 and 10); Left lower 3rd milk molar PQ AD 635 (Guérin, 2000, fig. 3: 5 and 8); Left radius PQ AD 3099; Left astragalus AD 619'94 (Guérin, 2000, fig. 1: 2); Right calcaneum AD 50'97 (Guérin, 2000, fig. 1: 1); Right 3rd metatarsal AD 618'94 (Guérin, 2000, fig. 5: 1 and 2).

**Conservation:** all the material is preserved at the Geological Survey of Namibia in Windhoek (Namibia).

**Studied material:** 1 small fragment of skull; 9 fragments of mandibles (4 large and 5 small), each bearing part of the corresponding dentition; 3 lower incisors; 16 isolated upper cheek teeth; 16 isolated lower cheek teeth; 3 proximal fragments of scapula; 1 humerus; 4 radii, including a complete one, 3 ulnae including a complete specimen; 9 carpal (3 magnus, 2 pyramids, 2 semilunars, 1 trapezoid, 1 pisiform); 5 metacarpals (1 Mc II, 3 Mc III, 1 Mc IV); 1 tibia; 1 large fragment de fibula; 1 patella; 13 tarsals (3 astragali, 5 calcanea, 1 cuboid, 2 naviculars, 1 small cuneiform, 1 external cuneiform); 9 metatarsals (4 Mt II of which 2 are complete, 4 complete or almost complete Mt III, 1 entire Mt IV); 11 phalanges. All these specimens come from Arrisdrift.

**Description:** Skull. There is a fragment of the back of the skull corresponding to an almost complete occipital crest (the "chignon"). The transversal diameter, calculated by symmetry, is 220 mm; this value, greater than the maximum observed in *D. bicornis*, is close to the mean of extant *C. simum* (Guérin, 1980). The chignon is deeply excavated in its centre, forming a sort of saddle of which the arrow in the sagittal plane reaches 18 mm. The posterior surface (occipital) is excavated just below the large crest, for a height of a dozen cm, into a fossa subdivided into two by a vertical median crest; below the fossa, the occipital surface becomes vertically convex.

Mandible. Three large fragments of horizontal ramus possess part of the symphysis which was certainly remarkably short, even though none of them preserve the rostral part. The dorso-ventral symphysis diameter has a maximum value of 35 mm. In the three specimens, the posterior border of the symphysis is located at the level of the anterior third or in the middle of the p/2. The brevity of the symphysis suggests that the lower incisors were absent or vestigial, which is the usual case in Dicerotinae. At the break in the demi-mandible AD 505'99, one can see the root of a small i/2, the size and shape of which being similar to the isolated vestigial incisor AD 87'98.

On this hemi-mandible, the horizontal ramus has a weakly convex ventral edge, with an inflexion below the anterior edge of the ascending ramus, and the talon does not project behind the rear of the articular condyle.

In *Diceros douarsiensis* the symphysis is 100 mm long and 43 mm high, with the posterior margin located between p/2 and p/3; in *D. pachynathus* the same margin is at the level of the front of p/4. Even though the anterior part of the horizontal ramus of *D. australis* is lower, its posterior part (from the level of m/2) is comparable in size to those of *D. douarsiensis* (Tabl. 1). The African medium sized Aceratheriinae and Dicerorhininae are much smaller than *D. australis*, and *Paradiceros* is even more so; as for *Chilotheridium pattersoni*, its mandible has very different proportions.

Lower incisors. Three lower incisors were found. Recall that the tusks of rhinos are the i/2 and not canines, contrary to what was written by D.A. Hooijer (1971), who also described an internal margin sharpened by wear in *Chilotheridium* whereas this genus does not have upper incisors!

AD 88'98, unworn and in the shape of a button, has the appearance of a vestigial tooth and could be a di/1; its total length is 41.5 mm of which 11 mm comprise the crown; the transverse diameter at cervix is 9.5 mm for a dorso-ventral diameter of 11 mm.

AD 87'98 is a lower right tusk, only slightly worn (Pl.1, Fig. 1); the length is 84 mm of which 24 mm is the crown; the transverse diameter at cervix is 19 mm while the dorso-ventral measurement is 13.5 mm; The wear surface is oblique and is 11 mm long and 10 mm wide; the enamel

thickness suggests a permanent tooth, an i/2. The transverse section is an asymmetrical flattened ellipse. AD 22'99 is an unworn distal fragment of a similar tusk.

The tusks of *Chilotheridium* are much stronger; according to Hooijer (1971), their transverse sections reach respectively 22 x 17, 30 x 18, 30 x 15 and 40 x 25 mm, with a crown height in worn specimens of between 44 and 55 mm; their shape is different, more asymmetrical (Hooijer, 1971, pl. 6), and there is a cingulum. Because of their reduced size the Arrisdrift tusks could not have been very effective, whereas in all the other rhinos bearing tusks, they are much larger and were used by their possessors like bayonets.

I consider that the small adult tusks from Arrisdrift represent a stage towards the disappearance of the entire anterior dentition, which occurs in the Dicerotinae between the end of the Middle Miocene and the Upper Miocene. Such a disappearance also occurred in other rhino lineages, including the palearctic Dicerorhininae in which the tusks are functional until the end of the Miocene, but are reduced into vestigial buttons in *Dicerorhinus megarhinus* from the Ruscinian, before disappearing completely in later forms (Guérin, 1980).

Upper cheek teeth. There are two well preserved dM4/ and the ectoloph of a third specimen; AD 292'94 is lightly worn (Pl.1, Fig. 2), its maximum height is 38 mm which gives it a hypsodonty index of 81. A strong paracone fold

is the only relief on the external wall, and there is only one internal fold, the crochet. There is a discontinuous crenulated lingual cingulum, apparent mainly below the spout of the internal valley. AD 730'00, in medium wear, has similar morphology but does not have a lingual cingulum, and has a closed medifossette, probably due to the coalescence of the crochet and a crista. The dimensions of these two teeth are close to those of *D. douarsiensis* (Tabl. 2).

Four complete specimens of M1/ or M2/ were discovered:

- AD 228'97 is probably an M1/ in which the wear is slight, with a hypsodonty index of about 85. On the ectoloph there is a large paracone fold and a weak vertical median fold. The crochet, which is weakly developed, is the only internal fold. The protocone has a weak constriction on its anterior surface. A deep post-fossette is limited posteriorly by horizontally well developed posterior cingulum with a point and forming a horizontal enamel surface on the postero-lingual side of the tooth; the presence of this surface is constant in M1/ and M2/, the P/ and the dM/. A weak crenulated lingual cingulum occurs

Table 2: Comparisons of the measurements of the upper cheek teeth of *Diceros australis*. AP = antero-posterior; artic = articular; D = diameter; dist = distal; horiz = horizontal; prox = proximal; T = transversal.

ARRISDRIFT	<i>Diceros douarsiensis</i>												<i>B. snowi</i>												<i>A. campbelli</i>				
	AD 292'94	AD 578'98	AD 649'97	AD 228'97	AD 285'95	AD 339	PQ AD 2697	PQ AD 2661	PQ AD 1103	AD 720'00	AD 730'00	AD 259'99	AD 489'00	AD 490'00	AD 44,5	AD 46	AD 59	AD 63	AD 54	AD 64	AD 32,5	AD 41	AD 39	AD 43					
D 4/ L		47																											
DT		47																											
M 1/ L																59													
DT																56,5													
M 2/ L																58													
DT																62,5													
M 3/ L abs.																	64,5	65,5	66,5	65,5									
L anat.																	55	53	52	54									
DT																62	62	61	60										
P 2/ L																													
DT																													
P 3/ L																													
DT																													
P 4/ L																43,5	37,5												
DT																60,5	54												
<i>D. douarsiensis</i>																													
	n	mean	min.	max.												n	mean	min.	max.										
D 4/ L		1	44,00													2	61,50	55	68										
DT		1	49,00													3	71,00												
M 1/ L		2	60,25	59	61,5											2	67,00	63	71										
DT		3	61,33	59	64											2	76,50	74	79										
M 2/ L		3	62,83	60	67,5											3	61,17	59	64										
DT		2	68,75	66,5	71											1	63,00												
M 3/ L abs.		1	63,00													3	57,33	50	64										
L anat.		3	57,33	50	64											1	63,00												
DT		3	61,17	59	64																								
P 2/ L																													
DT																													
P 3/ L																													
DT																													
P 4/ L		2	40,50	37,5	43,5											1	49												
DT		2	57,25	54	60,5											1	69												
<i>B. snowi</i>																													
D 4/ L		1	44,00																										
DT		1	49,00																										
M 1/ L		2	60,25	59	61,5											2	61,50	55	68										
DT		3	61,33	59	64											1	71,00												

below the spout of the median valley.

- The other three specimens are probably M2/; their morphology is very similar, with a single fold on the ectoloph (the median swelling observed in the M1/ is not present), the paracone fold which projects strongly is separated from the parastyle which is perpendicular. AD 490'00 has a constriction on the anterior surface of the hypocone that is not present on any other cheek teeth from Arrisdrift.

The dimensions of these M1/ and M2/ (Tabl. 2) are slightly lower than those of *D. douaniensis*, and much smaller than those of *A. campbelli* but greater than those of *D. leakeyi*.

The four M3/s all have the same morphology (Pl. 1, Fig. 4): a strong paracone fold, a crochet being the only internal fold, protocone with weak constriction, a discontinuous lingual cingulum, a crenulated postero-labial cingulum covering the posterior quarter of the ectometaloph. The dimensions are close to those of the three M3/s of *D. douaniensis*.

The right P2/ AD 259'99 is deeply worn. The ectoloph, which is strongly oblique towards the apex (indicating the brachydont nature of this tooth) has only one relief feature, the paracone fold; there is an uncrenulated continuous lingual cingulum, which passes uninterrupted into the anterior cingulum. The greatest width is at the rear of the tooth; the dimensions are provided in Tabl. 2.

AD 720'00 is a well worn right P3/, in which the medifossette is closed and shows only a remnant of the crochet; the morphology is close to that of the preceding

tooth, with a stronger cingulum and larger size (Tabl. 2).

Two right P4/s were discovered, one in medium wear (AD 578'98) and the other deeply worn (AD 649'97). The ectoloph has a strong parastyle and a thick but not very projecting paracone fold; there is no mesostyle nor metacone fold. The only internal fold is the crochet, but specimen N° AD 649'97 has a closed medifossette. There is a strong continuous, crenulated lingual cingulum. The protocone has no sign of constriction. Such a morphology (Pl. 1, Fig. 3), and in particular the characters on the ectoloph - which constitute the best odontological criterion in the Rhinocerotidae, see Guérin (1980) - is very similar to that observed in *Diceros douaniensis* and *D. gr. pachygnathus-neumayri*, and this similarity applies also to the dimensions (Tabl. 2).

Lower cheek teeth. There are two deciduous molars available, a dm/2 and a dm/3. The latter tooth, which is slightly worn, has two internal valleys whose transverse profile is sharply V-shaped, and whose spouts are at the same level; there are traces of labial cingulum on the posterior lobe (Pl. 1, Figs 5 and 8). The dimensions (Tabl. 3) are similar to those (43 x 24 mm) of isolated dm/3s of *D. pachygnathus* from Pikermi preserved at the Bologna Museum (Italy).

Two isolated m/2s of which one is complete and three isolated m/3s in good condition were collected (Pl 1, Figs 7 and 10). The latter are wider than the only known m/3 of *D. douaniensis*, and almost the same size as those of *Brachypotherium heinzelini* and *B. snowi*; they are much larger than the m/3 of *Chilotheridium* and *Paradiceros*, as well as those of African medium-sized Aceratheriinae and Dicerorhininae (Tabl. 3). The anterior internal valley has a

sharp V-shaped profile; the posterior one is V-shaped in one case and U-shaped in two others; the difference in level of the spouts varies from medium to strong. There is no labial or lingual cingulum, but all the m/3s possess a posterior crenulated cingulum which varies considerably in shape.

Eight isolated lower premolars can be studied: three p/2s, four p/3s and one p/4. The p/3 n° AD 157'95, completely unworn, has a hypsodonty index of 108; the two internal valleys are V-shaped and show marked differences in the level of their spouts; there is no labial or lingual cingulum, but there are anterior and posterior ones, which extend a little onto the labial surface, a feature that also occurs in the p/2 (Pl 1, Figs 6 and 9). The p/3 AD 731'00 has a trace of an external cingulum on its labial surface at the base of the median synclinal.

The lower cheek teeth present in tooth rows show the same features. In the hemi-mandible AD 356'00 the m/2 and m/3 possess an anterior cingulum that extends on the lingual side as far as the centre of the anterior valley; the m/1 is too worn to yield any information on this character, which does not occur in the dental row of AD 556'94. The most complete tooth row, (AD 300'97) has a segment comprising "two last premolars" and a segment "molars" whose respective lengths are very close to those of *D. douaniensis* and *D. gr. pachygnathus-neumayri* (Tabl. 3).

Scapula. Three proximal fragments (= articular) of scapulae were found. The largest, AD 506'99, has a transverse articular diameter of 82.5 mm (it is greatest in the middle of the surface) for an antero-posterior diameter of 101 mm, and its tuberosity is massive. The other two specimens are smaller, with the greatest diameter located at the rear of the articulation, and they possibly represent immature individuals.

Humerus. A left humerus (AD 736'00) (Pl. 4) was recovered during the 2000 field season. It is incomplete proximally, the most proximal point being about mid-height of the greater trochanter; it measures 483 mm in length, and it was probably about 500 mm in total length. The transverse diameter of the diaphysis is 72.5 mm and the transverse and antero-posterior diameters of the distal

epiphysis are 170 and 127 mm respectively.

The humerus of *D. schleiermacheri* is shorter (426 to 448 mm, the mean of four: 435.7 mm), with a narrower diaphysis (60.5 to 63.5 mm) and a narrower distal epiphysis (130 to 146 mm) but almost as deep.

That of *D. gr. pachygnathus-neumayri* is also shorter (409 to 447 mm, mean of five: 429 mm), but its diaphysis is as wide (65 to 81 mm) and its distal epiphysis, for the same width (146 to 182 mm), is a bit less developed antero-posteriorly (102 to 120 mm).

Radius and ulna. Comparisons of the dimensions and proportions (Tabl. 4; Fig. 1) show that the radius is longer than the biggest specimens observed of *D. gr. pachygnathus-neumayri* and *D. schleiermacheri*, the latter being slimmer. The undifferentiated aceratheres-dicerorhine from Rusinga, which has the same size and proportions as the medium-sized late Miocene aceratheres of Europe, is smaller and has different proportions. The radius of *Chilotheridium* and above all, of *Brachypotherium snowi* is more robust and much shorter. The proximal articulation, comprising a lateral facet and a medial facet, does not have the undulating anterior border nor the large re-entrant angle at the level of the coronoid process which characterises the genus *Ceratotherium*. It shows, however, a lateral facet which extends further transversally and the anterior margin of which is located only slightly behind the anterior border of the medial facet; the posterior margin of the lateral facet, which is regularly concave, makes an obtuse angle with the posterior border of the medial facet. These latter characters are typical of the genus *Diceros* (Guérin, 1980).

Three ulnae were discovered at Arrisdrift of which only one (AD 273'97) is complete; its maximum length reaches 533 mm; its proximal articular diameter is 95 mm and its antero-posterior diameter is 157 mm. The other two specimens consist of a distal epiphysis of an adult, and a proximal extremity of a young individual.

Carpals. The only complete semi-lunar has an anterior surface in which the width and height are identical: 42 mm; the most distal point of this surface, close to the median line, is rounded. The length is 69.5 mm. Only one dimension can be obtained from fragment AD 824'99, the


Table 3: Dimensions comparées des jugales inférieures de *Diceros australis*. AP = antéro-postérieur; artic = articulaire; D = diamètre; dist = distal; horiz = horizontal; prox = proximal; T = transversal.

ARRISDRIFT	AD	PQ AD	AD	AQ	AD	AD	AD	PQ AD	AD	AD	AD	AD	AD	AD	AD	AD	AD	AD	AD	AD	AD	AD	AD	
	8'98	635	44'98	200'98	468'97	163'97	378'94	134	86'98	157'95	432'95	827	330	556'94	300'97	313'98	437'97	223'97	260'99	731'00	355'00	356'00	505'99	
D /2	L	30																						
	DT	18																						
D /3	L		45																					
	DT		25,5																					
D /4	L																							
	DT																							
M /1	L																							
	DT																							
M /2	L																							
	DT																							
M /3	L																							
	DT																							
P /2	L																							
	DT																							
P /3	L																							
	DT																							
P /4	L																							
	DT																							
L P/3-P/4																								
L M/1-M/3																								
<i>Diceros douaniensis</i>		n	mean	min.	max.	<i>B. heinzelini</i>		<i>B. snowi</i>		<i>Chil. pattersoni</i>		n	mean	min.	max.	AD	East Afr.	<i>P. mukiri</i>		<i>F. terner</i>				
D /2	L	2	48,00	47	49	<i>B. heinzelini</i>		<i>B. snowi</i>		<i>Chil. pattersoni</i>		2	42,00	40,5	43,5	39,00	35	<i>P. mukiri</i>		<i>F. terner</i>				
D /3	DT	1	30,50	27	34,5	<i>B. heinzelini</i>		<i>B. snowi</i>		<i>Chil. pattersoni</i>		2	30,00	27	33	30,00	27	<i>P. mukiri</i>		<i>F. terner</i>				
D /4	DT	1	36,75	55,5	58	<i>B. heinzelini</i>		<i>B. snowi</i>		<i>Chil. pattersoni</i>		3	54,50	50,5	57	47,00	42	<i>P. mukiri</i>		<i>F. terner</i>				
M /1	L	2	33,75	32	35,5	<i>B. heinzelini</i>		<i>B. snowi</i>		<i>Chil. pattersoni</i>		3	31,00	29,5	32,5	32,00	26	<i>P. mukiri</i>		<i>F. terner</i>				
M /2	L	1	54,00			<i>B. heinzelini</i>		<i>B. snowi</i>		<i>Chil. pattersoni</i>		2	59,50	56	63	54,00	43,5	<i>P. mukiri</i>		<i>F. terner</i>				
M /3	L	1	3																					

**Table 4:** Comparisons of the measurements of the radius of *Diceros australis*. AP = antero-posterior; artic = articular; D = diameter; dist = distal; horiz = horizontal; prox = proximal; T= transversal.

ARRISDRIFT		<i>B. snowi</i>				IDA			
	PQ AD 3099	AD 731'99	AD 506'00	AD ss n°	Djebel Zeiten	Rusinga			
Length	435				286,5	305			
DT prox.	113		114		86,5	95			
DAP prox.	70,5		68,5		51,5	57,5			
DT dia.	64,5		68		48,5	51,5			
DAP dia.	42		39,5		40	47,5			
DT dist.	115		107		86	92			
DAP dist.	83		73,5		53,5	69			
DT artic. dist.	103	102	100,5		80,5				
DAP artic. dist.	54	51	52,5		41,5				
<i>D. gr. pachygnathus/neumayri</i>									
n	Mean	min.	max.	éc.-type	coeff. var.	n	Mean	min.	max.
Length	9	364,22	342	375	9,536	1	327,00	327	327
DT prox.	9	100,61	95	107	3,790	2	94,50	94	95
DAP prox.	9	65,00	55	72	5,315	1	53,00	53	53
DT dia.	9	59,11	55	63	2,667	2	45,00	44	46
DAP dia.	9	38,78	33	47	4,402	2	32,50	32	33
DT dist.	8	104,00	95	109,5	4,488	2	93,00	91	95
DAP dist.	9	65,78	60	71,5	3,833	2	56,75	54,5	59
DT artic. dist.						2	85,50	85	86
DAP artic. dist.									
<i>D. schleiermacheri</i>									
n	Mean	min.	max.	éc.-type	coeff. var.	n	Mean	min.	max.
Length	5	369,30	351,5	396	18,620	5	341,17	316	348
DT prox.	4	99,38	94,5	104,5	4,328	9	86,67	78,5	100
DAP prox.	3	64,00	63	66	1,732	10	54,25	50,5	59,5
DT dia.	5	54,20	48	57,5	3,785	6	48,33	44	52,5
DAP dia.	5	40,60	38	43	1,782	4,39	6	37,92	33
DT dist.	5	95,20	87	100	5,707	6	85,17	75	103,5
DAP dist.	5	65,60	61	69	3,991	7	54,50	50	61
DT artic. dist.	4	82,12	79	88,5	4,385	5	72,60	68,5	79,5
DAP artic. dist.	4	44,75	44	45	0,500	5	38,70	36,5	42,5
ACERATHERES						5	38,70	36,5	42,5

Bivariate plot magnums



**Figure 2:** Dispersion diagram of anterior breadth as a function of sub-articular height (abscissa) of three magnums of rhinos from the Miocene of Namibia, other Miocene rhinos and extant *Diceros bicornis*.

height of the anterior surface, which is slightly greater than 40 mm.

Only one of the two pyramidal collected is well preserved (PQ AD 3173); it is remarkably large, and clearly wider (71.5 mm) than high (63 mm).

The only pisiform preserved is 70 mm long and 29 mm wide with a height of 52 mm.

Two out of the three magnums found at Arrisdrift are complete, of which one (AD 538'97) is attributable to *D. australis* (Pl. 3, Fig. 4). In anterior view the bone has a rhomboidal outline which is rounded distally, and it is higher than wide. The distal articulation is about as wide in front as it is behind. The dimensions are as follows:

Total length: 104 mm

Anterior width: 56 mm

Anterior height: 45 mm

Maximum height: 74 mm

Sub-articular height: 72 mm

These dimensions and the proportions differ from those of *D. schleiermacheri* but are similar to three magnums of *Diceros gr. pachygnathus-neumayri*; they are completely different from those of the other complete magnum found at the site (Pl. 3, Figs 3-5 and text-fig. 2).

The only trapezoid known (AD 141'95) measures 44.5 mm in length and 26 mm width for a height of 35.5 mm.

**Metacarpal II.** The only specimen, a left one, is longer than the biggest known specimens of *Diceros gr. pachygnathus-neumayri* and *Dicerorhinus schleiermacheri*, but is clearly slimmer. That of *Chilotheridium* is very short and thick, whereas that of true aceratheres is shorter and has very different proportions (Tabl. 5).

The proximal articulation is long and narrow, with a crescent shaped outline, with a clear notch on its posterior margin. On the lateral surface of the epiphysis there is a single articular facet, constricted in its median part but extending right across the bone in its median part. The transverse section of the diaphysis is a rounded triangle.

**Metacarpal III.** Two complete left and one right Mc III were found (Pl. 2, Figs 3 and 4) and the largest one is bigger than the maximal values known in *Diceros gr. pachygnathus-neumayri* and *Dicerorhinus schleiermacheri*, but it is somewhat slimmer. The Mc III of *Brachypotherium snowi* and *B. heinzelini* are much shorter and thicker, and it is the same for *Chilotheridium*. Those of true aceratheres have more or less similar proportions but are much smaller (Tabl. 6).

A Simpson diagram (Fig. 3) shows that Mc IIIs from Arrisdrift do not belong to a brachypothere, nor to an aceratheres, but that it has clear analogies to those of *Diceros gr. pachygnathus-neumayri* and *Dicerorhinus schleiermacheri*.

The proximal articulation is very wide, triangular with a straight anterior margin. On the lateral surface of the proximal epiphysis there are two articular facets; the anterior one is pentagonal and subdivided into two parts of which the inferior one is more or less elongated; the posterior facet, which is lower than the anterior one, is a rounded triangle, of which the width varies from individual to individual. On the medial surface of the proximal epiphysis there is an articular facet in the shape of a sleeping S which is variable in height. The transverse section of the diaphysis is trapezoidal with a weakly convex anterior border and a slightly concave posterior

**Table 5:** Comparisons of the measurements of the Mc II of *Diceros australis*. AP = antero-posterior; artic = articular; D = diameter; dist = distal; horiz = horizontal; prox = proximal; T= transversal.

ARRISDRIFT									
AD 536'97									
Length	188								
DT prox.	41								
DAP prox.	54								
DT dia.	32,5								
DAP dia.	25								
DT max. dist.	43								
DT artic. dist.	42								
DAP dist.	45,5								
<i>D. gr. pachygnathus/neumayri</i>									
n	Mean	min.	max.	éc.-type	coeff. var.	n	Mean	min.	max.
Length	9	156,33	140,5	169	8,842	5,66	2	129,50	129,5
DT prox.	9	44,67	37	54	5,385	12,06	2	43,00	42
DAP prox.	6	46,50	40	51	4,231	9,10	2	37,25	37
DT dia.	9	40,78	33,5	44,5	4,374	10,73	2	31,75	31
DAP dia.	9	24,33	20	26	1,750	7,19	2	16,50	16
DT max. dist.	9	48,89	40	54	5,355	10,95	2	39,75	37
DT artic. dist.	9	43,17	34,5	48,5	4,644	10,76	2	34,00	31
DAP dist.	9	42,39	38,5	46	2,583	6,09	2	36,25	36
<i>Chilotheridium pattersoni</i>									
n	Mean	min.	max.	éc.-type	coeff. var.	n	Mean	min.	max.
Length	2	129,50	129,5	129,5		2	43,00	42	44
DT prox.	2	43,00	40	47		2	37,25	37	37,5
DAP prox.	2	37,25	37	37,5		2	31,75	31	32,5
DT dia.	2	16,50	16	17		2	16,50	16	17
DAP dia.	2	39,75	37	42,5		2	34,00	31	37
DT max. dist.	2	34,00	31	37		2	36,25	36	36,5
DT artic. dist.	2	34,00	31	37		2	36,25	36	36,5
DAP dist.	2	36,25	36	36,5					
<i>D. schleiermacheri</i>									
n	Mean	min.	max.	éc.-type	coeff. var.	n	Mean	min.	max.
Length	3	168,83	156,5	179	11,405	6			

**Table 6:** Comparisons of the measurements of the Mc III of *Diceros australis*. AP = antero-posterior; artic = articular; D = diameter; dist = distal; horiz = horizontal; prox = proximal; T= transversal.

ARRISDRIFT			<i>B. snowi</i>		<i>B. heinzelini</i>	
	AD 52 '97	AD 243 '95	Jebel Zeiten	Rusinga		
Length	200	212	219	159,5	149,5	
DT prox.	68	63,5	64,5	74,5	66	
DAP prox.	58	51,5	56,5	57	52	
DT dia.	57,5	58,5	61,5	60,5	53	
DAP dia.	27	26,5	26	22,5	24,5	
DT max. dist.	65	63,5	67,5	73,5	72,5	
DT artic. dist.	60	60	58	58	58,5	
DAP dist.	48	44*	54,5	46,5		
<i>D. gr. pachygnathus/neumayri</i>						
n	mean	min.	max.	éc.-type	coeff. var.	
Length	9	187,17	181	198	5,668	3,03
DT prox.	6	65,50	62,5	74,5	4,461	6,81
DAP prox.	8	54,94	52	59	2,427	4,42
DT dia.	11	63,55	59	69	3,020	4,75
DAP dia.	9	24,56	22	26,5	1,333	5,43
DT max. dist.	9	71,22	66,5	76	2,705	3,80
DT artic. dist.	10	56,15	52	59	2,174	3,87
DAP dist.	10	48,05	45	53	2,619	5,45
<i>Chilotheridium pattersoni</i>						
n	mean	min.	max.	éc.-type	coeff. var.	
Length	3	157,67	150	169		
DT prox.	3	54,33	50	61		
DAP prox.	3	41,67	37	45		
DT dia.	3	40,83	39	43		
DAP dia.	3	19,17	17,5	21,5		
DT max. dist.	3	54,83	51	61		
DT artic. dist.	3	46,33	44,5	49,5		
DAP dist.	3	38,17	37,5	39		
<i>D. schleiermacheri</i>						
n	mean	min.	max.	éc.-type	coeff. var.	
Length	6	195,42	181	204	8,546	4,37
DT prox.	9	62,67	58	69	3,700	5,90
DAP prox.	5	50,30	47,5	52	1,987	3,95
DT dia.	9	55,78	49,5	66	5,363	9,61
DAP dia.	8	23,38	22	25,5	1,188	5,08
DT max. dist.	7	63,50	60	69	3,149	4,96
DT artic. dist.	7	52,14	49	56	2,478	4,75
DAP dist.	7	46,71	45	49	1,410	3,02
<i>Aceratheres</i>						
n	mean	min.	max.	éc.-type	coeff. var.	
Length	10	163,75	139,5	181	13,382	8,17
DT prox.	10	53,35	50	59,5	3,092	5,80
DAP prox.	10	43,80	40	47,5	2,406	5,49
DT dia.	11	45,55	42	49	2,252	4,95
DAP dia.	10	20,20	17	24,5	2,163	10,71
DT max. dist.	10	55,25	49,5	60,5	3,676	6,65
DT artic. dist.	10	47,00	38,5	52	3,894	8,29
DAP dist.	7	41,71	38	44,5	2,563	6,15


one; the straight lateral margin is longer than the medial one which is more or less straight.

Metacarpal IV. The bone is slimmer and much longer than the biggest known specimens of *Diceros* gr. *pachygnathus-neumayri* and *Dicerorhinus schleiermacheri* (Tabl. 7).

The proximal articulation is triangular, a little longer than wide; this length/width ratio is the opposite in rhinos

with short feet such as *Brachypotherium* and *Chilotheridium*. The medial surface of the proximal epiphysis has two articular facets; the anterior one, semi-elliptical in outline, is long and low, whereas the posterior one is a vertical ellipse which is much higher than wide.

Tibia. The only tibia recovered is very damaged; in particular at its proximal end which prohibits appreciating whether its morphology is typical of Dicerotinae or not, but



**Figure 3:** *Diceros australis*: Simpson diagram of Mc III compared with other Miocene rhinos. The reference is *Diceros bicornis*. 1: Length; 2: DT proximal; 3:DAP proximal; 4: DT diaphysis; 5: DAP diaphysis; 6: DT maximal distal; 7: DT articular distal; 8: DAP distal.

**Table 7:** Comparisons of the measurements of the Mc IV of *Diceros australis*. AP = antero-posterior; artic = articular; D = diameter; dist = distal; horiz = horizontal; prox = proximal; T= transversal.

ARRISDRIFT		GRILLETAL		IDA	
	AD 404'97		GT 31'01	N	moyenne
Length	ca 188		127	2	158,75
DT prox.	55			1	52,00
DAP prox.	51,5		41*	1	46,00
DT dia.	37,5		29,5	2	34,00
DAP dia.	27		21	2	21,25
DT max. dist.	51		42	2	46,25
DT artic. dist.	45		38,5	2	43,00
DAP dist.	53		37*	2	38,75
<i>D. gr. pachygnathus/neumayri</i>		<i>Chilotheridium pattersoni</i>		<i>Chilotheridium pattersoni</i>	
n	mean	min.	max.	éc.-type	coeff. var.
Length	8	144,81	134	156	9,059
DT prox.	8	47,81	43	53	3,535
DAP prox.	8	42,12	37	47,5	3,410
DT dia.	8	37,69	31,5	41	3,162
DAP dia.	8	24,75	20	29	2,726
DT max. dist.	8	46,69	39	52,5	4,166
DT artic. dist.	8	43,06	37,5	47	3,590
DAP dist.	8	41,00	37,5	46	2,712
<i>D. schleiermacheri</i>		<i>ACERATHERES</i>		<i>ACERATHERES</i>	
n	mean	min.	max.	éc.-type	coeff. var.
Length	1	145,00			
DT prox.	2	49,25	47,5	51	
DAP prox.	1	43,00			
DT dia.	2	32,50	32,5	32,5	
DAP dia.	2	21,50	21	22	
DT max. dist.	2	45,25	43	47,5	
DT artic. dist.	2	42,25	38,5	46	
DAP dist.	2	40,50	40	41	

**Table 8:** Comparisons of the measurements of the tibia of *Diceros australis*. AP = antero-posterior; artic = articular; D = diameter; dist = distal; horiz = horizontal; prox = proximal; T= transversal.

ARRISDRIFT		<i>Chilotheridium pattersoni</i>		
	PQ AD 561	min.	max.	
Length	440	319	355	
DAP dia.	57	47		
DT dist.	100	88		
DAP dist.	80	66		
DT artic. dist.	87			
DAP artic. dist.	75			
<i>D. gr. pachygnathus/neumayri</i>		<i>D. cf. douariensis</i>		
		Baccinello V3		
n	mean	min.	max.	
Length	5	390,40	368	420
DAP dia.	7	58,36	49	73
DT dist.	6	107,92	99	115
DAP dist.	7	79,00	71	99
DT artic. dist.		10,724		13,57
DAP artic. dist.			58	101
<i>D. schleiermacheri</i>		<i>ACERATHERES</i>		
		n	mean	
Length	2	398,00	386	
DAP dia.	2	58,25	57,5	
DT dist.	2	103,50	101	
DAP dist.	2	72,25	69	
DT artic. dist.	1	82,50		
DAP artic. dist.	1	60,00		
<i>ACERATHERES</i>				
n	mean	min.	max.	
Length	9	361,22	328	383
DAP dia.	9	47,39	38	53
DT dist.	9	86,17	72,5	91,5
D				