

**Examine the Possibilities of Enlarging/Managing
Larger Habitats for the Re-introduced Rhino with
Planning details in Dudhwa Tiger Reserve**



SATYA PRAKASH, IFS, PG DIPLOMA TRAINEE

SUPERVISOR

Dr. SATYA PRIYA SINHA

A Report – Up Dated (Oct,2002)

**WILDLIFE INSTITUTE OF INDIA, DEHRADUN
(UTTARANCHAL STATE)**

JUNE 2002

CONTENTS

ABSTRACT	1
INTRODUCTION	2
BACKGROUND OF RHINO RE-INTRODUCTION PROGRAMME IN DUDHWA TIGER RESERVE, UTTAR PRADESH	6
OBJECTIVE	14
METHODOLOGY	14
STUDY AREA	15
DISCUSSION	16
PROPOSED ACTION PLAN	21
THE EXPECTED OUTCOME	25
CONCLUSION	26
BIBLIOGRAPHY	27
APPENDIX -CLIPPING	32
ACKNOWLEDEMENT	33

"Informed conservationists generally agree that an endangered species whose habitat type has by human impact been reduced and split up into separated "islands", should live in as many of those islands as possible. This reduces the risk of extinction."

Dr. R. Schenkel

Abstract

The Great Indian One Horned Rhinoceros once inhabited throughout the Indus, Brahmaputra and Gangetic flood plains and nearby foothills of South Asia. Due to rampant poaching and loss of suitable habitat, rhinoceros are now restricted to a few isolated pockets of protected areas in Assam and West Bengal in India and Nepal as well. Few Great Indian One-Horned Rhinoceros roaming in the Terai area of Central North India are believed to have disappeared in 1878 near Pilibhit, Uttar Pradesh. One of the conservation measures adopted for the species is the establishment of additional viable populations in suitable areas in the former ranges of its distribution to increase the areas under rhinoceros, thereby reducing the risk of extinction. In the years 1984 and 1985, rhinos from Assam and Nepal were re-introduced in the Dudhwa Tiger Reserve in Uttar Pradesh to establish a new viable breeding population. The present study was carried out as a part of management term paper exercise in Dudhwa Tiger Reserve from 5th to 15th March 2002. The methodology adopted for the study was rapid survey of the area, interaction with the various levels of field functionaries and the secondary information from the literature and records. During the study, it was found that the newly established population in Dudhwa seems to be doing well as the population rose to twenty from the seed population of seven. All these seventeen population are progenies single dominant male as others are unable to assist in the presence of this dominant male. Intra-specific aggression amongst the male in the fenced area is another problem. Considering the fact that there is increase in population over the years, the problem of inbreeding depression and hostility amongst the male are inevitable. Fortunately, the authorities in the department could foresee this serious problem well in time and another area, Badhi- Churaila sector in Belriyen Range of the Park has been identified where few females and the males from the existing area will be translocated. It is also proposed few more females and male rhinos of different genetic base will also be released in this newly created area. Thus, the problem of inbreeding depression and male hostility will be solved.

To make the rhino re-introduction programme successful, various Action Plans have been proposed. The urgent requirement is the awareness coupled with effective protection measures and adequate funding mechanism for the infrastructure development. Last but not the least, if our wild life habitat are not to assume the character of "open air museum", the rhino habitats should be extended gradually and endeavor has already been made by the conception of Corridor Restoration and Development Project and Terai Arc Landscape Programme which will not only help rhinos and tigers but the entire terai ecosystem.

Conservation of
the rhino
Bansari

**EXAMINE THE POSSIBILITIES OF ENLARGING/MANAGING LARGER
HABITATS FOR THE RE-INTRODUCED RHINO WITH PLANNING
DETAILS IN DUDHWA TIGER RESERVE**

INTRODUCTON

In ancient times, all the three Asiatic species of rhinoceros roamed over the Indus, Gangetic and Brahmaputra flood plains of the Indian sub- continent. In the relics of Mohenjo-Daro era, some rhino seals were found which are preserved in the Indian National Museum, New Delhi. The records say that the invading Emperor Timur hunted and killed many rhinos on the frontier of Kashmir in AD 1398 and there are evidences that rhino existed in parts of the west of subcontinent as far northwest as Peshawar till 16th Century. Babur, the founder of Mughal Empire in India in his famous memoirs – the Baburnamah described how he hunted rhino in bush country near the Indus as late as 1519 AD

Out of three species of rhino that roamed over the Indo-Gangetic and Brahmaputra floodplains, two species namely Javan Rhinoceros (*Rhinoceros sondaicus*) which was once "fairly common" in the Sundarbans became extinct in India about 1900 AD and Sumatran Rhino (*Didermoceus sumatrensis*) disappeared from the Lushai hills of Assam in about 1935.

The only species of Asiatic Rhinoceros that exists in Indian subcontinent is the Great Indian One-Horned Rhinoceros (*Rhinoceros unicornis*). The *Rhinoceros unicornis* were once widely distributed throughout Indo-Gangetic and Brahmaputra floodplains of the subcontinent.

The causes of disappearance of Great Indian One Horned Rhinoceros from several locations of its former range of distribution and decline of population were primarily the following:

- Destruction and fragmentation of Rhino habitat primarily for extension of agriculture
- Hunting of Rhino for sports during Mughal period and early days of British Rule in India
- Poaching of Rhino for horns and other parts attributed to have magical medicinal values



It will be interesting to note that in Assam Col. Pollock a Military Engineer engaged in laying of roads in Brahmaputra Valley almost shot a Rhino or a Wild Buffalo for breakfast every day. A sportsman in Bengal Dooars, possibly an English Planter fires

about 100 shots at a number of Rhinoceros in a day, killing five and seriously wounding more than twenty five. Maharaja Nirpendra Narayan of Coochbehar shot 208 Rhinoceros between 1871 to 1907.

The Great Indian One Horned Rhinoceros also would undoubtedly have ceased to exist, but for the strict protection given to it when its population fell to a very low level at the beginning of this century. Even though there is no precise documentation regarding the population of Rhino that existed in India at the turn of the century, its population was believed to be around 100 in the beginning of the current century. Being roughly 50 to 60 in Assam and 40-50 in West Bengal. At present the Great Indian One Horned Rhinoceros has total population of about 2500 animals in the world, that too only in India and Nepal. In India, currently its population is restricted to four natural population in Assam viz Kaziranga, Manas, Orang and Pobitora, two natural population in West Bengal viz Jaldapara and Gorumara and one re-introduced population in Dudhwa NP and one migratory population in Katerniaghat in Uttar Pradesh. There are few Rhino exist in Bhutan adjacent to Manas Tiger Reserve, Assam.

In Nepal, the three rhino populations are in Royal Chitwan NP, Royal Bardia N.P and Sukhiaphanta WLS. The Rhino of Royal Chitwan N.P. is a natural population while Royal Bardia NP and Sukhiaphanta WLS has the re-introduced population. The Kaziranga National Park in Assam (India) has the highest population of Rhino (about 1600) and Pobitora WLS (85 rhinos in 18 sqkm area) followed by Royal Chitwan N.P. in Nepal (about 800 rhinos).

In 1979, the Asian Rhino Specialist Group of IUCN Species Survival Commission emphasized the need for continuous efforts in protection and monitoring of the species and "to establish additional viable population in suitable areas, preferably in the former distributional range of the rhino". Thus, on the basis of this logic the IUCN Rhino Specialist Group and the Rhino sub-committee of the Indian Board of Wildlife (IBWL) recommended the establishment of additional Rhino population in India and the Rhinos were re-introduced in Dudhwa Tiger Reserve, U.P. in the year 1984.

Table 1: Indian Rhino Population in India and Nepal (1999)

Country	State	PA	Estimated Population
India	Assam	• Kaziranga NP	1649
		• Manas WLS	5
		• Orang WLS	46
		• Pobitara WLS	76
	West Bengal	• Jaldapara WLS	53
		• Garumara NP	19
U.P.	• Dudhwa TR	13	
	• Katerniaghat WLS	4	
Nepal		• Royal Bardia NP	52
		• Royal Chitwan NP	600
Total			2524

Source: Report on the Regional Meeting for India and Nepal of the IUCN/SSC Asian Rhino specialist Group, Kaziranga, Assam, India, 21-27 February 1999.

BACKGROUND OF RHINO RE-INTRODUCTION PROGRAMME IN DUDHWA TIGER RESERVE, UTTAR PRADESH

The Need for Rhino Re-introduction

1. **Restricted Location:** Conservationists generally agree that an endangered species whose habitat type has by human impact been reduced and split up into separated "islands", should live in as many of those "islands" as possible.



This reduces the risk of extinction. The same applies to Great Indian Rhinoceros and it became imperative to create as many "islands" as possible especially in its former ranges of distribution (*Schenkel 1981*). Re-introduction is necessary mainly to establish new viable breeding populations and safeguard this species from poaching and natural calamities (*Mishra and Dinerstein, 1987*).

2. **Increasing Population of Rhinos in Assam leading to conflict:** Due to devoted efforts to protect Rhinos in Kaziranga National Park, Pobitora WLS and other Protect Areas in Assam, the population pressure is high and during recent years population surplus has been observed. These animals have spread up and down and across the Brahmaputra valley. However, most areas into which the Rhinos move are used for agriculture (*Schenkel, 1981*). The Kaziranga has unusually high density of population and Pobitora has about 85 rhinos in 16 sq. km. area. From the point of view of conservation,

only alternative is to translocate them to other Protected Areas where it can thrive well.

- 3. Protection from Natural Calamities:** In spite of the protective measures, the persecution of this animal continues due to exceptionally high price of its horn. In Kaziranga National Park during the period between 1983-1989, 235 animals were killed by poachers and in the same period 368 animals died due to natural causes related to flood, old age and illness (*Dutta, 1991*). In the recent past poachers wiped out the entire population of Laokhowa wild sanctuary. Despite the extremely strong protection measures, rhinos are also poached from other protected areas as well.

Natural calamities like **flood** is also equally disastrous for Rhinoceros population in Kaziranga cattle grazing are an other problem for rhino. Cattle presence in wild increases the chances of disease transmission. However, there has not been such record of disease transmission (*Sinha, 1999*).

One horned rhinoceros is a hardiest of animal and seldom falls prey to **disease** that affects other herbivores, domestic or wild. But any epidemic outbreak can cause great loss to the animal. In 1944 and 1947 due to suspected case of Anthrax and another unidentified disease several Rhinos died in Kaziranga National Park (*Dutta, 1991*). In 1979 too, hemorrhage septicemia was detected in Kaziranga National Park in 10 cases (*Sinha, 1999*).

- 4. Increasing the Rhino Habitat in as many "islands" as possible:** Habitat destruction is another major problem for rhino. The prime rhino habitats are under various threats from encroachments, cattle grazing, agriculture activities, illegal felling and flood. It had been cleared in past for agricultural purposes and developmental activities. The whole area of terai and flood plains of Brahmaputra is very fertile belt. Encroachment of forestland for agricultural activity is a wide spread problem more or less everywhere in India. Increasing the rhino habitat with strong protection is the only alternative.

Thus, by considering the current highly restricted distribution with poaching pressure, habitat specificity, and in consideration to the scattered small population, it became imperative to re-introduce the species in suitable habitats in its former range of distribution as one of the measures to be adopted for the long-term survival of the Indian Rhinoceros. The IUCN Rhino Sub-Committee of the Indian Board of Wildlife (IBWL) recommended the establishment of an additional Rhino population in India and consequently rhinos were re-introduced in Dudhwa National Park in 1984.

Selection of Dudhwa Tiger Reserve for Rhino Re-introduction

Following up on the recommendation of the Asian Rhino specialist group, the wildlife status evaluation committee of the Indian Board of Wildlife appointed a sub committee to consider alternative areas for establishing a rhino population by translocation in suitable habitats. This sub committee considered the ecological requirements of potential areas for the re-introduction of Indian rhinoceros and established the following criteria:

- Diversity of Habitat, including flooded grasslands with a variety of food plants
- Ample shade and water bodies for wallowing and drinking especially in the hot season.
- Protection from all forms of human disturbance and harassment, including pollution, poaching and the introduction of disease via domestic stock. It is equally important that conflict with cultivation adjacent to areas of re-introduction be avoided, especially in view of rhino's liking for crops such as paddy and sugarcane.
- Translocation to an area which are not yet inhabited by rhinos, but the area falls under the former range of rhino distribution in the past.



On the basis of above criteria, possible alternative habitat suggested were Dudhwa National Park (U.P.), Jaldapara (W.B.), Champaran (Bihar), Intaki in Nagaland and Laligabari Sanctuary of Arunachal Pradesh. Among these various areas considered by this Sub-Committee, Dudhwa National Park was thought to be the most promising which met all the above-mentioned criteria.

- Dudhwa National Park was found to be the most suitable because of the significant similarities to habitats of Kaziranga National Park. Dudhwa National Park contains diversity of habitat, including flooded grasslands with a variety of food plants and ample shade and water for wallowing and drinking.
- Adequate protection is available in Dudhwa Tiger Reserve because of its status National Park and later as Tiger Reserve.
- The area is a portion of the historic range of the rhino. The last one having shot in 1878 in Pilibhit district, which is in proximity to now Dudhwa Tiger Reserve.

Prof. Schenkel confirmed the suitability; the then chairperson of Asian Rhino Specialist Group and Prof. Schenkel observed, "*Dudhwa is the area most suitable for establishing a new local population of Indian rhinoceros. The area is protected, large enough and contain suitable habitat*" and this was further supported by the vegetation survey conducted by the Botanical Survey of India that revealed the presence of several food species of Rhino that were common to Dudhwa, Kaziranga and Manas National Park. These are:

Grasses	10 species
Herbs and climbers	6 species
Hydrophytes	6 species
Shrubs and under shrubs	7 species
Trees	12 species

Rhinos generally prefer **grasses** like *Saccharum spp.*, *Cynodon dactylon*, *Arundo donax*, *Polytoca digitata*, *Hygroryza aristata*, *Vetiveria zizanioides*, *Imperata cylindrica*, *Themeda spp.*, *Chrysopogen aciculatus*, *Puspalidum flavilum*, *Naranga porphyrocoma*, *Phyragmites karka*, etc.

The preferred **sedges** like *Cyperus spp.* and herbs, shrubs and saplings of species like *Polygonum plebeium*, *Ageratum conyzoides*, *Erigeron spp.*, *Artemisia nilagirica*, *Eupatorium odoratum*, *Solanun spp.*, *Colebrookia oppositifolia*, *Murraya koenigii*, *Trewia nudiflora*, *Litsaea spp.*, *Premna sp.*, etc.

They also prefer **aquatic plants** like *Hydrilla verticillata*, *Vallisneria spiralis*, *Hygroryza aristata*, *Potamogeton sp.*, etc.. It is estimated that the rhino population in Kaziranga takes about 77% grasses and 23% herbs and shrubs. Wide range of materials eaten by rhinos suggests that the animal is not very specific in its choice. However, majority of above mentioned food plants are available in the Dudhwa National Park (Hazra and Shukla, 1982).

The U.P. Forest Department came forward with an area of approximately 90km² in the south west part of the park (South Sonaripur and Bellarian ranges) It was estimated an area of 90 sq. km. could support an ultimate maximum number of 90 rhinos considering one rhino per sq. km. (Sale and Singh, 1987). In addition there are other areas of good rhino habitat in Dudhwa Tiger Reserve. In order to prevent released rhinos from wandering out of Dudhwa into adjacent cultivation and to assist their initial establishment in optimal habitat, a 27 sq. km. area was enclosed by a stranded electric fence. A 9 km critical section near the park boundary was additionally protected against accidental escape by construction of rhino and elephant proof trench outside the electric fence. Holding stockades for the new arrivals were built within the fences zone, and Dudhwa field staff was sent to Assam for training in rhino management.

carrying capacity

Finally, re-introduction of Rhino in Dudhwa took place in two phases – one in 1984 and another in 1985. The Rhino Re-introduction in India could become possible due to the keen interest of the former **Prime Minister of India, Smt. Indira Gandhi** and her full support to the rhino re-introduction programme.

The 1984 translocation from Assam

Early in 1984 a group of about 10 rhinos living outside Pobitora Wild Life Sanctuary in Assam was selected by the Assam Forest Department for the capture operation. The fact that these rhinos were causing considerable damage to cultivated crops and

proving difficult to protect adequately provided enhanced justification for the translocation experiment. This remains a mystery that why the translocation of rhinos were done from Pabitora WLS because one of the main reasons of translocation and creation of another viable breeding population was to reduce the high density of rhino from Kaziranga National Park and to prevent rhino population from natural calamities. Between 11 and 21 March 1984, six animals were captured by drug immobilization. Anyway, these six animals were crated, revived, transported to stockades a few kilometers from the capture area and released. After release animals were encouraged to wallow and in most cases satisfactory feeding was established within two to three days. A team of veterinarians rendered necessary health care, mostly consisting of treatment of superficial lacerations received during the capture. The first animal captured, large male, escaped from its stockade during the night.

On 30th March, the five remaining animals (a sub-adult, two elderly females, a young adult and one older male) were crated, driven in trucks to Guwahati Airport and headed into Aeroflot 1176 cargo aircraft chartered by the Government of India. Three of the animals were lightly sedated prior to loading and they all remained calm during the two and a half hour flight from Guwahati to New Delhi. After food and water at Delhi Airport, the rhinos were driven through the night to Dudhwa National Park, where they were uncrated into individual stockades. One female died due to stressful abortion after 11 days but the remaining four settled well; three were released from the stockades on 20 April 1984 and the large male was released on 9 May, after being fitted with radio collar. Another female died on July 31 1984 after a bid to tranquilize her to treat a wound. Thus, there left only three rhinos, one female and two males

The 1985 translocation from Nepal

To establish a rigorous breeding nucleus of rhinos in Dudhwa, it was decided to introduce more stock from a different population. The collaboration of the Majesty's Government of Nepal was obtained in the exchange of four adult female rhinos, from the Royal Chitwan National Park, for 16 domesticated Indian elephants. By selecting only females, the reproductive potential in Dudhwa would be more than doubled and eventual mating of these animals with the totally unrelated Assam males would

ensure maximum genetic vigor. Capture took place on 28-31 March 1985. All four female rhinos, estimated to be between 5 and 7 years old, were immobilized and sieged into crates, which they were revived. They were immediately driven 720 km to Dudhwa and all withstood the 24-hour journey. They were released into the wild after a week.

Present status – A success story

Of the total of nine rhinos translocated to Dudhwa Tiger Reserve seven survived in excellent health and these consisted of the young female and both the males of 1984 translocation from Assam, and all four young females of the 1985 from Nepal. Thus, these seven rhinos constituted the seed population of rhinos at Dudhwa National Park. In 1988, one adult male from Assam died after fought with another dominant male. Again in 1991, a female, from Nepal died due internal infection and abortion. She also lost her male calf in 1993 killed by the dominating male.

The first evidence of breeding in the re-introduced population was detected in the form of remains of a newly born calf in a patch of tall grass in August 1987. There were no signs of predation; hence it may have been a case of premature birth or any such natural circumstances. The first successful calving occurred in early 1989. Three more calves followed this in the same year. The breeding success followed then and the seed population of 6 has since increased to 20 till date (*Sinha et al., 2001*).

Table 2: Demographic Status of existing Rhinoceros in Dudhwa TR (2002)

	SEX		
	Male	Female	Calf
AGE	1. 25 years 2. 11 years 3. 10 years 4. 4 years	1. 23 years 2. 22 years 3. 21 years 4. 19 years 5. 13 years 6. 13 years 7. 10 years 8. 8 years 9. 5 years 10. 5 years 11. 3 years	1. 1 year 2. 1 year 3. 6 month 4. 2 month 5. 2 month
TOTAL	4	11	5

All the rhinos are kept in an area of 27 sq.km. Area encircled by a power fence in south Sonaripur Range. Everyday, four riding elephants are used for monitoring. During rainy season, boats in the Suheli River do monitoring the fence in the southern part of the RRA. There are two boats for this purpose. Seldom were all rhinos sighted everyday due to poor sighting in the tall grassland condition. All the adult rhinos are identified individually by recording different physical traits. The fortnightly report is submitted to the Deputy Director of the park and then forwarded upto the CWLW. Monitoring is done in relation to their movements, feeding and social behaviour. In addition to these, the special protection in this Rhino Re-introduction Area (RRA) is given top priority. The all round increase in the status of protection of the area has helped other endangered wildlife such as *Hispid Hare* and *Bengal Florican*. The herbivore population especially Swamp Deer has increased tremendously in the RRA.

87 11.02
7 00

OBJECTIVES

- To examine the possibilities of enlarging the habitats for the re-introduced rhinos,
- To identify the potential rhino habitats for enlarging the habitats gradually,
- To identify the strengths and weaknesses in managing these larger rhino habitats.

METHODOLOGY

- A rapid survey was conducted in the Dudhwa Tiger Reserve to collect information on present distribution of rhinos, identification of potential habitats for enlarging the rhino habitat in the near future, land use pattern, ground verification of vegetation and major threats.
- Interaction with the various levels of forest staff with a view to collect information related to current situation in RRA, problems and issues, threats etc. in potential rhino habitat were carried out.
- Various secondary information related to rhinos were collected from literature and forest records.

STUDY AREA

The Dudhwa National Park declared as Dudhwa Tiger Reserve under Project Tiger in 1987 in the Kheri district of Uttar Pradesh and lies between 28 18 North latitude and 80 57 E longitude, approximately 30 km south of the Nepal Himalayas. It is 490 sq km in extent with buffer zone of 124 sq km under the park administration.

The park is a compact block of approximately 50 km in length by 10 km in width. The Mohana and Suheli Rivers constituted respectively the northern and the southern – boundaries. An area of 25 sq km within Kakraha block surrounded by a power fence constitutes the Rhino Re-introduction Area (RRA). A section of the fence perimeter has a parallel stretch of trench. The RRA habitat is a mix of tall wet grassland, woodland complex with ten perennial swamps. South of the RRA flows the river Suheli.

The area identified for creation of a satellite population falls in the Bhadi- Churaila sector of the Reserve, which fulfills all habitat requirements for the Rhino rehabilitation. The area has already been surveyed for its habitat suitability and found fit for rehabilitation of Rhinos. The extent of area to be fenced in the Bhadi- Churaila sector has been decided on the basis of estimated number of animals in the population to be built in due course of time and habitat requirement of Rhinos for different purposes. This area comprises of Sal forest (336 ha), Grassland (608.23 ha) and Wetland/ Swamp (129.00 ha), which fulfill the need of rhino, and is one the prime rhino area in Dudhwa NP.

As whole vegetation consist of some of the best Forest of Sal (*Shorea robusta*) in India, mixed moist forest, riparian communities, tall wet grassland with patches of short grasses. Interspersed within the grassland are a number of swamps within the park, grassland covers 20% of the total area. So far, 75 species of trees, 21 species of shrubs, 17 species of climbers, 77 species of grass and grassland plants, 179 species of aquatic plant have been listed.

DISCUSSION

Need for Enlarging Rhino Habitat

1. Augmenting Genetic Variability:

The original objective was aimed at releasing 30 rhinos in Dudhwa National Park. The precise reason why this was not carried out after release of two batches is both financial and administrative. However, due to breeding the number has increased from seven to seventeen. Out of the seven seed population, only two were male named Raju and Bankey. Initially, Raju had asserted his dominance but with passage of time Bankey became dominant and killed Raju in one of their regular fights in October 1988. During 1991 an attempt was made to counter the anticipated problem of inbreeding by introducing a male, named Rohit, in captivity from Kanpur Zoo. He was repeatedly attacked by Bankey and severely injured. This animal had to be taken out, treated and returned to the zoo after it had recovered. The existing rhino population in RRA has two more adult males but they are unable to assert themselves in front of the dominant male Bankey. Bankey continues to be the dominant male even today. All the calves born till today are progeny of this dominant male – Bankey. Because all these calves surviving now are born after 1989, there is little chance of mating with Raju who died in 1988. In absence of any other male even the sexually mature daughters are mating with their father. The two adult males, though unable to assert are also the progenies of same rhino Bankey and if in the future they will mate with the females, same type of genes will be transmitted to the next generation. If this situation prevails for few successive generations, it may cause inbreeding depression, which is a threat for genetic viability. So, there is an urgent need to introduce some other male with different genetic base. And keeping in view the past experiences with Bankey's behaviour newly translocated male rhinos should be kept in separate enclosure to avoid the chance fighting among them. And hence, there is an urgent need of enlarging rhino habitat (Sinha, 1999).

2. To Reduce conflict amongst males in existing RRA:

Bankey has driven one of the two adult males in existing population out from the main fence on several occasions. Therefore, separate contiguous minifence has been created to provide safe heaven to one of these animals. The two males are often seen parading on their respective sides of the common section of the fence. The operation failures of the fence and even functional fence also are unable to restrict the rhinos from fighting. Thus, there is an urgent need to create another rhino habitat (Sinha, 1999).

3. Increasing Population in the existing RRA:

Increasing population of rhino in the existing area is also a major problem. In this case, either new area can be selected for rhino translocation in Dudhwa Tiger Reserve or the enclosure will be removed. In case of the removal of the enclosures, there are chances of crop depredation by rhinos in the adjoining areas, which is very close to the southern boundary of the Rhino Re-introduction Area (RRA). In response to these situation, it is better to keep the rhino under enclosure for few more years, at least till the people accept them as their neighbors. And hence, there is a need to select some new area inside the Dudhwa National Park. In the light of above facts, long back in 1991, Dr. S.P. Sinha, Project Coordinator, Corridor Project of USWLS, strongly recommended for the urgency of creating another viable breeding population of rhino in Dudhwa Tiger Reserve. He proposed Bhadi-Churaila area in Belreiyen Range of Dudhwa Tiger Reserve. Dr. Sinha recommendation and proposal has been accepted and Bhadi-Churaila sector of Dudhwa National Park has been identified for creation of additional rhino habitat as this fulfills all habitat requirements for the rhino rehabilitation.

The area selected for enlarging Rhino habitat

On the basis of recommendation of Dr. S.P. Sinha in 1991, the area is identified for creation of another viable population of rhino in Dudhwa Tiger reserve. Finally, in 2002, the work is in progress and proposed area is to be enclosed in power fence for the creation of another viable population of rhino inside the Dudhwa Tiger Reserve. The area falls in Bhadi-Churaila sector of Belreiyen Range. This area provides water for drinking and wallowing, shade and an adequate variety of plants known to be

eaten by rhino elsewhere. One of the essential prerequisite for rhino re-introduction is vast grassland with water for drinking and wallowing. This Bhadi-Churaila sector has two permanent water bodies known as Bhadi Taal and Churaila Taal respectively. Apart from these two permanent and large water bodies, there are various other smaller water bodies, which are permanent as well as, seasonal. The Rhino Subcommittee of Indian Board for Wildlife (IBWL) has identified this Bhadi-Churaila sector in Belreiyar Range as one of the possible sites for re-introduction of rhino in their original recommendation. Feasibility study with regard to the habitat availability for the re-introduction of rhino in Dudhwa by the Botanical Survey of India led by Dr. Hajra carried out a detailed survey of the vegetation of Dudhwa in relation to the rhino feeding ecology. The detailed study clearly established a number of floral elements common to Dudhwa National Park (U.P.) and Kaziranga (Assam), both of which are excellent rhino habitats.

The advantage of the area selected is that it is in the central location of Belreiyar Range. There is no danger of rhino wandering in the human occupation and cultivation in case of operation failure of power fence. While in case of existing rhino re-introduced area, it is adjacent to the southern boundary of the park, which lacks a buffer zone and outside of which is an area of dense human occupation and cultivation. Bhadi-Churaila sector is centrally located and sufficient buffer is available.

This Bhadi-Churaila sector comprises an area of 10.74 Sq.km. and the fencing work has already been started and target was to complete the fencing operation by the end of this financial year. The work was in progress when the study was carried out.

Table 3: The Bhadi-Churaila Habitat

Range	Block/Compartment	Sal Forest (ha.)	Grassland (ha.)	Wetland/Swamps (ha.)
Belreiya Range	Bhadi-2	180.09	-	-
	Bhadi-3a (Part)	-	30.00	105.00
	Bhadi-3C	18.21	354.88	105.00
	Bhadi-6a	88.63	-	-
	Bhadi-6b	-	106.43	-
	Laudaria-2 (Part)	35.39	-	-
	Laudaria - 3B	4.04	96.62	24.00
	Laudaria - 4 (Part)	9.72	20.30	-

(Source: Proposed Plan for Bhadhital Rhino Area, 2002)

Total Area = 1,073.31 ha = 10.733 sq. km

Total length of the fence = 17 km

THE OTHER POSSIBLE SITES FOR ENLARGING IN THE FUTURE:

There are excellent "phantas" (grassland) and water bodies in other ranges as well. Dudhwa and Sathiana Ranges are blessed with vast expanse of grasslands with number of viable swamps. These areas can also be rehabilitated with rhinos in due course of time when the rhino population increases. There exists a viable connectivity between the existing rhino and other grasslands. All these can be joined together in due course of time with the increase in rhino population and the fenced areas be extended in the boundaries of the park only.

PROPOSED ACTION PLAN

1. **Fencing the newly identified RRA:** As per the plans, the fencing of newly identified RRA i.e. Bhadi-Churaila sector of RRA will be completed by the end of the financial year 2001-2002 i.e. by end of March 2002. Work is in progress when the study was carried out. The erection of fencing posts were in progress and about 3000 posts were required for this purpose. The three-strand power fence will cover entire proposed area of 17 km long stretch of perimeter before releasing the rhinos. 3-strand
2. **Releasing rhinos in the newly created RRA from the existing RRA:** After the completion of power fencing, three females and one male from the existing RRA will be released in this newly created Bhadi-Churaila sector of RRA. This will provide immediate relief to the conflict situation in the existing RRA and enhance breeding in the newly created RRA.
3. **Releasing fresh batch of rhinos to augment the genetic variability:** Since all the rhinos born in the existing re-introduction area are progenies of single bull Bankey, the problem of inbreeding depression needs to be addressed timely. In order to overcome this problem, there is a proposal to introduce fresh batch of rhinos – one male and three females, either from Royal Bardia NP in Nepal or Assam to broaden the genetic base (*Sinha, Mishra and Tiwari, 2002*)
4. **Monitoring:** Presently four elephants are engaged in the monitoring work. Mahouts and staff scan the area and send the report to the headquarter and office of the Deputy Director. Monitoring is an important activity to see their health condition, movement pattern and other behavioral aspects. With the creation of new RRA, more number of elephants and Mahouts as well as monitoring staff is required. 4 elephants
5. **Infrastructure support:**
 - A road will be built along the inner side of the entire fence to familiarize patrolling and fire management
 - The monitoring parties are provided with walkie-talkie sets so that they are always in contact while conducting monitoring.

6. **Habitat Management:** This is one of most important aspect of the proposed action plan. It has several components:-
- (i) **Grassland Management:** For the management of grassland, annual burning is one of the most important exercise. This discourages the invasion of woodland and facilitates new shoots there by increasing nutritional value. But the annual burning is to be made in consideration of the other endangered species like Bengal Florican, Hispid Hare, prey species of tiger and sensitivity to breeding requirements of species that are obligate of the grasslands. For example, the grasslands utilized by Bengal Florican, the grass is cut first prior to burning. Burning is done in patches while the areas used by the hispid hare are protected from the fire (Sinha, Sawarkar and Tiwan, 2001). Controlled and cool burning plays an important role in management of grasslands especially relating to timing and pattern of prescribed burning.
 - (ii) **Wetland management:** Water is the lifeline of rhino though here are permanent swamps in the newly created RRA, creation of additional wallowing grounds and waterholes as well as supplementing in the existing taals (water bodies), new borings and pumping sets are required.
 - (iii) **Weed Removal:** Removal of weeds from the "phantas" (grasslands) and water bodies also is an important aspect of habitat management.
 - (iv) **Woodland manipulation:** The newly created RRA contains predominantly of Sal trees and other species as well. Though felling and removal of tree from PA is totally banned, the problem needs to be addressed to check the invasion of trees into grasslands. The removal of trees is required from the point of view of extension of rhino habitat.
7. **Research Work:** With the passage of time, more and more research is required in relation to its habitat, food availability intraspecific behaviour and health aspects.
8. **Veterinary Care:** In Dudhwa NP, at present there are working elephants for monitoring purpose as well as for tourists. These working elephants need proper care and upkeep. Veterinary care also needed from time to time for rhinos. So, establishment of veterinary units is required.

9. **Local Awareness:** Villages located at the fringes are already facing the problem of elephants and other herbivores straying into their agricultural field. Re-introduction of rhino is another added burden to their problem. Due to proper power fencing and constant monitoring, rhinos generally do not stray into agricultural fields and there is no such report till date from the existing RRA. But with the creation of new RRA, these rhino population is bound to increase. Therefore, to make the conservation programme more successful, participation of fringe villagers is of utmost importance, so that they may compromise with the situation, at least their hostility can be reduced.
10. **Ecotourism:** The last rhino disappeared in this area was around 1878. Re-introduction of rhinos in this area has fascinated the tourists. Now every tourist who visits the National Park wants to have a glimpse of this mighty animal. More elephants and mahouts are required for this purpose.
11. **Eco-development:** The fringe villagers are the worst sufferers in the process of building up the PAs. They are deprived of the forest produces required by them for their day-to-day livelihood. In addition, they become the victim of wildlife deprecations including loss of life. Thus, the people have to face severe economic hardships. To ameliorate the economic hardships of these people, economic support in the form of eco-development programme has become very essential.
12. **Anti-poaching measure:** Protection is the *sine qua non* for any PA. Some conservationists give importance to the protection to the extent that PA doesn't require any input except protection. This is true to the great extent. The success story of increased rhino population in Kaziranga National Park, Assam, Royal Chitwan NP, Nepal and other rhino bearing PAs is primarily due to effective anti-poaching network built up in these areas. Dudhwa Tiger Reserve shares 56 kms of international boundary with Nepal. With the re-introduction of rhinos, the determined poachers may be attracted in their venture as the rhino horns fetch good revenue in the international market. Moreover, Tharu villages located in the Northern fringe of the park have relation in Nepal. For them, it can be a new and wonderful avenue for getting fast and handsome revenue. The facilities provided presently are not at all adequate. Sophisticated arms and training of anti-poaching personnel is of utmost importance to make the Rhino Re-introduction Programme successful.

13. **Setting up of field Head Quarter in the RRA:** There is a requirement of setting up of head quarter, possibly near chhanga nala area for better monitoring of newly re-introduced rhinos. This will also require appropriate fund.
14. **Construction of watchtowers:** Watch towers need to construct in this new area for monitoring activities.
15. **Requirement of more elephants:** More elephants are required for monitoring the rhinos in the newly re-introduced rhino area for monitoring purpose as well as for the tourists. At present the elephants are less in number and to fulfill the requirements of tourists and monitoring activities they are put in to overworking which badly affects their health.
16. **Financial Aspect:** All the above mentioned points in Proposed Action Plan will be light of the day only if adequate financial provision is available. Initially, the Government of India has provided financial support for the translocation and conservation of the rhino in newly reintroduced area. But with the passage of time, no funding is available from the Central Government for the conservation of rhino in Dudhwa. At present, the entire effort for conservation of rhino in Dudhwa is being born by the State Government as a part of the Forestry budget. Thus, the financial input is obviously very small. With the increase of rhino population in this protected area i.e. Dudhwa National Park and gradual change of management strategies, the requirement of financial support is very essential. Dr. S.P. Sinha, Project Coordinator, Corridor Restoration and Development Project of USWLS approached various organizations for the financial support with a proposal of Rs 65 Lakh for ten years. Dr. Sinha, approached International Rhino Foundation, USA, SOS Rhino, USA, US Fish & Wildlife Service, ITC-India, and Weeden Foundation New York, USA for sponsorship. Memorandum of Understanding is on the process and talks are going on some of these organizations have responded for the cause of conservation of rhino. The Government always welcomes such type of help.

THE EXPECTED OUTCOME

1. The creation of the Bhadi-Churaila sector of RRA shall reduce the prevailing conflict situation among the males in the existing population. Fresh batch of rhinos either from Nepal or Wild captive rhinos will also be reintroduced in this area to broaden the genetic base.
2. The intensive management efforts to be applied for Rhinos are bound to benefit the existing take off population of the highly endangered swamp deer in the area. These expectations are based on the observations recorded in the existing RRA. The existing RRA is located on the south Sonaripur Range and this Range has recorded the highest number of all the five deer's found including swamp deer and hog deer amongst the nine ranges of Dudhwa Tiger Reserve. (UP Forest Department, 2001)
3. There exists a viable connectivity between the existing and newly created RRAs through the grasslands of Bankey Tal area. Thus two populations can also be joined in due course of time when the Rhino population increases by extending the fenced areas.
4. Last but not the least, the conservationists' ultimate objective of creating as many separated viable "islands" as possible to reduce the risk of extinction will be fulfilled when the re-introduced rhinos of Dudhwa once again haunts the entire Terai area.

CONCLUSION

The re-introduction of rhinos in Dudhwa NP has resulted in the first viable population of rhino in these Terai areas of U.P. since the last century. This was the first experiment of re-introduction of Asian rhinos in the world and it is a success story now. Following India's footsteps the Government of Nepal also has re-introduced rhinos in Royal Bardia National Park from Royal Chitwan National Park. Both these re-introduction programmes proved highly successful and there is every hope that it will undoubtedly lead to further use of this approach to repopulate the selected areas of the rhino's former ranges of distribution. Our ultimate endeavor to cover more and more areas of former ranges of rhino distribution, if we do not want the present rhino habitat to assume the character of "open air museum" where endangered species are kept under rigorous protection. With the concept of landscape level of habitat management as the Restoration and Development of Corridor Project and Terai Arc Landscape Programme. There is hope that posterity will appreciate our effort and this mighty animal will once again roam around freely in his ancestor's home that is encroached by man. The areas need to be extended and provided with more corridor types outlets so that animals can roam around in different areas freely.

Last but not the least, as long as the myth of the aphrodisiac and medicinal value of the rhino horn persists, the animal will never be safe from poaching, which has become a lucrative business. Strong protection and mass awareness seems to be the only and the best alternative.

BIBLIOGRAPHY

- A. Kumar, 1996.** Forest type mapping of Dudwa National Park using remote sensing and GIS. P. G. Diploma in Forest Ecology. IIRS, Dehradun.
- Altken, B. 1988.** A Dudhwa diary. Cub Vol.5(4):18-22.
- Anonymous.** Some observations on habitat of the Dudhwa national park. 1-7
- Bakhsh, N. 1995.** Checklist of Birds of Dudwa National Park. Unpublished.
- Berkmuller, K. and Mukherjee, S. 1987.** Buffer zones in the service of ecodevelopment. Symposium on tropical Ecology. December 1987: 1-18
- Chaturvedi, A.N. and Mishra, C.M. 1985.** Ecological survey of grasslands at Dudhwa national Park. Indian Foresters. 111 (8): 579-582
- Coe, Malcom. 1979.** A look at wildlife management in India. Oryx 15(1) : 60-65
- Hajra, P.K. and Shukla, U. 1982.** Dudhwa National Park. Botanical Survey of India. (Some botanical aspects of the proposed new habitat for rhino). Howrah. April 1982:1-12 (1): 41-48
- Halloway, C. (1973).** Swamp deer in Uttar Pradesh. Oryx. 12 (1): 41-48
- Jain, S.K. and Sastry, A.R.K. 1983.** Botany of some tiger habitats in India. Botanical Survey of India. 70pp
- Javed, S. 1996.** Elephants in Dudhwa. Garjah 16:17-22.
- Kotwal, P.C. and Parihar, A.S. 1990.** Management of hard ground Barasingha in Kanha national park. Paper presented in National seminar on management of Barasingha in Dudhwa national park, Lakhimpur- Kheri.
- Leith, H. 1975.** Primary production of major vegetation units of the world. 203-215. In: Leith, H & R.H. Whittaker. Eds. Primary productivity of the Biosphere. Springer-Verlag, New York.
- Martin, E and Vigne, L. 1996.** Numbers of Greater one-horned rhino continue to rise. News and views. Oryx 30(3).

Mishra, G.C. 1989. Rhino rehabilitation in Dudhwa National Park. Van Vigyan Vol. 27(1): 60-64

Mishra, C.M. and Chaturvedi, A.N. 1985. Ecological survey of grasslands at Dudhwa National Park. Indian Forester Vol. 111(8): 579-582

Martin, C. (1975). Status and Ecology of the Barasingha (*Cervus duvaucelli branderi*) in Kanha National Park (India). Organ. Kolb. Zwich. 134 pp

Qureshi, Q. and Sawarkar, V.B. 1991. Protected area design and conservation: Dudhwa national park. A case study. National Symposium of Environment and wildlife. Indian Soc. of Environment.

Qureshi, Q., Sawarkar, V.B. and Mathur, P.K. 1990. Status and management of swamp deer in Dudhwa National Park. Seminar on management of Swamp deer in Dudhwa Tiger Reserve. INT ACH and WWF - India.

Qureshi, Q., Sawarkar, V.B. and Mathur, P.K. 1990. Ecology and management of Swamp deer *Cervus devauecell devauecell* in Dudhwa Tiger Reserve. Project report of Wildlife Institute of India.

Qureshi, Q., Sawarkar, V.B. and Mathur, P.K. Dudhwa National Park - The ark must be kept afloat. Unpublished Paper

Raj, T. and Sinha, N.K. 1988. Some of the problems of management of Dudhwa National Park. Cheetal 28(3):41-44

Rahamani, A. 1996. Present status of the Bengal Florican *Houbaropsis bengalensis* in Dudhwa Tiger reserve. A Report.

Rizvi, S.H.A. 1979. Working plan for the South Kheri Forest Division. Vol 1 (part 1 & 2) Rizvi, S.H.A. 1979. Working plan for the South Kheri Forest Division. Vol 2.

Rodger, W.A. 1988. The wild grazing ungulates of India: An ecological review. Proceedings of the second International rangeland congress, New Delhi.

Sale, J.B. and Singh, S. 1987. Reintroduction of Greater Indian rhinoceros into Dudhwa National Park. Oryx Vol. 21(2):81-84

Sale, J.B. 1986. Rhinos re-established in Uttar Pradesh. Indian Forester 112 (10):945-48

- Sale, J.B. 1986.** Reintroduction in Indian wildlife management, Indian Forester, 112(10):867-73
- Sankaran, R. 1988.** The Bengal florican at Dudhwa national Park. The floricans. Bombay Natural History Society, Annual Report No.4:25-32
- Sankaran, R. 1988.** Studies of the Swamp deer in Dudhwa national Park (1988-1989), Endangered species project. Bombay Natural History Society. Technical report No.14
- Sankaran, R. 1989.** Status of the swamp deer in Dudhwa National Park 1988-1989. Bombay Natural History Society. Technical Report No. 14 25
- Sankaran, R. 1990.** The Bengal florican in Dudhwa National Park. In: Status and Ecology of Lesser Bengal Florican with report on Jerdon Courser and Mountain Quail. Bombay Natural History Society. 45-54.
- Sawarkar, V.B. 1988.** Status of the northern Swamp deer *Cervus duvauceli duvauceli* in the Dudhwa National Park. A report Wildlife Institute of India.
- Schaaf, D. and Singh, A. 1977.** Barasingha in the Dudhwa sanctuary. Oryx Vol. 13(5):495-498
- Schaller, G.B. 1967.** The deer and the Tiger. Chicago Um Press, Chicago. 370 pp.
- Schenkel, S. 1981.** Report on the suitability of Dudhwa National Park as potential site for reintroduction of the Indian rhinoceros. 21-24.
- Singh, A. 1978.** The status of the Swamp deer (*Cervus duvauceli duvauceli*) in the Dudhwa National Park. In: Peter Scott Chairman, Threatened Deer. International Union Conservation of Nature & Natural Resource (IUCN) , Morges, Switzerland. 132-142.
- Singh, A.N. and Singh, A. 1985.** A checklist of birds of Dudhwa national Park. A report by the UP Forest Department, Lucknow.
- Singh, L. 1991.** Present status of Barasingha *Cervus duvauceli duvauceli* in India and its conservation. Paper presented at the International Seminar on Veterinary medicine in wild and captive animals. 8-10 November 1991, Bangalore.
- Singh, R.L. 1994.** Dudhwa Rashtriya Udhyan . In : Van Vanyajeev aur Vanvasi. Pub Bishen Singh Mahendra Pal Singh, Dehra Dun.

- Singh, R.L. 1983.** Management Plan of Dudhwa National Park. Uttar Pradesh Forest Department (Unpublished).
- Singh, R.L. 1986.** Conflict in the wild. India Today.
- Singh, V.B. 1978.** The elephants in UP - a resurvey of its status after 10 years. Journal of Bombay natural History Society. Vol. 75(1): 71-82
- Singh, V.P. 1982.** Bio-ecological studies on *Cervus duvauceli duvauceli*, Swamp deer (Barasingha) in Dudhwa near Indo-Nepal border. Ph.D thesis, Kanpur University (unpublished) Singh, V.P. 1987. Dudhwa National Park. Animal Citizen Vol. 26(1):48
- Sinha, S.P. and Sawarkar, V.B. 1994.** Ten years of the Asian Rhino reintroduction programme in Dudhwa national Park, India. IUCN reintroduction Specialist Group Newsletter. No.9
- Sinha, S.P. and Sawarkar, V.B. 1991.** Management of the reintroduced Indian One-homed rhinoceros in Dudhwa National Park.- A report Wildlife Institute of India.
- Sinha, S.P. and Sawarkar, V.B. 1991.** Management of the reintroduced Great one-homed rhinoceros (*Rhinoceros unicornis*) in Dudhwa National Park. UP, India. International Rhino Conference, Rhinoceros Biology and Conservation, San Diego, California, USA:9-11 May 1991.
- Sinha, S.P. and Sawarkar, V.B. 1991.** Habitat and dietary selection by the population of reintroduced Great Indian one homed rhinoceros *Rhinoceros unicornis* in Dudhwa National Park. International Rhino Conference, Rhinoceros Biology and Conservation, San Diego, California, USA: 9-11 May 1991.
- Sinha, S.P. and Sawarkar, V. B. 1992.** Management of the reintroduced Indian Great one- homed rhinoceros (*Rhinoceros unicornis*) in Dudhwa National Park. IV World Congress of National Parks and protected areas, Caracas, Venezuela. 10-21 February 1992.
- Sinha, S.P., Sawarkar, V.B. and Mukherjee, S.K. 1995.** Bibliography of Indian One-homed Rhinoceros (*Rhinoceros unicornis*). Wildlife Institute of India, Dehra dun

Sinha, S.P. and Sawarkar, V.B. 1996. Monitoring and Management of reintroduced rhinoceros in Dudhwa national Park. International Conference on Sustainable Use of Natural resources, Budapest, Hungary. 26-29 August 1996.

Sinha, S.P., Sawarkar, V.B and Ashish Tewari. 2001. Management of Re-Introduced Greater One-horned Rhinoceros (*Rhinoceros unicornis*) in Dudhwa National Park & Tiger Reserve, Uttar Pradesh, India. Proc of the International Elephant and Rhino Research Symposium, Vienna, June 7-11, 2001

Sinha, S.P. 1991. Rhino reintroduction programme in Dudhwa National Park, an update. IUCN reintroduction Specialist Group Newsletter No.(2), Nov 1991.

Sinha, S.P. 1991. Hispid hare *Caprologus hispid* rediscovered in Dudhwa National Park. Zoo Print. 4(4), Nov 21.

Tilak, R. and Ray, P. 1990. Range extension and notes on *Rana erythraea* from Dudhwa National Park, Uttar Pradesh. Journal of Bombay Natural History Society. 87(3): 380-384.

- **1982.** Report of the committee to study the problems of man-eating tigers in Kheri district of Uttar Pradesh. Ministry of Agriculture and Irrigation. Govt Of India. New Delhi.

- **1981.** Final recommendations on the translocation of Great Indian One Horned Rhinoceros, by the rhino sub committee. Wildlife status evaluation of IBWL. Govt of India. New Delhi



THE TIMES OF INDIA, NEW DELHI

A new home for rhinos at Dudhwa sanctuary

By Mohit Dubey
Times News Network

LUCKNOW: Hostility between two male rhinos and fear of vendetta from a calf on an elder — the lone reproducing male rhino at the Dudhwa national park — has jolted the state forest department into action.

The outcome: A sprawling new 20-square km home, dotted by swamps and grasslands, for the 17 Indian Rhinoceros housed here.

Confirming that the second phase of the 'Rhino rehabilitation project' had begun after it was undertaken in 1984, chief wildlife warden R. L. Singh said on Monday that electric fencing of the new area had begun.

Back from a two-day visit of Dudhwa, Singh said that they had decided to "get going" after reports that the hostility among male members was posing a major safety hazard for the regionally extinct race of Indian rhinos. The new area being carved out in the thickets of Dudhwa involves two swamp areas of Bhadi and Churaila Tsal. The second phase entails an expenditure of Rs 15 lakh that involves the setting up of a field headquarters in the area, some watch towers and the construction of the electric fencing.

The department has cleared renovation of a dilapidated and abandoned rest house at *Changa Nala* which would serve as the field headquarters. Dudhwa houses 17 rhinos which is the largest concentration in India except for the Brahmaputra valley in North-East.

The second phase of the "Rhino rehabilitation" project has also been hurried through since the incidence of tiger preying have increased manifold in the area. Fears of inbreeding have also forced foresters hands. Six animals have fallen to the wild cat in the last two decades.

ACKNOWLEDGEMENT

We are highly grateful to Prof. V.B. Sawarkar, Director and Dean, Faculty of Wildlife Sciences at WII, Dehradun from whose inspiration and his encouragement to develop the art of scientific paper writing, as Prof. Sawarkar believes in more meaning and sense in the fewest short words. He has kindly accepted as one of the author of this report.

There are plenty of materials already published on rhino-re-introduction programme in India, which helped us in many ways while comparing the situation and its development.

The field staffs of Dudhwa NP are well experienced with wide range of knowledge in the field of wild animal distribution in the park. Mr. A.B. Tiwari, ROF, and Girdhar, Forester, South Sonaripur (where existing RRA is located) and Mr. Mohammad Azam Khan, ROF and Ambika Prasad, Forester of Belrelyen Range (where newly created RRA is located) imparted with their knowledge and experience and dealt in detail regarding the problems and issues. We unhesitatingly acknowledge that bulk of the material for this has been gleaned from the interactions conducted with these individuals. Departmental records also were consulted. Our thanks to all of them.

Mr. Ashish Tiwari, Deputy Director, Dudhwa NP deserves gratitude for sharing the problems and issues related to RRA.

We are highly indebted to Dr. Bitapi C.Sinha for her guidance and support in bringing out this.

Fellow Officer Trainees also have contributed indirectly in writing this through meaningful discussion at various occasions. Our thanks to them. We are highly thankful to Mr. Kuldeep Chauhan for his cooperation and assistance in bringing this document.

Authors

Authors Address:

***** Prof. V.B.Sawarkar, IFS, Director & Dean, Faculty of
Wildlife Sciences, Wildlife Institute Of India,
Chandrabani, Dehra Dun -248001 Uttaranchal State
INDIA**

Email: omega@wii.gov.in

Fax: 91-135-640117

*** Mr. Satya Prakash*, IFS. Former PG Diploma Wildlife
Management Trainee of Wildlife Institute of India, Forest
HQ Ranchi, Jharkhand Sate, INDIA**

**** Dr.Satya Priya Sinha, Project Coordinator, Corridor
Project Of US Fish & WLS C/O Wildlife Institute of India
Chandrabani Campus HNO.IV, Dehra Dun -248001
Uttararchal State INDIA**

Email: sinhasp@hotmail.com & sinhasp@yahoo.com