

RESEARCH PAPER

Postinduction butorphanol administration alters oxygen consumption to improve blood gases in etorphine-immobilized white rhinoceros

Peter Buss^{a,b,c}, Michele Miller^d, Andrea Fuller^{b,c}, Anna Haw^b, Eliza Stout^e, Francisco Olea-Popelka^e & Leith Meyer^{b,c}

^aVeterinary Wildlife Services, South African National Parks, Kruger National Park, Skukuza, South Africa

^bDepartment of Paraclinical Sciences, Faculty of Veterinary Science, University of Pretoria, Onderstepoort, South Africa

^cBrain Function Research Group, School of Physiology, Faculty of Health Sciences, University of the Witwatersrand, South Africa

^dDepartment of Science and Technology/National Research Foundation Centre of Excellence for Biomedical Tuberculosis Research, Medical Research Council Centre for TB Research, Division of Molecular Biology and Human Genetics, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa

^eColorado State University, College of Veterinary Medicine and Biomedical Science, Department of Clinical Sciences, Fort Collins, CO, USA

Correspondence: Peter Buss, Veterinary Wildlife Services, South African National Parks, Kruger National Park, Private Bag X402, Skukuza 1350, South Africa. E-mail: peter.buss@sanparks.org

Abstract

Objective To investigate the effects of post-induction butorphanol administration in etorphine-immobilized white rhinoceros on respiration and blood gases.

Study design Randomized crossover study.

Animals A group of six sub-adult male white rhinoceros.

Methods Etorphine, or etorphine followed by butorphanol 12 minutes after recumbency, was administered intramuscularly [2.5 mg etorphine, 25 mg butorphanol (1000–1250 kg), or 3.0 mg etorphine, 30 mg butorphanol (1250–1500 kg)]. Sampling started at 10 minutes after initial recumbency, and was repeated at 5 minute intervals for 25 minutes. Arterial blood gases, limb muscle tremors, expired minute ventilation and respiratory frequency were measured at each sampling point. Calculated values included alveolar–arterial oxygen gradient [$P(A-a)_{O_2}$], expected respiratory

minute volume (\dot{V}_E), tidal volume (V_T), oxygen consumption (\dot{V}_{O_2}) and carbon dioxide production (\dot{V}_{CO_2}).

Results Etorphine administration resulted in an initial median (range) hypoxaemia [arterial partial pressure of oxygen 25.0 (23.0–28.0) mmHg], hypercapnia [arterial partial pressure of carbon dioxide 76.2 (67.2–81.2) mmHg], increased $P(A-a)_{O_2}$ [41.7 (36.6–45.1) mmHg, \dot{V}_{O_2} [11.1 (10.0–12.0) L minute⁻¹] and muscle tremors. Butorphanol administration was followed by rapid, although moderate, improvements in arterial partial pressure of oxygen [48.5 (42.0–51.0) mmHg] and arterial partial pressure of carbon dioxide [62.8 (57.9–75.2) mmHg]. In rhinoceros administered butorphanol, \dot{V}_{O_2} [4.4 (3.6–5.1) L minute⁻¹] and \dot{V}_{CO_2} [4.2 (3.8–4.4) L minute⁻¹] were lower than in those not administered butorphanol. Increased arterial oxygen tension was associated with lower oxygen consumption ($p = 0.002$) which was positively associated with lower muscle tremor scores ($p < 0.0001$).

Conclusions and clinical relevance Hypoxaemia and hypercapnia in etorphine-immobilized rhinoceros resulted from an increased $[P(A-a)O_2]$ and increased $\dot{V}O_2$ and $\dot{V}CO_2$ associated with muscle tremors. Rather than being associated with changes in $\dot{V}E$, it appears that improved blood gases following butorphanol administration were a consequence of decreased $\dot{V}O_2$ associated with reduced muscle trembling.

Keywords butorphanol, etorphine, oxygen consumption, white rhinoceros.

Introduction

Chemical capture is an essential tool in the management of free-ranging white rhinoceros (*Ceratotherium simum*) (Wenger et al. 2007). Etorphine, the opioid preferentially used in immobilization of rhinoceros, results in areflexia without total loss of consciousness, resulting from central nervous system depression within a few minutes following intramuscular (IM) administration (Portas 2004; Swan 1993). Etorphine is combined commonly with azaperone, a butyrophenone tranquillizer, which reduces induction times and opioid-associated hypertension (Portas 2004). Unfortunately, hypoxaemia, hypercapnia and acidaemia are well documented adverse effects associated with the use of etorphine and azaperone, and mortalities have been associated with immobilization in rhinoceros (Kock et al. 1995; Haw et al. 2015). Butorphanol, a mixed opioid agonist–antagonist, is administered intravenously (IV) in immobilized rhinoceros to mitigate these adverse cardiorespiratory effects. However, inconsistent changes in blood gases following butorphanol administration have been reported. Different explanations for the effect of butorphanol in white rhinoceros, including improved ventilation or possible alterations in metabolic activity, have been proposed (Miller et al. 2013; Boardman et al. 2014; Haw et al. 2014; Buss et al. 2015). The objectives of this study were to determine the effects of etorphine on respiration in immobilized rhinoceros and changes that occur following IV administration of butorphanol. Cardiovascular effects of etorphine and IV butorphanol in these study rhinoceros have been previously reported (Buss et al. 2016).

Materials and methods

The study had ethical approval from SANParks Animal Use and Care Committee (Ref. no. 14-2) and University of the Witwatersrand Animal Ethics Screening Committee (Ref. no. 2014/15/C). Management of the rhinoceros was conducted according to the South African National Parks (SANParks) Standard Operating Procedures for the Capture, Transportation and Maintenance in Holding Facilities of Wildlife.

A group of six white rhinoceros, subadult (age, 5–6 years) males, were captured in Kruger National Park (23° 49' 60 S, 31° 30' 0 E; alt. 317 m), South Africa, and habituated to captivity in holding pens over a period of 4 months. The study was a crossover design with two interventions allocated using computer generated random numbers with a 2 week washout period between treatments: 1) etorphine hydrochloride (9.8 mg mL⁻¹, M99; Elanco, GT, South Africa) plus hyaluronidase (5000 i.u., Kyron Laboratories, GT, South Africa), administered IM; and 2) etorphine hydrochloride plus hyaluronidase administered IM followed by butorphanol (50 mg mL⁻¹, Kyron Laboratories) administered IV. Doses were 2.5 mg etorphine, 5000 i.u. hyaluronidase, 25 mg butorphanol (1000–1250 kg), and 3.0 mg etorphine, 5000 i.u. hyaluronidase, 30 mg butorphanol (1250–1500 kg) (Haw et al. 2014; Buss et al. 2015). Azaperone was not included in the immobilizing drug combination because of potential confounding respiratory and cardiovascular effects (Portas 2004).

Etorphine plus hyaluronidase were administered using a 3.0 mL plastic dart with a 60 mm uncollared needle propelled from a compressed air rifle (DAN-INJECT, International S.A., South Africa). Once an animal could be safely handled, it was blindfolded and placed in sternal recumbency for 1 minute to facilitate initial instrumentation and subsequently rolled into lateral recumbency. The influence of variable induction times on physiological measurements was reduced by conducting a trial only if the rhinoceros became recumbent and could be safely handled within 15 minutes after darting (Haw et al. 2014). Recumbency was used as an indicator of immobilization level equivalency between trials. Data collection started 10 minutes after initial recumbency ($t = 0$), and was repeated at 5 minute intervals over a 25 minute study period. In those rhinoceros allocated

treatment 2, butorphanol was administered IV at 2 minutes ($t = 2$) as this allowed for instrumentation of study animals and most closely approximated the time at which butorphanol is administered in field-immobilized rhinoceros. In treatment 1, sterile saline was administered at $t = 2$. At the end of each trial, all rhinoceros were weighed and naltrexone was administered (40 mg mL^{-1} , Kyron laboratories) IV at 20 times the etorphine dose.

Expired minute ventilation (corrected for body temperature and saturated pressure; $\dot{V}_{E\text{BTPS}}$; L minute^{-1}) was measured via shortened equine endotracheal (ET) tubes (V KRUUSE I.D. 28, Cat. No. 282270; Jørgen Kruuse A/S, Denmark) inserted into each nostril with the cuffs inflated to create an airtight seal. A two-way Y-shape nonrebreathing valve (2730 Series; Hans Rudolph, Inc., OK, USA) was connected to each ET tube at the nares external margin which allowed inspired air to enter the tube and directed expired air to a PowerLab Exercise Physiology System (ML870B80; ADInstruments, NSW, Australia). The design and function of equipment collecting the expired air was tested in a pilot study to ensure that it did not impede respiratory function. The $\dot{V}_{E\text{BTPS}}$ was determined via a respiratory flow head (MLT1000L) linked to a spirometer (ML140) and a gas mixing chamber (MLA245). The respiratory flow head had a $1000 \text{ L minute}^{-1}$ capacity and was calibrated daily according to manufacturer's instructions. Expired air temperature was recorded by a thermistor pod (ML309) in the mixing chamber.

Expired air was collected into a Douglas bag for 1 minute at the end of each sampling interval and analysed using a Cardiocap/5 (Datex-Ohmeda/GE Healthcare, Finland) for mixed-expired carbon dioxide pressure ($P_{E\text{CO}_2}$; mmHg) and expired oxygen fraction ($F_{E\text{O}_2}$; %). The same monitor was used to analyse expired air from one of the ET tubes to determine end-tidal carbon dioxide pressure (mmHg) and oxygen fraction (%), and measure respiratory frequency (f_R). Body temperature (T ; °C) was measured using a rectal thermometer (BAT-12; Physitemp Instruments, NJ, USA).

A 22 gauge intravenous catheter was inserted into an auricular artery and blood samples collected into heparinized 1 mL syringes and immediately analysed using a portable blood gas analyser (iSTAT 1 Handheld Clinical Analyzer; Heska Corporation, CO, USA) and CG4+ cartridge (iSTAT CG4+ cartridges; Heska Corporation). The alveolar–arterial oxygen gradient [$P_{(A-a)\text{O}_2}$] (mmHg) was calculated using the formula

$F_{i\text{O}_2}(P_B - P_{\text{H}_2\text{O}}) - P_{a\text{CO}_2} - P_{a\text{O}_2}$, with inspired oxygen fraction ($F_{i\text{O}_2}$; %) standardized to 20.9% and barometric pressure (P_B ; mmHg) measured by the portable blood gas analyser prior to each immobilization ($P_{a\text{O}_2}$ = arterial partial pressure of oxygen; $P_{a\text{CO}_2}$ = arterial partial pressure of carbon dioxide). Alveolar vapour pressure of saturated air ($P_{\text{H}_2\text{O}}$; mmHg), at a specific T , was determined using the formula $4.58 \exp [(17.27 T) / (237.3 + T)]$ (Meyer et al. 2010). The expected respiratory minute ventilation (\dot{V}_E ; L minute^{-1}) in the rhinoceros prior to immobilization was estimated from body mass using the formula $0.518 \text{ BM}^{0.802}$ (Bide et al. 1997). Actual respiratory minute ventilation was considered to be equivalent to $\dot{V}_{E\text{BTPS}}$, which was divided by f_R to calculate tidal volume (V_T ; L breath^{-1}).

The Enghoff modified Bohr's equation $[(P_{a\text{CO}_2} - P_{E\text{CO}_2})/P_{a\text{CO}_2}]V_T$ was used to determine physiological dead space ($\dot{V}_{D\text{PHYS}}$; L breath^{-1}) (Tusman et al. 2012). The ($\dot{V}_{D\text{PHYS}}$) was corrected by 300 mL for the volume of the two ET tubes extending beyond the rhinoceros' nostrils.

Oxygen consumption (\dot{V}_{O_2} ; L minute^{-1}) was calculated as the difference between inspired and expired oxygen fractions as a proportion of expired minute ventilation at standard temperature and dry pressure ($\dot{V}_{E\text{STPD}}$), i.e. $\dot{V}_{\text{O}_2} = (F_{i\text{O}_2} - F_{E\text{O}_2}) / 100 \times (\dot{V}_{E\text{STPD}})$ (McArdle et al. 1986). The $\dot{V}_{E\text{BTPS}}$ was multiplied by $(273/310) [(P_B - 47)/760]$ to convert from BTPS to STPD (West 2008). Since inspired and expired minute ventilation were not equivalent (depending on the respiratory quotient), and $\dot{V}_{E\text{STPD}}$ was used to determine both $F_{i\text{O}_2}$ and $F_{E\text{O}_2}$, the Haldane transformation was used to correct the inspired oxygen volume, i.e. $\dot{V}_{\text{O}_2} = \dot{V}_{E\text{STPD}} \{F_{i\text{O}_2} [(1 - F_{E\text{O}_2} + F_{E\text{CO}_2}) / 1 - (F_{i\text{O}_2} + F_{i\text{CO}_2})] - F_{E\text{O}_2}\}$ (McArdle et al. 1986).

Carbon dioxide production (\dot{V}_{CO_2} ; L minute^{-1}) was calculated as the product of $\dot{V}_{E\text{STPD}}$ and the difference between expired and inspired carbon dioxide fractions. Carbon dioxide production was determined using the formula $\dot{V}_{\text{CO}_2} = \dot{V}_{E\text{STPD}}(F_{E\text{CO}_2} - 0.03\%)$. Inspired fractions for oxygen and carbon dioxide were standardized at $F_{i\text{O}_2} = 20.9\%$ and $F_{i\text{CO}_2} = 0.03\%$ (McArdle et al. 1986).

Skeletal muscle tremors, especially of the limbs, head and shoulders, in the immobilized rhinoceros were subjectively evaluated and scored by a single observer at each time point according to criteria in Appendix A. The scores ranged from 1) no visible tremors to 5) severe tremors. Total muscle tremor scores were calculated as the sum of all the scores for that treatment at each time point.

Data analyses

Stata (Stata Statistical Software: Release 14, College Station, TX, USA) was used for statistical analyses. Descriptive statistics were calculated to assess data distribution for each treatment at different sampling points. Due to the relatively small sample size ($n = 6$), nonparametric statistical tests were used to compare median blood gases and respiratory values at specific sampling points within each treatment. The Kruskal–Wallis test was used to assess whether median values for blood gases and respiratory parameters differed over sampling points. Based on these findings, differences in medians between matched pairs of values at $t = 0$ to $t = 10$ were compared using the Wilcoxon rank signed test. To confirm that no further changes occurred after 10 minutes, linear regression (using ranks) was used to assess changes in blood gases and respiratory parameters after 10 minutes using $t = 10$ as the reference value. Correlations between blood gases, respiratory parameters and muscle tremor scores were evaluated using linear regression. To evaluate differences in blood gases and respiratory parameters between treatment groups, linear regression (using ranks) was used to compare median blood gases and respiratory parameters while adjusting for the effect of time. Statistical significance was set at $p < 0.05$ for all statistical tests.

Results

All rhinoceros in both treatments became sternally recumbent within 15 minutes of etorphine administration, and sample and data collection started 10 minutes later. All study animals recovered with no ill effects.

Treatment 1: etorphine

Table 1 shows the blood gas and respiratory values for treatment 1. Median arterial PaO_2 and PaCO_2 did not change significantly during the first 10 minutes ($t = 0$ – 10) or subsequent 15 minutes ($t = 10$ – 25 ; Fig. 1).

The median \dot{V}_E ($t = 0$) was $163 \text{ L minute}^{-1}$. \dot{V}_{EBTPS} declined between $t = 0$ – 10 and $t = 10$ – 25 ; however, these changes were not statistically significant (Fig. 2). The f_R decreased significantly ($p = 0.034$) over the first 10 minutes, but did not change between $t = 10$ and $t = 25$. No changes over time ($t = 0$ – 10 , $t = 10$ – 25) were observed in V_T , P(A-a)O_2 or \dot{V}_{DPHYS} .

The \dot{V}_{O_2} decreased between $t = 0$ and $t = 10$ ($p = 0.046$; Fig. 3a); however, the decrease in \dot{V}_{CO_2}

over the same period was not statistically significant (Fig. 3b). There was a positive correlation between PaCO_2 and \dot{V}_{CO_2} ($p = 0.021$, $r^2 = 0.15$), and inverse correlation between PaO_2 and \dot{V}_{O_2} ($p = 0.038$, $r^2 = 0.12$). Both \dot{V}_{O_2} and \dot{V}_{CO_2} were associated with muscle tremor scores ($p < 0.0001$, $r^2 = 0.52$; $p = 0.0001$, $r^2 = 0.57$, respectively), which decreased over the study period (Fig. 4).

Treatment 2: etorphine and butorphanol

Table 1 shows the blood gases and respiratory values for treatment 2. In etorphine-immobilized rhinoceros administered butorphanol IV, there was a significant increase in PaO_2 ($p = 0.027$) and decrease in PaCO_2 ($p = 0.046$) between $t = 0$ and $t = 10$ with no further significant changes (Fig. 1). Between $t = 0$ and $t = 10$, \dot{V}_{EBTPS} declined ($p = 0.03$) with a decrease in f_R ($p = 0.032$), a reduction in V_T ($p = 0.046$) and drop in \dot{V}_{DPHYS} ($p = 0.03$), then remained unchanged ($t = 10$ – 25 ; Fig. 2). Following butorphanol administration, there was a transient increase in \dot{V}_{EBTPS} , which reached a maximum value at $t = 3$, but returned to preinjection values at $t = 5$. The P(A-a)O_2 did not change significantly during the trial. The \dot{V}_{O_2} declined between $t = 0$ and $t = 10$ ($p = 0.027$) with no further changes over time (Fig. 3a). Trends in \dot{V}_{CO_2} were similar to \dot{V}_{O_2} with declines between $t = 0$ and $t = 10$ ($p = 0.027$), although there was also a decrease from $t = 10$ to $t = 25$ ($p = 0.014$; Fig. 3b). There was a positive association between PaCO_2 and \dot{V}_{CO_2} ($p = 0.0002$; $r^2 = 0.34$), and inverse correlation between PaO_2 and \dot{V}_{O_2} ($p = 0.002$, $r^2 = 0.25$). Both \dot{V}_{O_2} and \dot{V}_{CO_2} were positively associated with muscle tremor scores ($p < 0.0001$, $r^2 = 0.72$; $p < 0.0001$, $r^2 = 0.65$, respectively). Muscle tremors decreased rapidly between $t = 0$ and $t = 5$ and remained low for the rest of the trial period (Fig. 4).

Comparison of treatments 1 and 2

Table 2 shows overall distribution of blood gases and respiratory values between treatments 1 and 2. The PaO_2 was higher ($p < 0.001$) and PaCO_2 was lower ($p = 0.001$) when comparing overall median values in treatment 2 versus treatment 1 (Fig. 1). There were no differences in median values for \dot{V}_{EBTPS} or \dot{V}_{DPHYS} between treatments (Fig. 2). Median f_R was statistically higher ($p = 0.045$) and V_T significantly lower ($p = 0.008$) in animals administered butorphanol. In treatment 2, median P(A-a)O_2 values were lower compared to treatment 1 ($p = 0.019$). Overall \dot{V}_{O_2}

Table 1 Distribution of blood gases and respiratory parameters, median and interquartile range (25–75th percentile), at sampling periods 0, 5, 10, and 25 minutes in six captive male white rhinoceros (age 5–6 years) for two treatments: 1 [(etorphine intramuscularly (IM)); and 2 [etorphine IM and butorphanol intravenously (IV)]

	Treatment ^a	0 minutes	5 minutes	10 minutes	25 minutes
PaO ₂ (mmHg) (kPa)	I	25.0 (23.0–28.0)	25.0 (23.0–28.0)	27.5 (23.0–29.0)	26.0 (25.0–29.0)
	II	3.3 (3.1–3.7)	3.3 (3.1–3.7)	3.7 (3.1–3.9)	3.5 (3.3–3.9)
PaCO ₂ (mmHg) (kPa)	I	25.5 (22.0–26.0)	48.0 (42.0–50.0)	48.5 (42.0–51.0)	43.5 (38.0–46.0)
	II	3.4 (2.9–3.5)	6.4 (5.6–6.7)	6.5 (5.6–6.8)	5.8 (5.1–6.1)
PaCO ₂ (mmHg) (kPa)	I	76.2 (67.2–81.2)	71.1 (60.9–79.6)	72.3 (66.9–81.8)	78.5 (70.8–82.4)
	II	10.2 (9.0–10.8)	9.5 (8.1–10.6)	9.6 (8.9–10.9)	10.5 (9.4–11.0)
V̇ _E BTPS (L minute ⁻¹)	I	81.7 (76.1–89.3)	58.2 (54.7–68.4)	62.8 (57.9–75.2)	63.7 (62.8–65)
	II	10.9 (10.1–11.9)	7.8 (7.8–9.1)	8.4 (7.7–10.0)	8.5 (8.4–8.7)
f _R (breaths minute ⁻¹)	I	164.0 (126.6–182.2)	137.6 (102.5–153.7)	118.7 (88.5–130.6)	96.1 (66.8–101.4)
	II	151.4 (139.0–172.2)	153.0 (125.7–160.5)	89.5 (85.0–98.5)	83.0 (76.6–86.5)
V _T (L breath ⁻¹)	I	10 (9–10)	9 (8–9)	7 (6–7)	6 (5–8)
	II	10 (8–10)	11 (10–13)	8 (6–8)	6 (6–7)
P(A-a)O ₂ (mmHg) (kPa)	I	18.0 (14.1–21.5)	16.1 (14.0–16.7)	18.3 (17.7–18.7)	14.2 (12.7–15.5)
	II	16.8 (15.2–20.4)	12.2 (11.7–16.1)	11.9 (11.0–16.9)	13.4 (11.7–14.4)
P(A-a)O ₂ (mmHg) (kPa)	I	41.7 (36.6–45.1)	48.9 (39.0–53.3)	44.3 (33.8–46.9)	39.0 (33.6–44.1)
	II	5.6 (4.9–6.1)	6.5 (5.2–7.1)	5.9 (4.5–6.3)	5.2 (4.5–5.9)
V̇ _D PHYS (L minute ⁻¹)	I	37.0 (33.3–41.5)	38.9 (32.9–43.1)	37.0 (30.7–39.3)	36.7 (35.0–39.1)
	II	4.9 (4.4–5.5)	5.2 (4.4–5.7)	4.9 (4.1–5.2)	4.9 (4.7–5.2)
V̇ _O ₂ (L minute ⁻¹)	I	59.5 (35.8–69.8)	42.1 (19.3–60.5)	39.5 (23.5–43.7)	35.6 (26.4–48.2)
	II	47.9 (46.9–50.9)	35.9 (17.5–52.2)	31.4 (21.9–41.2)	29.0 (26.1–34.5)
V̇ _{CO} ₂ (L minute ⁻¹)	I	11.1 (10.0–12.0)	8.7 (7.7–9.8)	7.8 (5.8–8.6)	6.4 (3.7–7)
	II	10.9 (9.1–12.0)	6.8 (5.5–8.0)	4.4 (3.6–5.1)	4.2 (4.0–4.6)
V̇ _{CO} ₂ (L minute ⁻¹)	I	8.3 (7.8–11.4)	6.8 (5.9–7.8)	6.0 (4.9–7.1)	4.3 (3.5–5.3)
	II	9.3 (7.2–10.4)	6.9 (6.2–7.7)	4.2 (3.8–4.4)	3.7 (2.7–3.8)

PaO₂, arterial partial pressure oxygen; PaCO₂, carbon dioxide; V̇_EBTPS, expired minute ventilation; f_R, respiratory rate; V_T, tidal volume; P(A-a)O₂, alveolar–arterial oxygen gradient; V̇_DPHYS, physiological dead space; V̇_O₂, oxygen consumption; V̇_{CO}₂, carbon dioxide production.

^aTreatment 1, etorphine IM (2.5 mg etorphine, 1000–1250 kg; 3.0 mg etorphine, 1250–1500 kg); Treatment 2, etorphine IM (2.5 mg etorphine 1000–1250 kg; 3.0 mg etorphine 1250–1500 kg) and butorphanol IV (25 mg butorphanol, 1000–1250 kg; 30 mg butorphanol, 1250–1500 kg).

and V̇_{CO}₂ were lower in treatment 2 compared to treatment 1 animals ($p = 0.001$; **Figs 3a, b**).

Discussion

In this study, immobilization of white rhinoceros with etorphine resulted in hypoxaemia and hypercapnia. Contrary to previous reports, these changes were not associated with a decrease in respiratory minute ventilation but rather an increase in alveolar–arterial oxygen gradient and oxygen consumption. Administration of IV butorphanol was followed by improvements in arterial oxygen and carbon dioxide tensions, although animals remained hypoxaemic and hypercapnic. Improved blood gases appeared to result primarily from a decrease in oxygen consumption associated with decreased muscle tremors, rather than from changes in ventilation.

Hypoxaemia and hypercapnia are frequently reported in etorphine-immobilized rhinoceros and extremes of blood gases measured in our study (PaO₂ of

about 25 mmHg and PaCO₂ of about 80 mmHg) suggest they can be life-threatening (Yaksh & Wallace 2011; Haw et al. 2014; Boardman et al. 2014). A PaO₂ ≤ 80 mmHg indicates hypoxaemia in an anaesthetized animal and animals with values < 60 mmHg normally require supportive treatment (Read 2003). A PaCO₂ > 70 mmHg may produce myocardial depression, arrhythmias and impaired metabolism because of respiratory acidosis, and ventilator support is advocated (Moens 2013). In addition, opioid chemoreceptor depression can profoundly depress hypercapnic and hypoxic ventilatory responses (Pattinson 2008).

In etorphine-immobilized rhinoceros, initial median V̇_EBTPS was clinically similar to V̇_E at rest. These results suggest that the marked hypoxaemia and hypercapnia measured in the first arterial sample were not a consequence of a reduction in ventilation, as has been previously advocated (Kock et al. 1995; Miller et al. 2013; Boardman et al. 2014; Haw




Figure 1 Median and interquartile range of arterial partial pressures of (a) oxygen (PaO₂) and (b) carbon dioxide (PaCO₂) at sampling periods 0, 5, 10, 15, 20 and 25 minutes in six captive male white rhinoceros (age, 5–6 years) for treatment 1 [etorphine intramuscularly (IM)] or treatment 2 [etorphine IM and butorphanol intravenously (IV)]. The dashed line indicates the time at which butorphanol was administered. * indicates a significant ($p < 0.05$) difference within treatment between $t = 0$ and $t = 10$, +, indicates a significant ($p < 0.05$) difference within treatment between $t = 10$ and $t = 25$, #, indicates a significant ($p < 0.05$) difference in overall median values between treatments 1 and 2 from 5 to 25 minutes.

et al. 2014). It is unlikely that the absence of reduced $\dot{V}_{E_{BTPS}}$ was because of a delay in maximum respiratory opioid effect as etorphine had caused sufficient central nervous system depression to induce a state of immobilization and recumbency for 10 minutes prior to sampling. Meyer *et al.* (2015) reported similar results in that hypoventilation was not the primary cause of hypoxaemia and hypercapnia in etorphine-immobilized domestic goats.

The initial median f_R (10 breaths minute^{-1}) was substantially lower than the rate (16–23 breaths minute^{-1}) reported for standing unrestrained captive white rhinoceros (Citino and Bush 2007). These values suggest that $\dot{V}_{E_{BTPS}}$ was maintained by an increase in V_T in the immobilized rhinoceros. In the study rhinoceros, the increased V_T may have been a compensatory response to hypoxaemia and/or, hypercapnia. It is also possible that initial

sympathetic stimulation associated with darting influenced respiratory minute ventilation (Heistad *et al.* 1972).

In rhinoceros immobilized only with etorphine, hypoxaemia and hypercapnia did not change significantly over the 25 minute study interval. Although median $\dot{V}_{E_{BTPS}}$ also did not change significantly, the initial and final values of 164 and 96 L minute^{-1} , respectively, may reflect a clinically relevant reduction in minute volume. A decrease in $\dot{V}_{E_{BTPS}}$ is contradictory to the expected response to hypercapnia and hypoxia (Pattinson, 2008). Opioids suppress the ventilatory response to hypercapnia and hypoxia through reduced excitation of chemosensory neurons and chemoreceptor bodies (Yaksh & Wallace 2011). It is possible that etorphine receptor binding increased over time, causing further depression of respiratory rhythmogenesis resulting in hypopnoea;




Figure 2 Median and interquartile range of expired minute ventilation ($\dot{V}_{E_{BTPS}}$) measured at 1 minute intervals between 0 and 25 minutes in six captive male white rhinoceros (age, 5–6 years) for treatment 1 [(etorphine intramuscularly (IM)] or treatment 2 [etorphine IM and butorphanol intravenously (IV)]. The dashed line indicates the time at which butorphanol was administered. *, indicates a significant ($p < 0.05$) difference within treatment between $t = 0$ and $t = 10$; +, indicates a significant ($p < 0.05$) difference within treatment between $t = 10$ and $t = 25$; #, indicates a significant ($p < 0.05$) difference in overall median values between treatments 1 and 2 from 5 to 25 minutes.

however, it is likely that peak respiratory perturbations had been reached at initial sample collection in the immobilized rhinoceros at approximately 20 minutes post-dart (Yaksh & Wallace 2011). The decrease in $\dot{V}_{E_{BTPS}}$ may have been because of an opioid-induced diminished hypoxic drive associated with a lowered set-point. An arterial oxygen tension of 25 mmHg in immobilized rhinoceros should have stimulated an increase in ventilation, which was not observed. In opioid-immobilized rhinoceros, the hypothetically lowered hypoxic threshold would have resulted in a decreased respiratory stimulus at $PaO_2 > 25$ mmHg with a subsequent fall in ventilation and limited changes in PaO_2 (Pattinson 2008).

The elevated $P(A-a)O_2$ observed at $t = 0$ may have contributed to decreased PaO_2 and increased $PaCO_2$. The normal resting A-a gradient is unknown for white rhinoceros; however, a gradient of 41.7 mmHg in the immobilized rhinoceros, compared to 10 mmHg in horses at rest, suggests a clinically relevant finding (Doherty & Valverdeis 2008). An elevated A-a gradient is indicative of ventilation/perfusion mismatching, a physiologic right-to-left shunt, or impaired diffusion of gases between alveoli and perfusing blood (West 2008). A decreased \dot{V}/\dot{Q} ratio does not usually result in hypercapnia since increasing $PaCO_2$ stimulates respiration. However,

this response may be limited in opioid-immobilized rhinoceros because of alterations in chemoreceptor sensitivity (Buss et al. 2015). Etorphine-induced pulmonary hypertension may contribute to hypoxaemia by hindering gas exchange across alveolar–capillary membranes because of pulmonary congestion with interstitial oedema or a decrease in blood flow passage time through pulmonary vasculature (Meyer et al. 2015).

Oxygen consumption is a measure of metabolic activity and maintenance of homeostatic processes in mammals (Porter & Brand 1995). The \dot{V}_{O_2} was elevated in etorphine-immobilized rhinoceros ($8.23 \text{ mL kg}^{-1} \text{ minute}^{-1}$) compared to horses ($3 \text{ mL kg}^{-1} \text{ minute}^{-1}$) at rest and may have contributed significantly to hypoxaemia (Evans & Rose 1988). The difference in \dot{V}_{O_2} may be the result of species differences; however, metabolic rate per unit body mass tends to decrease with increased size, so a rhinoceros should consume less oxygen per unit body mass than should a horse (Porter & Brand 1995). Increased skeletal muscle activity may be an important contributor to metabolic demands, an idea supported by our findings of a strong correlation between \dot{V}_{O_2} and muscle tremor scores in immobilized rhinoceros.

The overall decrease in $\dot{V}_{E_{BTPS}}$ over time in treatment 2 supports the hypothesis that IV administration




Figure 3 Median and interquartile range of (a) oxygen consumption ($\dot{V}O_2$) and (b) carbon dioxide production ($\dot{V}CO_2$) calculated for sampling periods 0, 5, 10, 15, 20 and 25 minutes in six captive male white rhinoceros (age, 5–6 years) for treatment 1 [(etorphine intramuscularly (IM)] or treatment 2 [etorphine IM and butorphanol intravenously (IV)]. The dashed line indicates the time at which butorphanol was administered. *, indicates a significant ($p < 0.05$) difference within treatment between $t = 0$ and $t = 10$; +, indicates a significant ($p < 0.05$) difference within treatment between $t = 10$ and $t = 25$; #, indicates a significant ($p < 0.05$) difference in overall median values between treatments 1 and 2 from 5 to 25 minutes.




Figure 4 Muscle tremor scores at sampling periods 0, 5, 10, 15, 20 and 25 minutes were the sum of all the scores (1 to 5) at each time point in six captive male white rhinoceros (age, 5–6 years) for treatment 1 [(etorphine intramuscularly (IM)] or treatment 2 [etorphine IM and butorphanol intravenously (IV)]. The scores ranged from 1) no visible tremors to 5) severe tremors.

Table 2 Overall distribution of blood gases [arterial partial pressure oxygen (PaO₂) and carbon dioxide (PaCO₂), expired minute ventilation ($\dot{V}_{E\text{BTPS}}$), respiratory rate (f_R), tidal volume (V_T), alveolar–arterial oxygen gradient (P(A-a)O₂), physiological dead space ($\dot{V}_{D\text{PHYS}}$), oxygen consumption (\dot{V}_{O_2}), and carbon dioxide production (\dot{V}_{CO_2})] and respiratory parameters, median and interquartile range (25–75th percentile), over the sampling period 5–25 minutes in six captive male white rhinoceros (age, 5–6 year) for two treatments: 1 [(etorphine intramuscularly (IM)); and 2 [etorphine IM and butorphanol intravenously (IV)]

^a Treatment	1		2	
PaO ₂ (mmHg)	26	(23–29)	42.5	(30.5–48.5)
(kPa)	3.5	(3.1–3.9)	5.7	(4.1–6.5)
PaCO ₂ (mmHg)	75.5	(67.4–82.3)	64.3	(61.1–76.1)
(kPa)	10.1	(9.0–11.0)	8.6	(8.1–10.1)
$\dot{V}_{E\text{BTPS}}$ (L minute ⁻¹)	105.9	(84.3–137.9)	90.0	(84.7–137.9)
f_R (breaths minute ⁻¹)	7.0	(5–7)	7.0	(6–9)
V _T (L minute ⁻¹)	16.3	(14.0–18.3)	14.1	(11.8–16.2)
P(A-a)O ₂ (mmHg)	41.5	(34.1–45.8)	36.8	(33.2–40.3)
(kPa)	5.5	(4.5–6.1)	4.9	(4.4–5.4)
\dot{V}_{O_2} (L minute ⁻¹)	7.3	(5.6–8.7)	4.8	(4.1–6.8)
\dot{V}_{CO_2} (L minute ⁻¹)	5.8	(4.2–7.2)	4.1	(3.8–6.3)

^aTreatment 1, etorphine IM (2.5 mg etorphine, 1000–1250 kg; 3.0 mg etorphine, 1250–1500 kg); Treatment 2, etorphine IM (2.5 mg etorphine 1000–1250 kg; 3.0 mg etorphine 1250–1500 kg) and butorphanol IV (25 mg butorphanol, 1000–1250 kg; 30 mg butorphanol, 1250–1500 kg).

of butorphanol in etorphine-immobilized rhinoceros does not improve respiratory minute ventilation. The $\dot{V}_{E\text{BTPS}}$ increased for a few minutes following the administration of butorphanol then subsequently decreased, similar to treatment 1, so overall there was no significant difference between the two treatments. The $\dot{V}_{D\text{PHYS}}$ was also similar between treatments, suggesting changes in deadspace ventilation did not influence blood gases following butorphanol administration. The P(A-a)O₂ was generally lower in treatment 2 compared to treatment 1, but an initial lower value and no significant change following butorphanol administration, suggests that this finding is of limited clinical importance. Further investigations are needed to confirm the consistency and significance of this difference between treatments as changes in alveolar ventilation: perfusion ratios, shunting, and gas diffusion rates across alveolar–capillary membranes could alter arterial blood gases (Haw et al. 2014; West 2008).

The results of this study suggest that the improvements in PaO₂ and PaCO₂, following butorphanol administration in etorphine-immobilized white rhinoceros, arose from changes in metabolic oxygen consumption associated with decreased muscle tremors. An inverse correlation between PaO₂ and \dot{V}_{O_2} , and a positive correlation between \dot{V}_{O_2} and muscle tremor scores, suggest that these three variables are interrelated. We propose that an increase in PaO₂ following IV butorphanol

administration is a result of decreased metabolism associated with reduced muscle tremors. Carbon dioxide production followed a similar trend to that of oxygen consumption which further supports the hypothesis that blood gas changes following butorphanol administration result from reduced metabolic activity. In shivering humans recovering from induced hypothermia, it has been shown that carbon dioxide production and oxygen consumption are positively associated with muscular activity and metabolic rate (Ralley et al. 1988). A reduction in muscle tremors associated with butorphanol administration may be mediated through antagonism of etorphine-induced sympathetic nervous system activity. Muscle tremors in immobilized rhinoceros are significantly associated with increased plasma catecholamine concentrations (De Lange 2015). The muscle tremors did not typify opioid-induced changes in locomotion often observed during induction in white rhinoceros. An alternative explanation for the improved arterial blood gases following butorphanol administration may be increased pulmonary perfusion and reduction in \dot{V}/\dot{Q} inequality rather than changes in oxygen consumption (Wagner 2008). However, heart rate and systemic blood pressure decreased in etorphine-immobilized rhinoceros following butorphanol administration, suggesting a decrease in cardiac output (Buss et al. 2016).

The transient increase in ventilation following the administration of butorphanol is of clinical

significance in that it has the potential to mislead the uninformed observer. It may appear that butorphanol administration improves ventilation by increasing respiratory rate; however, within a few minutes, respiratory minute ventilation was the same as in opioid-immobilized rhinoceros that were not administered butorphanol.

The use of only six rhinoceros because of welfare considerations and logistical challenges is a study limitation. The small sample size may have resulted in lack of statistical differences and masked potentially clinically important physiological changes. Inability to determine alveolar ventilation, cardiac output, pulmonary artery pressures, shunt fractions and \dot{V}/\dot{Q} ratios, in part, limited a comprehensive understanding of physiological mechanisms influencing arterial blood gases. Physiological differences may exist between free-ranging and captive rhinoceros; therefore, further studies should compare these conditions.

Conclusion

Our findings support the hypothesis that butorphanol does not improve respiratory minute ventilation in etorphine-immobilized white rhinoceros. However, butorphanol did improve arterial oxygen and carbon dioxide tensions, probably as a consequence of reduced metabolism and muscle tremors. We also showed that hypoxaemia and hypercapnia following etorphine administration were not a result of decreased respiratory minute ventilation. An increase in the alveolar–arterial oxygen gradient probably contributed to hypoxia in immobilized white rhinoceros. However, the impact of and underlying physiological processes leading to changes in alveolar–arterial gradient require further elucidation. Our findings provide evidence that hypoxia and hypercapnia in immobilized rhinoceros result also from increases in metabolic oxygen consumption and carbon dioxide production with inadequate ventilatory compensation.

Acknowledgements

The authors thank Markus Hofmeyr, Marius Kruger, Milandie Kruger, Leana Rossouw, Guy Hausler and holding-facility staff of Veterinary Wildlife Services, Kruger National Park. The assistance of members of the Brain Function Research Group, University of Witwatersrand Medical School, and staff and students of the Faculty of Veterinary Science, University of Pretoria, is also acknowledged. Financial and in-kind support was provided by South African

National Parks. The project was supported by funding grants from the South African Veterinary Foundation and University of Witwatersrand (00125485211015 12110500000000000000004992).

Authors contributions

PB: project designed and implementation, data collection, collation and analyses, and preparation of manuscript. MM: project implementation, including data collection, and preparation of manuscript. AF: project design and implementation, including data collection, and preparation of manuscript. AH: project implementation, including data collection, and preparation of manuscript. ES: statistical analysis of data and preparation of manuscript. FOP: project design, statistical analysis of data and preparation of manuscript. LM: project design, project implementation, including data collection, and preparation of manuscript.

Conflict of interest statement

There are no conflicts of interests by the authors.

References

- Bide RW, Armour SJ, Yee E (1997) Estimation of human toxicity from animal inhalation toxicity data: 1. Minute volume-body weight relationships between animals and man. Suffield Report 673, Defense Research Establishment Suffield Ralston, Canada.
- Boardman WSJ, Caraguel CGB, Raath JP, Van Zijl Langhout M (2014) Intravenous butorphanol improves cardiopulmonary parameters in game-ranched white rhinoceroses (*Ceratotherium simum*) immobilized with etorphine and azaperone. *J Wildl Dis* 50, 849–857.
- Buss P, Miller M, Fuller A et al. (2016) Cardiovascular effects of etorphine, azaperone and butorphanol combinations in chemically immobilized captive white rhinoceros (*Ceratotherium simum*). *J Zoo Wildl Med* 47, 834–843.
- Buss P, Olea-Popelka F, Meyer L et al. (2015) Evaluation of cardiorespiratory, blood gas, and lactate values during extended immobilization of white rhinoceros. *J Zoo Wildl Med* 46, 224–233.
- Citino SB, Bush M (2007) Reference cardiopulmonary physiologic parameters for standing, unrestrained white rhinoceroses (*Ceratotherium simum*). *J Zoo Wildl Med* 38, 375–379.
- De Lange SS (2015) Tremors in white rhinoceros (*Ceratotherium simum*) during chemical immobilization. Master's dissertation, Faculty of Veterinary Science, University of Pretoria.
- Doherty T, Valverdeis A (2008) *Manual of Equine Anesthesia*. Wiley-Blackwell, NJ, USA.

- Evans DL, Rose RJ (1988) Cardiovascular and respiratory responses in thoroughbred horses during treadmill exercise. *J Exp Biol* 134, 397–408.
- Haw A, Hofmeyr M, Fuller A et al. (2014) Butorphanol with oxygen insufflation corrects etorphine-induced hypoxaemia in chemically immobilized white rhinoceros (*Ceratotherium simum*). *BMC Vet Res* 10, 1–9.
- Haw A, Hofmeyr M, Fuller A et al. (2015) Butorphanol with oxygen insufflation improves cardiorespiratory function in field-immobilised white rhinoceros (*Ceratotherium simum*). *J S Afr Vet Assoc* 86, E1–E10.
- Heistad DD, Wheeler RC, Mark AL et al. (1972) Effects of adrenergic stimulation on ventilation in man. *J Clin Invest* 51, 1469–1475.
- Kock M, Morkel P, Atkinson M, Foggin C (1995) Chemical immobilization of free-ranging white rhinoceros (*Ceratotherium simum simum*) in Hwange and Matobo national parks, Zimbabwe, using combinations of etorphine (M99), fentanyl, xylazine, and detomidine. *J Zoo Wildl Med* 26, 207–219.
- McArdle WD, Katch FL, Katch VL (1986) Exercise Physiology: Energy, Nutrition, and Human Performance (2nd edn). Lea & Febiger, PA, USA, pp. 637–639.
- Meyer LCR, Hetem R, Fick L et al. (2010) Effects of serotonin agonists and doxapram on respiratory depression and hypoxemia in etorphine-immobilized impala (*Aepyceros melampus*). *J Wildl Dis* 46, 514–524.
- Meyer LCR, Hetem RS, Mitchell D, Fuller A (2015) Hypoxia following etorphine administration in goats (*Capra hircus*) results more from pulmonary hypertension than from hypoventilation. *BMC Vet Res* 11, 18.
- Miller M, Buss P, Joubert J et al. (2013) Use of butorphanol during immobilization of free-ranging white rhinoceros (*Ceratotherium simum*). *J Zoo Wildl Med* 44, 55–61.
- Moens Y (2013) Mechanical ventilation and respiratory mechanics during equine anesthesia. *Vet. Clin Equine* 29, 51–67.
- Pattinson KTS (2008) Opioids and the control of respiration. *Br J Anaesth* 100, 747–758.
- Portas TJ (2004) A review of drugs and techniques used for sedation and anaesthesia in captive rhinoceros species. *Aust Vet J* 82, 542–549.
- Porter RK, Brand MD (1995) Cellular oxygen consumption depends on body mass. *Am J Physiol* 269, R226–R228.
- Ralley FE, Wynands JE, Ramsay JG et al. (1988) The effects of shivering on oxygen consumption and carbon dioxide production in patients rewarming from hypothermic cardiopulmonary bypass. *Can J Anaesth* 35, 332–337.
- Read MR (2003) A review of alpha2 adrenoceptor agonists and the development of hypoxemia in domestic and wild ruminants. *J Zoo Wildl Med* 34, 134–138.
- Swan GE (1993) Capture drugs. In: The Capture and Care Manual. McKenzie AA (ed.). Wildlife Decision Support Services, RSA, pp. 1–69.
- Tusman G, Sipmann FS, Bohman SH (2012) Rationale of dead space measurement by volumetric capnography. *Anesth Analg* 114, 866–874.
- Wagner PD (2008) Causes of a high physiological dead space in critically ill patients. *Crit. Care* 12, 148.
- Wenger S, Boardman W, Buss P et al. (2007) The cardiopulmonary effects of etorphine, azaperone, detomidine, and butorphanol in field-anesthetized white rhinoceros (*Ceratotherium simum*). *J Zoo Wildl Med* 38, 380–387.
- West JB (2008) Respiratory physiology: the essentials (8th edn). Lippincott Williams & Wilkins, PA, USA, pp. 172–173.
- Yaksh TL, Wallace MS (2011) Opioids, analgesia, and pain management. In: Goodman & Gilman's The Pharmacological Basics of Therapeutics (12th edn). Brunton LL, Chabner BA, Knollman BC (eds). McGraw Hill Medical, NY, USA, pp. 481–523.

Received 17 July 2016; accepted 30 March 2017.

Available online 17 April 2017

Appendix A. Muscle tremor scores. Criteria for subjectively scoring muscle tremors

Degree of muscle tremor
Level 5 Severe tremors – resulting in whole body and head movement
Level 4 Moderate tremors – resulting in severe shoulder, chest, leg and foot movement
Level 3 Slight tremors – resulting in minor shoulder, chest and severe leg and foot movement
Level 2 Mild tremors – resulting in minor leg and foot movement
Level 1 No visible tremors
