



Ophthalmology of Perissodactyla: Zebras, Tapirs, Rhinoceroses, and Relatives 34

Brian C. Gilger and Andrew G. Matthews



© Chrisoula Skouritakis

Introduction

In phylogenetic systematics, Perissodactyla represents an order of the class Mammalia and comprises the odd-toed ungulates. The order is conventionally divided into two Sub-Orders: Hippomorpha and Ceratomorpha (Table 34.1). Hippomorpha comprises the family Equidae with eight extant species of horses, asses, and zebras in the single genus *Equus*. Ceratomorpha comprises the monogeneric family Tapiridae, with five extant species, and the Rhinocerotidae, with five currently surviving species of rhinoceros in four genera (Table 34.1).

In terms of conservation status, with the exception of the domestic horse and the kiang, the members of the order represent some of the planet's more threatened species,

with some being classed currently as critically endangered (IUCN Red List of Threatened Species: <https://www.iucnredlist.org>).

The early ancestors of modern Perissodactyls divided into Hippomorpha and Ceratomorpha during the early Eocene period (58-52 million years ago; mya), at which time they appear to have been an abundant and highly successful group of mammals with extensive speciation widely distributed across North America, Europe, and Asia. The subsequent division of the Ceratomorpha into Tapiridae and Rhinocerotidae occurred during the late Eocene and Oligocene periods (50-25 mya) and in the case of the tapir species, evolution continued into the Miocene and late Pliocene (11-2 mya). As radiation and speciation progressed within the order, the fossil record shows concurrent extinctions and geographic migration of vulnerable species, influenced by climate change, habitat loss, competition for nutrients and, possibly, human predation, resulting in the distribution and population types now extant. In the case of the Hippomorpha, both fossil records and karyotyping studies (Trifonov et al. 2008) have shown extensive evolutionary radiation during the Miocene, but the modern horse (*Equus* spp.), the only

B. C. Gilger (✉)
College of Veterinary Medicine, North Carolina State University,
Raleigh, NC, USA
e-mail: brian_gilger@ncsu.edu

A. G. Matthews
Bents Cottage, LunanBay, Inverkeilor, Angus, Scotland, UK

Table 34.1 Species diversity of currently extant members of Order Perissodactyla (After Steiner and Ryder 2011)

Sub-order	Family	Species	Common name
Hippomorpha	Equidae	<i>Equus caballus</i>	Domestic horse
		<i>E. przewalskii</i>	Przewalski's wild horse
		<i>E. asinus</i>	African wild ass
		<i>E. kiang</i>	Kiang
		<i>E. hemionus</i>	Asiatic wild ass (Onager)
		<i>E. zebra</i>	Mountain zebra
		<i>E. quagga</i> ^a	Plains zebra ^a
		<i>E. grevyi</i>	Grevy's zebra
Ceratomorpha	Tapiridae	<i>Tapirus indicus</i>	Malayan/Asian tapir
		<i>T. terrestris</i>	Lowland/Brazilian tapir
		<i>T. pinchaque</i>	Mountain tapir
		<i>T. bairdii</i>	Baird's tapir
	Rhinocerotidae	<i>T. kabomani</i>	Little black tapir
		<i>Diceros bicornis</i>	Black rhinoceros
		<i>Ceratotherium simum</i>	White rhinoceros
		<i>Rhinoceros unicornis</i>	Indian rhinoceros
		<i>Dicerorhinus sumatrensis</i>	Sumatran rhinoceros
		<i>Rhinoceros sondaicus</i>	Javanese rhinoceros

^aFormerly known as *E. burchellii*. Burchell's zebra is now classed a subspecies of *E. quagga*

Table 34.2 Defining anatomical and physiological characteristics of members of the order Perissodactyla

	Molar dentition	Alimentary tract	Forelimb phalanges (no)	Hindlimb phalanges (no)
Equidae	Hypsodont	Simple stomach and fermentative hindgut	1	1
Tapiridae	Brachydont	Simple stomach and fermentative hindgut	4	3
Rhinocerotidae	Hypsodont	Simple stomach and fermentative hindgut	3	3

extant genus, appeared relatively late, during the Pliocene (2–4 mya) (Steiner and Ryder 2011).

All Perissodactyls are characteristically mesaxonic, where the sagittal plane of symmetry of the distal limb passes through the third phalanx. Their geographic range in the wild is restricted to mainly subtropical or tropical lowland habitats, ranging from grassland savannahs to rainforests. Outliers are the Mountain tapir, found in the central Andean cloud forest, the Mountain zebra, whose preferred habitat is the more mountainous areas of southern Africa, and the Kiang found on the Tibetan Plateau. All members of the order are herbivores, either predominantly grazers (Equidae), predominantly browsers (Tapiridae) or both grazers and browsers (Rhinocerotidae), this representing anatomical and physiological adaptation to their preferred habitats (Coimbra and Manger 2017). The defining anatomical and physiological characteristics of the Perissodactyls are shown in Table 34.2.

Ocular Anatomy and Function

In terms of foraging behavior, Perissodactyla are either cathemeral (i.e., irregularly active at any time) or diurnal (Equidae, Rhinocerotidae, *T. terrestris* and *T. pinchaque*), or are nocturnal crepuscular (i.e., twilight active) feeders

(other Tapiridae) (García et al. 2012; Banks et al. 2015). These patterns of feeding activity reflect the nature of the habitat and the need to avoid potential predatory threats and will determine to a significant extent visual requirement and ocular structure and function (Peichl 2005; Veilleux and Lewis 2011; Veilleux and Kirk 2014; Banks et al. 2015).

In general, terrestrial mammal visual systems reflect specialization for habitat variation in ambient light intensity (Veilleux and Lewis 2011). Decreasing corneal diameter relative to the axial length of the eye reduces visual sensitivity but enlarges the retinal image and ameliorates peripheral distortion of the image, and is recognized as potentially conferring some survival advantage in open habitat prey ungulates. However, studies on Perissodactylae have failed to identify differences in corneal diameter: ocular axial length ratios, irrespective of their occupying primarily either open or afforested habitats (Veilleux and Lewis 2011; Veilleux and Kirk 2014).

Like the domestic horse, non-domestic *Equidae* have relatively large, laterally placed eyes with horizontal pupils and well-developed *granula iridica* (Johnson 1901; Banks et al. 2015). The horizontal, oblate pupil increases the horizontal depth of field and minimizes blurring of peripheral vision resulting from astigmatism of oblique incidence (Banks et al. 2015) (Fig. 34.1), resulting in increased image quality both of



Fig. 34.1 Zebra eye. (a) Note the horizontal, oblate pupil similar to that observed in a domestic horse. (b) Dilated zebra eye, with the pupil maintaining only a slightly ovoid shape upon full mydriasis. (a) Courtesy of Dr. Ronald K. Cott. (b) Courtesy of Dr. Bret A. Moore

the ground immediately in front and of the area behind. In addition, using head pitch, the animal appears able to align the long axis of the pupil with the horizon (Banks et al. 2015). These evolutionary adaptations will confer a significant survival advantage to a cathemeral prey species occupying open grasslands.

The anatomic ocular dimensions, including anterior chamber depth, lens thickness, axial length and corneal curvature, have been measured in 12 eyes of eight Grevy's zebra, aged 4–14 years, using ultrasound biometry (Evans et al. 2009). The mean (\pm SD) axial length (40.7 ± 1.1 mm), anterior chamber depth (6.9 ± 0.30 mm), and lens thickness (13.83 ± 4.24 mm) were essentially similar to those recorded in the domestic horse (Grinninger et al. 2010). Mean intraocular pressure (IOP) using applanation tonometry was 21.77 mmHg (Evans et al. 2009) (Appendix C). In another study using applanation tonometry, IOP recorded in six eyes of Plains zebras (*Equus burchellii*) was 29.5 ± 3.4 mmHg (Ofri et al. 1998) (Appendix C).

The *Equidae* fundus is paucicellular and with an extensive fibrous tapetum. The tapetal reflex in non-domestic *Equidae* varies from blue-yellow to green-yellow (Johnson 1901) (Fig. 34.2). The neurosensory retina is rod dominated, with increasing cone and ganglion cell numbers in the photoreceptor dense macular areas. These comprise the near contiguous *area centralis* and the visual streak, which are located dorsotemporally to the optic disc and in outline approximate to the shape of the horizontal pupil (Sandmann et al. 1996; Ehrenhofer et al. 2002; Peichl 2005). The macular areas are thought to be used for binocular vision and will permit high acuity dichromatic blue/yellow, color vision.

The *Rhinocerotidae* eye is small relative to the domestic horse, with an ultrasonographically measured mean axial length of 2.61 ± 0.11 cm (Bapodra and Wolfe 2014) compared with 4.05 ± 0.27 cm in the horse (Grinninger et al. 2010). The rounded pupil of *Rhinocerotidae* spp. is typically associated with cathemeral foraging and the absence of significant predation risk (Banks et al. 2015). Anecdotal commentary, presumably made by smug survivors of a close encounter, has it that rhinoceroses are notoriously short sighted. However, rhinoceroses are mildly hyperopic (Howland et al. 1993), and ganglion cell topographic studies on both the black and the white rhinoceros have shown the presence of a visual streak with an *area centralis* both temporally and nasally. In the white rhinoceros, the visual streak lies dorsal to the optic disc, and in the black the optic disc lies within the visual streak (Pettigrew and Manger 2008; Coimbra and Manger 2017) (Fig. 34.3). It is thought that the appearance of a nasally located *area centralis* may compensate for limited lateral head movement in these species (Coimbra and Manger 2017). Calculations based on peak ganglion cell density and axial length of the eyes indicate that *Rhinocerotidae* has a visual resolution of 6–7 cycles/degree. This compares to 60 cycles/degree in humans, 25 cycles/degree in horses and 6 cycles/degree in rabbits, and allows for the prediction that rhinoceroses, in optimal conditions of contrast and luminance, can discern an adult human at between 100–200 m (Pettigrew and Manger 2008; Coimbra and Manger 2017). The greater visual resolution in the horse is likely to be a consequence of larger eye size rather than increased ganglion cell density (Coimbra and Manger 2017). *Rhinocerotidae* have heavily pigmented




Fig. 34.2 Normal fundus images of an (a) adult and (b) foal zebra. Note the paucirangiotic vasculature pattern. The blood vessels are fine and present only in the direct area of the optic disc and cross its margin to extend a short distance into the retina. The optic disc is horizontally oval, and the tapetal reflex is green to green-yellow. In the foal, the optic nerve is hyperemic

Fig. 34.3 Topographic flat-mount retinal map demonstrating retinal ganglion cell densities in the White Rhinoceros (*C. simon*). There is a well-defined horizontal streak of increased retinal density from 200 cells/mm² to a peak density temporally (approximately 2000 cells/mm²) and nasally (approximately 1800 cells/mm²). The black circle near the center of the figure represents the optic disc. T temporal, V ventral. Modified, with permission, from Coimbra and Manger (2017) *Journal of Comparative Neurology* 525: 2484–2498




Fig. 34.6 Rhinoceros with a chronic, intermittently ulcerative, keratitis, which was suspected to be immune-mediated and triggered by UV light exposure. (Photograph courtesy of Dr. Gil Ben-Shlomo)




Fig. 34.7 White rhinoceros with a presumptive diagnosis of eyelid Habronemiasis resulting in granulomatous blepharitis and conjunctivitis. (Photograph courtesy of Dr. José Ricardo Pachaly)



Tapiridae

Ocular surface inflammation, especially ulcerative keratitis and conjunctivitis, is reported to be common in captive tapir most likely caused by trauma and exposure to UV light, which free-ranging tapirs have little exposure to since their

natural habitat is primarily dense jungles (Montiani-Ferreira 2001; Da Silva et al. 2013).

Two female South American Tapirs (*T. terrestris*) at a zoological park were observed to have unilateral blepharospasm and epiphora as a result of deep central corneal ulcers. Successful treatment consisted of topical tobramycin and serum. Medication of ocular surface within the deep-set



Fig. 34.8 Chronic ulcerative keratitis in a Malayan tapir (*T. indicus*). Chronic ulcerative stromal disease developed bilaterally (a). Two months following placement of an episcleral cyclosporine implant (b)

demonstrating a central corneal scar, but the resolution of the stromal keratitis. This chronic keratitis is suspected to be caused or perpetuated by excessive UV light exposure

orbits was facilitated by use of a flexible intravenous catheter inserted into the medial canthus (Da Silva et al. 2013).

Malayan tapirs (*T. indicus*) at a zoological park in South Florida develop chronic, proliferative, and bilateral corneal changes typically by one year of age or within one year of arrival at the park. Biopsy of the raised irregular corneal lesions suggested a diagnosis of papilloma. In a 26-year-old, wild caught, male Malayan tapir, presenting with similar corneal lesions, 5-fluorouracil (25 mg in 0.5 mL) was injected subconjunctivally and repeated at 3, 7, 12, and 17 weeks. The lesions decreased in size, and an improvement in visual function was present for up to 2 years following the series of injections (Karpinski and Miller 2002).

Malayan tapirs (*T. indicus*) have also been observed (Karpinski L, personal communication) to develop chronic bilateral keratopathy, including opacification and ulceration, associated with excessive UV exposure, similar to that observed in marine pinnipeds (Miller et al. 2013). As in the case of the affected pinnipeds, the Malayan tapirs responded to topical or episcleral cyclosporine (Fig. 34.8).

Summary

Very little objective information has been published on the incidence and management of ocular diseases among members of the order Perissodactyla. In terms of general clinical guidance, given the common early phylogenetic origin of the order's species and their likely broad similarity in immunopathological responses to ocular injury or insult, any approach to the diagnosis and therapy of ocular disease should be based upon protocols used at present in the

domestic horse (*E. caballus*) (Gilger 2017). In the case of captive tapirs and possibly rhinoceros, the clinician should be alert to the potential role of UV exposure in the genesis of chronic destructive keratitis, as described in captive marine pinnipeds in the USA (Miller et al. 2013). In any event, the principles guiding the clinical management of the acute ocular disease are:

- Prompt and accurate diagnosis of the generic nature of the disease process and the ocular tissues involved in that process.
- Where appropriate, identify any extraneous causal agent, e.g., fungi or other microbial pathogens, foreign material. This may involve microbial sampling, biopsy, or aspiration of intraocular content (Stoppini and Gilger 2017; Dwyer 2017).
- Formulate and implement an effective therapeutic strategy targeting the affected tissue and directed towards the elimination of any causal agent or other impediments to healing, e.g., foreign body, necrotic corneal or other tissue, or towards suppression of aberrant or dysregulated immunoinflammatory responses, e.g., keratolysis and ulceration driven by innate hydrolases derived from leucocytes and macrophages sequestered on the ocular surface.
- Promote ocular repair, directed towards restoring the physical integrity and normal function of the injured ocular tissues.

In the case of most non-domestic species, the above must be accomplished within the constraints imposed by limited patient contact and compliance and the perceived risks to

the patient of multiple general anesthesia events. In circumstances of good management facilities and relatively tractable patients, effective topical therapy may be possible via a transpalpebral lavage system allowing remote administration of the agent. However, in many cases, effective topical therapy typically requires short interval dosing, which may limit its suitability for use. Continuous delivery pumps can be used in conjunction with subpalpebral lavage systems and may have a benefit in enhancing the bioavailability of some topical medication such as hydrolase inhibitor preparation in corneal ulceration. However, some medications, such as ophthalmic suspensions (e.g., natamycin), or drugs that require refrigeration (e.g., voriconazole, cefazoline), are not suitable to be used in continuous delivery systems, and the clinician should consider this where such systems are considered.

With the probable exception (in some instances) of the inflamed uveal tract or ocular surface, most systemically administered therapeutic agents, including those administered orally, cannot be relied upon alone to achieve and maintain therapeutic levels within the eye or on the ocular surface (Matthews 2004, 2009). This necessitates early consideration of the use of depot formulated drug preparations, where appropriate and where available, administered by local administration techniques such as subconjunctival, intracorneal, or intracameral injection. Consideration should also be given to the use of slow-release embedded devices, e.g., suprachoroidal or episcleral Cyclosporine A implants, in managing immune-mediated uveal or corneal disease, or cisplatin beads as adjunctive therapy in managing periocular tumors.

When addressing a severe corneal wound condition (e.g., keratomalacia, deep stromal ulcer, descemetocoele), early recourse to single intervention conjunctival grafting surgeries (Brooks et al. 2017) in managing corneal ulceration is likely to accelerate corneal healing and repair and may preserve the eye, albeit at the expense of end-stage corneal clarity, and should be a primary consideration in any therapeutic strategy.

References

- Archie EA (2013) Wound healing in the wild: stress, sociality and energetic costs affect wound healing in natural populations. *Parasite Immunol* 35(11):374–385
- Banks MS, Sprague WW, Schmoll J et al (2015) Why do animal eyes have pupils of different shapes? *Sci Adv*. <https://doi.org/10.1126/sciadv.1500391>
- Bapodra P, Wolfe BA (2014) Baseline assessment of ophthalmic parameters in the greater one-horned rhinoceros (*Rhinoceros unicornis*). *J Zoo Wildl Med* 45:859–865
- Brooks D, Matthews A, Clode A (2017) Diseases of the equine cornea (Chapter 7). In: Gilger B (ed) *Equine ophthalmology*, 3rd edn. Wiley Blackwell, Philadelphia, pp 1252–1368
- Coimbra JP, Manger PR (2017) Retinal ganglion cell topography and spatial resolving power in the white rhinoceros (*Ceratotherium simum*). *J Comp Neurol* 525:2484–2498
- Da Silva M-AO, Hermoza C, Rojas G, Freundt JM (2013) Identifying an effective treatment for corneal ulceration in captive tapirs. *Tapir Conserv* 22:12–14
- Dwyer A (2017) Practical field ophthalmology (Chapter 3). In: Gilger B (ed) *Equine ophthalmology*, 3rd edn. Wiley Blackwell, Philadelphia, pp 72–111
- Ehrenhofer MCA, Deeg CA, Reese S et al (2002) Normal structure and age-related changes of the equine retina. *Vet Ophthalmol* 5:39–47
- Esson DW, Wellehan JFX, Lafortune M et al (2006) Surgical management of a malacic corneal ulcer in a greater one-horned Asian rhinoceros (*Rhinoceros unicornis*) using a free island tarsocconjunctival graft. *Vet Ophthalmol* 9:65–69
- Evans AL, Carter RT, Marlar AB, Citino SB (2009) Determination of the appropriate size intraocular lens for cataract surgery in the Grevy's zebra (*Equus Grevyi*). In: Proceedings AAZV AAWV Joint Conference
- Francois J, Neetens A (1962) Vascularisation of the retino-oculic block. In: Davson H (ed) *Vegetative physiology and biochemistry: the eye*, vol 1. Academic Press, New York, pp 372–380
- Gandolf AR, Willis AM, Blumer ES, Atkinson MW (2000) Melting corneal ulcer management in a greater one-horned rhinoceros (*Rhinoceros unicornis*). *J Zoo Wildl Med* 31:112–117
- García MJ, Medici EP, Naranjo EJ et al (2012) Distribution, habitat and adaptability of the genus tapirus. *Integr Zool* 7:346–355
- Gerkema MP, Davies WL, Foster RG et al (2013) The nocturnal bottleneck and the evolution of activity patterns in mammals. *Proc R Soc B Biol Sci* 280:20130508. <https://doi.org/10.1098/rspb.2013.0508>
- Gilger BC (2017) *Equine ophthalmology*, 3rd edn. Wiley Blackwell, Philadelphia
- Gilger BC, Hollingsworth SR (2017) Diseases of the uvea, uveitis, and recurrent uveitis. In: *Equine ophthalmology*, 3rd edn. Wiley Blackwell, Philadelphia
- Grininger P, Skalicky M, Nell B (2010) Evaluation of healthy equine eyes by use of retinoscopy, keratometry, and ultrasonographic biometry. *Am J Vet Res* 71:677–681
- Holmes EC, Ellis SA (1999) Evolutionary history of MHC class I genes in the mammalian order Perissodactyla. *J Mol Evol* 49:316–324
- Horowitz IH, Dubielzig RR, Botero-Anug AM et al (2016) Conjunctival habronemiasis in a square-lipped rhinoceros (*Ceratotherium simum*). *Vet Ophthalmol* 19:161–166
- Howland HC, Rowland M, Murphy CJ (1993) Refractive state of the rhinoceros. *Vision Res* 33:2649–2651
- Johnson GL (1901) I. Contributions to the comparative anatomy of the mammalian eye, chiefly based on ophthalmoscopic examination. *Philos Trans R Soc Lond Ser B*. [https://doi.org/10.1016/1352-2310\(95\)00219-7](https://doi.org/10.1016/1352-2310(95)00219-7)
- Karpinski LG, Miller CL (2002) Fluorouracil as a treatment for corneal papilloma in a Malayan tapir. *Vet Ophthalmol* 5:241–243
- Kenny DE, Dugan SJ, Knightly F, Baier J (2003) Intracapsular lens removal in a Przewalski's wild horse (*Equus caballus przewalskii*). *J Zoo Wildl Med* 34:284–286
- Kumar A, Suryadevara N, Hill TM et al (2017) Natural killer T cells: an ecological evolutionary developmental biology perspective. *Front Immunol* 8(1858):1–19
- Martin LB, Hopkins WA, Mydlarz LD, Rohr JR (2010) The effects of anthropogenic global changes on immune functions and disease resistance. *Ann N Y Acad Sci* 1195:129–148
- Matthews AG (2004) Ophthalmic therapeutics (Chapter 13). In: Bertone JJ, Horspool LJ (eds) *Equine clinical pharmacology*. WB Saunders, Philadelphia, pp 217–246

- Matthews AG (2009) Ophthalmic antimicrobial therapy in the horse. *Equine Vet Educ* 21:271–280
- Miller S, Colitz CM, St Leger J, Dabielzig R (2013) A retrospective survey of the ocular histopathology of the pinniped eye with emphasis on corneal disease. *Vet Ophthalmol* 16:119–129
- Montiani-Ferreira F (2001) Ophthalmology. In: Fowler M, Cubas Z (eds) *Biology, medicine and surgery of South American wild animals*. Iowa State University Press, Ames, pp 437–456
- Muraille E (2014) Generation of individual diversity: a too neglected fundamental property of adaptive immune system. *Front Immunol* 5(208):1–7
- Ofri R, Horowitz IH, Kass PH (1998) Tonometry in three herbivorous wildlife species. *Vet Ophthalmol* 1:21–24
- Peichl L (2005) Diversity of mammalian photoreceptor properties: adaptations to habitat and lifestyle? *Anat Rec A Discov Mol Cell Evol Biol* 287A:1001–1012
- Pettigrew JD, Manger PR (2008) Retinal ganglion cell density of the black rhinoceros (*Diceros bicornis*): calculating visual resolution. *Vis Neurosci* 25:215–220
- Sandmann D, Boycott BB, Peichl L (1996) Blue-cone horizontal cells in the retinae of horses and other equidae. *J Neurosci* 16:3381–3396
- Steiner CC, Ryder OA (2011) Molecular phylogeny and evolution of the Perissodactyla. *Zool J Linnean Soc.* <https://doi.org/10.1111/j.1096-3642.2011.00752.x>
- Stoppini R, Gilger B (2017) Equine ocular examination basic techniques (Chapter 1). In: Gilger B (ed) *Equine ophthalmology*, 3rd edn. Wiley Blackwell, Philadelphia, pp 1–39
- Trifonov VA, Stanyon R, Nesterenko AI et al (2008) Multidirectional cross-species painting illuminates the history of karyotypic evolution in Perissodactyla. *Chromosome Res* 16:89–107
- Ujvari B, Klaassen M, Raven N et al (2018) Genetic diversity, inbreeding and cancer. *Proc R Soc B* 285:20172589. <https://doi.org/10.1098/rspb.2017.2589>
- Veilleux CC, Kirk EC (2014) Visual acuity in mammals: effects of eye size and ecology. *Brain Behav Evol* 83:43–53
- Veilleux CC, Lewis RJ (2011) Effects of habitat light intensity on mammalian eye shape. *Anat Rec (Hoboken)* 49:316–324