

*Int. Zoo Yb.* (2007) **41**: 52–74  
 DOI:10.1111/j.1748-1090.2007.00017.x

## The many uses of non-invasive faecal steroid monitoring in zoo and wildlife species

F. SCHWARZENBERGER

*Department of Natural Sciences, Institute of Biochemistry, University of Veterinary Medicine, Vienna, Austria*

*E-mail: Franz.Schwarzenberger@vu-wien.ac.at*

During the past two decades, techniques for faecal steroid analysis have been developed and have been used for research with mammalian, bird and, to a lesser extent, reptile, amphibian and fish species. Various techniques for the analysis of reproductive (oestrogen, androgen, progesterone) and adrenocortical (glucocorticoids) steroid hormones have been established and have been applied to a wide range of research questions studying captive and free-ranging wildlife, as well as domestic and laboratory species. Because of species-specific differences in steroid metabolism in even closely related species, careful validation of assay methods is necessary in order to generate meaningful and accurate results. For future research and management of free-ranging and captive wildlife, the great potential of non-invasive endocrine monitoring will be utilized more than ever. In light of this, captive wildlife species are ideal research subjects, as longitudinal sample collection is possible and studies connecting physiology, endocrinology, reproduction and stress with various social and/or environmental factors can be carried out and can be analysed to determine how they impact animal health.

*Key-words:* adrenal steroid hormones; assay validation; biodiversity conservation; excreted steroids; faecal steroid analysis; non-invasive; reproductive steroid hormones; steroid metabolism.

### INTRODUCTION

During the past two decades, techniques for faecal steroid analysis (oestrogen, androgen, progesterone and glucocorticoid metabolites) have been developed and have been used for research with mammals (for references, see Table 1), birds (i.e. Baltic *et al.*, 2005; Goymann, 2005; Möstl *et al.*, 2005; Palme *et al.*, 2005; Wasser & Hunt, 2005; Szoke *et al.*, 2006) and, to a lesser extent, reptile (i.e. Atkins *et al.*, 2002; Rittenhouse *et al.*, 2005), amphibian (Szymanski *et al.*, 2006) and fish (i.e. Turner *et al.*, 2003; Ellis *et al.*, 2004;

Wysocki *et al.*, 2006) species. Techniques have been applied to a wide range of research questions studying captive and free-ranging wildlife, as well as domestic and laboratory species. These techniques are now widely accepted and several reviews focusing on steroid metabolism and on the validation of faecal steroid assays have been published recently (Schwarzenberger *et al.*, 1996, 1997; Whitten *et al.*, 1998; Möstl & Palme, 2002; von der Ohe & Servheen, 2002; Monfort, 2003; Wielebnowski, 2003; Buchanan & Goldsmith, 2004; Graham, 2004; Millspaugh & Washburn, 2004; Möstl *et al.*, 2005; Palme, 2005; Palme *et al.*, 2005; Touma & Palme, 2005; Ziegler & Wittwer, 2005; Keay *et al.*, 2006; Lane, 2006). In addition, reviews emphasizing the importance of faecal steroid analysis for the elucidation of endocrine physiology in certain taxa (carnivores: Young, K. M. *et al.*, 2004; Brown, 2006; primates: Shimizu, 2005; rhinoceros: Roth, 2006), and as an important tool for the management and conservation of wildlife (Pukazhenti & Wildt, 2004; Cockrem, 2005; Andrabi & Maxwell, 2007), or as reliable indicators of stress (Wasser *et al.*, 2000; Möstl & Palme, 2002; Millspaugh & Washburn, 2004; Queyras & Carosi, 2004; Touma & Palme, 2005; Keay *et al.*, 2006; Lane, 2006) have been published.

The aim of this review is briefly to summarize studies using faecal steroid analysis in captive and free-ranging mammalian wildlife species. The focus is on recently published literature summarizing reproductive (oestrogens, androgens and progesterones)

TAXA/SPECIES	APPLICATION (WHAT HAS BEEN STUDIED?)	CAPTIVE/ FREE-RANGING (WILD), ♂/♀	HORMONES DETERMINED IN FAECES	REFERENCES
<b>PRIMATES</b>				
<b>HOMINIDAE</b>				
Chimpanzee <i>Pan troglodytes</i>	oestrous cycle, sexual swelling dominance rank, challenge hypothesis immune endocrine interactions, intestinal parasites social rehabilitation of former laboratory chimpanzees, animal welfare	captive, ♀ wild, ♂ wild, ♂ & ♀	E, P A A, GC	Emery & Whitten (2003) Muehlenbein <i>et al.</i> (2004) Muehlenbein (2006)
Lowland gorilla <i>Gorilla gorilla</i>	reproductive senescence; oestrous cycle, sexual behaviour	captive, ♂	GC	Reimers <i>et al.</i> (2007)
<b>HYLOBATIDAE</b> White-handed gibbon <i>Hylobates lar</i>	oestrous cycle, sexual swelling, ovulation, paternity confusion	wild, ♀	E, P	Atsalis <i>et al.</i> (2004)
<b>OLD WORLD MONKEYS</b>				
<b>CERCOPITHECIDAE</b>				
Baboon <i>Papio cynocephalus</i>	peripartum physiology (life-history stages) sexual maturation (infancy, puberty); oestrous cycle, sexual swelling and consortship pregnancy, foetal loss	wild, ♀ wild, ♂ & ♀ wild, ♀	E, A, P, GC E, A, P, GC E, P, GC	Altmann <i>et al.</i> (2004) Gesquiere <i>et al.</i> (2005, 2007) Beehner <i>et al.</i> (2006)
Chama baboon <i>Papio hamadryas ursinus</i>	seasonal reproduction and environmental stressors ♂ hierarchy, immigrant ♂, aggression, stress confusion	wild, ♂ & ♀ wild, ♂ wild, ♀	GC GC P	Weingrill <i>et al.</i> (2004) Bergman <i>et al.</i> (2005) Heistermann <i>et al.</i> (2001)
Hanuman langur <i>Semnopithecus entellus</i>	oestrous cycle, concealed ovulation, paternity confusion	wild, ♀	P	
Japanese macaque <i>Macaca fuscata</i>	oestrous cycle, conception, paired urine and faecal samples social rank, aggression, stress peripartum physiology (maternal style) reproductive behaviour	captive & wild, ♀ wild, ♂ captive, ♀ semi-wild, ♀	E, P A, GC E, P E, P	Fujita <i>et al.</i> (2001) Barrett <i>et al.</i> (2002) Bardi <i>et al.</i> (2003) O'Neill <i>et al.</i> (2004)
Long-tailed macaque <i>Macaca fascicularis</i>	oestrous cycle, sexual swelling, post-conception mating, paternity confusion	wild, ♀	E, P	Engelhardt <i>et al.</i> (2007)

Table 1. Continued

TAXA/SPECIES	APPLICATION (WHAT HAS BEEN STUDIED?)	CAPTIVE/ FREE-RANGING (WILD), ♂/♀	HORMONES DETERMINED IN FAECES	REFERENCES
Red colobus <i>Ptilocolobus badius</i>	food availability, parasite infections, adrenal activity	wild, ♂ & ♀	GC	Chapman <i>et al.</i> (2006)
Red-shanked douc langur <i>Pygathrix nemaeus</i>	oestrous cycle, change in group composition	captive, ♀	E, P, GC	Heistermann <i>et al.</i> (2004)
Rhesus macaque <i>Macaca mulatta</i>	peripartum physiology (maternal style)	captive, ♀	E, P	Bardi <i>et al.</i> (2003)
Stump-tailed macaque <i>Macaca arctoides</i>	♂ reproductive behaviour in response to vaginal secretions	captive, ♀	E	Cerdas-Molina <i>et al.</i> (2006)
NEW WORLD MONKEYS				
ATELIDAE				
Black-handed spider monkey <i>Ateles geoffroyi</i>	oestrous cycles, foetal loss; paired urine and faecal samples	captive & wild, ♀	E, P	Campbell <i>et al.</i> (2001)
Mantled howler monkey <i>Alouatta palliata</i>	artificial insemination, vaginal cytology	captive, ♀	E, P	Hernández-López <i>et al.</i> (2007)
Muriqui monkey <i>Brachyteles arachnoides</i>	challenge hypothesis, aggression, mating system	wild, ♂	A	Cristobal-Azkarate <i>et al.</i> (2006)
Common marmoset <i>Callithrix jacchus</i>	seasonality, reproductive behaviour, oestrous cycle	wild, ♂ & ♀	E, A, P, GC	Strier <i>et al.</i> (2003)
Golden lion tamarin <i>Leontopithecus rosalia</i>	reproductive suppression, parent-daughter relations, oestrous cycle	captive, ♀	E, P, GC	Ziegler & Sousa (2002)
Tufted capuchin <i>Cebus apella nigritus</i>	social rank, challenge hypothesis	wild, ♂	A, GC	Bales <i>et al.</i> (2006)
White-faced capuchin <i>Cebus capucinus</i>	seasonality, reproductive behaviour, aggression	wild, ♂	A, GC	Lynch <i>et al.</i> (2002)
LEMURIFORMES	reproductive behaviour, oestrous cycles, concealed ovulation, paternity confusion	wild, ♀	E, P	Carnegie <i>et al.</i> (2005)
INDRIDAE	social rank, sexual behaviour, seasonality	wild, ♂	A	Brockman <i>et al.</i> (2001)
Verreaux's sifaka <i>Propithecus verreauxi</i>				

LEMURIDAE Red-fronted lemur <i>Eulemur fulvus rufus</i>	seasonality, aggression, challenge hypothesis pregnancy, foetal sex determination	wild, ♂ & ♀	E, A, P	Ostner <i>et al.</i> (2002), Ostner & Heistermann (2003)
Ring-tailed lemur <i>Lemur catta</i>	social rank, sexual behaviour, aggression, seasonality adrenal activity as predictor of mortality	wild, ♂ & ♀ wild, ♂ & ♀	GC GC	Cavigelli <i>et al.</i> (2003), Gould <i>et al.</i> (2005) Pride (2005)
<b>MARSUPIALS</b>				
DASYURIDAE Chuditch <i>Dasyurus geoffroii</i>	oestrous cycle, vaginal cytology	captive, ♀	E	Stead-Richardson <i>et al.</i> (2001)
MACROPODIDAE Tamar wallaby <i>Macropus eugenii</i>	animal well-being, immune function, paired plasma and faecal samples	captive, ♀	GC	McKenzie & Deane (2005)
PETAURIDAE Squirrel glider <i>Petaurus norfolcensis</i>	oestrous cycle, pregnancy, seasonality, urinary cytology	captive, ♀	E, P	Woodd <i>et al.</i> (2006)
TACHYGLOSSIDAE Short-beaked echidna <i>Tachyglossus aculeatus</i>	sex identification, oestradiol:androgen ratio; hCG stimulation, seasonality, sperm production	captive, ♂ & ♀	E, A	Oates <i>et al.</i> (2002), Johnston <i>et al.</i> (2007)
TARSIPEIDIDAE Honey possum <i>Tarsipes rostratus</i>	radiometabolism study, oestrous cycle, embryonic diapause, seasonality, reproductive stress	captive, ♂ & ♀	E, P, GC	Bradshaw <i>et al.</i> (2004), Oates <i>et al.</i> (2004, 2007)
VOMBATIDAE Common wombat <i>Vombatus ursinus</i> Southern hairy-nosed wombat <i>Lasiorhinus latifrons</i>	oestrous cycle, seasonality, sexual behaviour, paired blood and faecal samples	captive, ♀	P	Paris <i>et al.</i> (2002)
<b>PILOSA</b>				
BRADYPODIDAE Three-toed sloth <i>Bradypus variegatus</i>	oestrous cycle, pregnancy, paired blood-faecal samples	captive, ♀	E, P	Mühlbauer <i>et al.</i> (2006)
MYRMECOPHAGIDAE Giant anteater <i>Myrmecophaga tridactyla</i>	oestrous cycle, pregnancy, post-partum period, aging ♀	captive, ♀	E, P	Patzl <i>et al.</i> (1998)

Table 1. Continued

TAXA/SPECIES	APPLICATION (WHAT HAS BEEN STUDIED?)	CAPTIVE/ FREE-RANGING (WILD), ♂/♀	HORMONES DETERMINED IN FAECES	REFERENCES
<b>RODENTIA</b>				
<b>CHINCHILLIDAE</b>				
Chinchilla <i>Chinchilla lanigera</i>	radiometabolism study, pregnancy, seasonality	captive, ♀	E, P	Busso <i>et al.</i> (2007)
<b>CRICETIDAE</b>				
Syrian hamster <i>Mesocricetus auratus</i>	oestrous cycle, superovulation	captive, ♀	E, P	Chelimi <i>et al.</i> (2005)
<b>ERETHIZONTIDAE</b>				
Common porcupine <i>Erethizon dorsatum</i>	oestrous cycle, pregnancy, pseudopregnancy	captive, ♀	E, P	Bogdan & Monfort (2001)
<b>MURIDAE</b>				
Midday gerbil <i>Meriones meridianus</i>	population density, habitat type and ecological indices, adrenal activity	wild, ♂ & ♀	GC	Kuznetsov <i>et al.</i> (2004)
<b>SCIURIDAE</b>				
Cape ground squirrel <i>Xerus inauris</i>	pregnancy, assay validation	captive, ♀	E, P	Pettitt <i>et al.</i> (2007)
<b>LAGOMORPHA</b>				
<b>LEPORIDAE</b>				
European rabbit <i>Oryctolagus cuniculus</i>	physiological stress response, odour and presence of a predator, ACTH challenge long-term stress and fitness, body condition, survival probability of wild animals put into captivity	captive, ♂ & ♀ wild, ♂ & ♀	GC	Monclus <i>et al.</i> (2006), Cabezas <i>et al.</i> (2007)
<b>SIRENIA</b>				
<b>DUGONGIDAE</b>				
Dugong <i>Dugong dugon</i>	assessment of reproductive status	wild, ♂ & ♀	E, A	Lanyon <i>et al.</i> (2005)
<b>CETACEA</b>				
<b>BALAEINIDAE</b>				
North Atlantic right whale <i>Eubalaena glacialis</i>	adrenal activity, reproductive rank	wild, ♂ & ♀	E, P, A, GC	Rolland <i>et al.</i> (2005), Hunt <i>et al.</i> (2006)

<b>PROBOSCIDEA</b>									
<b>ELEPHANTIDAE</b>									
African elephant									
<i>Loxodonta africana</i>	oestrous cycle, pregnancy; paired faecal and urine samples	captive, ♀	E, P						Fiess <i>et al.</i> (1999)
	muth, intensity, duration, behaviour, sexual activity	captive & wild, ♂	A, GC						Ganswindt, Heistermann & Hodges (2005), Ganswindt, Rasmussen <i>et al.</i> (2005)
	ecological conditions (vegetation) and reproduction	wild, ♀	P						Wittmyer <i>et al.</i> (2007)
<b>PERISSODACTYLA</b>									
<b>EQUIDAE</b>									
Grevy's zebra	oestrous cycle, pregnancy, abortions	captive, ♀	E, P						Asa <i>et al.</i> (2001)
<i>Equus grevyi</i>	oestrous cycle, pregnancy, sexual behaviour	semi-wild, ♀	P						Scheibe <i>et al.</i> (1999)
Przewalski's horse	oestrous cycle, pregnancy, comparative study plasma and faeces, regumate treatment to prevent abortion	captive, ♀	E, P						Berkeley <i>et al.</i> (1997)
<i>Equus ferus przewalskii</i>	ultrasonography, oestrous cycles, pregnancy identification of faecal steroid metabolites, pregnancy, gas-liquid chromatography-mass spectrometry	captive, ♀	P						Radcliffe <i>et al.</i> (2001)
<b>RHINOCEROTIDAE</b>									
Black rhinoceros	oestrous cycle, seasonality, foetal loss, behavioural observation	wild, ♀	P						Lance <i>et al.</i> (2001)
<i>Diceros bicornis</i>	comparative study; oestrous cycle, pregnancy, testicular, adrenal activity	captive, ♂ & ♀	E, P, A, GC						Garnier <i>et al.</i> (2002)
White rhinoceros	comparative study; adrenal activity, environmental factors, matched plasma, urine and faeces	captive, ♂ & ♀	GC						Brown <i>et al.</i> (2001)
<i>Ceratotherium simum</i>	comparative study; adrenal activity, behavioural observation	captive, ♀	GC						Turner <i>et al.</i> (2002)
Black rhinoceros	comparative study; adrenal activity, behavioural, environmental factors, reproduction	captive, ♀	GC						Carlstead & Brown (2005)
<i>Diceros bicornis</i>	pregnancy diagnosis, embryonic loss, reproductive failure, infertility, genital health, ultrasonography	captive, ♀	P						Radcliffe <i>et al.</i> (1997), Hermes <i>et al.</i> (2006)
White rhinoceros	reproductive behaviour, territoriality, environmental factors, GnRH application	wild, ♂	A						Rachlow <i>et al.</i> (1998), Kretzschmar <i>et al.</i> (2004)
<i>Ceratotherium simum</i>	oestrous cycle, pregnancy, parturition, reproductive behaviour	captive, ♀	P						Schwarzenberger <i>et al.</i> (1998), Patton <i>et al.</i> (1999)
	pregnancy, parturition, vaginal electrical impedance	captive, ♀	E, P, GC						Bowers <i>et al.</i> (2005)

Table 1. Continued

TAXA/SPECIES	APPLICATION (WHAT HAS BEEN STUDIED?)	CAPTIVE/ FREE-RANGING (WILD), ♂/♀	HORMONES DETERMINED IN FAECES	REFERENCES
Indian rhinoceros <i>Rhinoceros unicornis</i>	oestrous cycle, pregnancy, post-partum period	captive, ♀	E, A, P	Schwarzenberger <i>et al.</i> (2000)
Sumatran rhinoceros <i>Dicerorhinus sumatrensis</i>	oestrous cycle, pregnancy, paired blood and faecal samples, ultrasonographic monitoring, regumate treatment to prevent abortion	captive, ♀	P	Roth <i>et al.</i> (2001, 2004)
<b>ARTIODACTYLA</b>				
<b>BOVIDAE</b>				
Arabian oryx <i>Oryx leucoryx</i>	oestrous cycle, pregnancy, post-partum period	captive, & wild, ♀	E, P	Ostrowski <i>et al.</i> (2005)
Highorn sheep <i>Ovis montanus</i>	social rank, reproductive behaviour	wild, ♂	A	Pelletier <i>et al.</i> (2003)
Bison <i>Bison bison</i>	pregnancy diagnosis, radiocollared animals	wild, ♀	E, P	Schoenecker <i>et al.</i> (2004)
	rut, dominance, stress, mating success, reproductive behaviour	semi-wild, ♂	A, GC	Mooring <i>et al.</i> (2004, 2006)
	seasonality, oestrous cycle, post-partum cycle, pregnancy, reproductive behaviour	semi-wild, ♀	P	Vervaecke & Schwarzenberger (2006)
Buffalo <i>Bubalus bubalis</i>	ovulation induction, oestrous synchronization, norgestomet implant,	captive, ♀	P	Hattab <i>et al.</i> (2000)
Chamois <i>Rupicapra rupicapra</i>	gastrointestinal and lung helminths, ♂-biased parasitism, endocrine-immune interaction	wild, ♂ & ♀	E, A, GC	Hoby <i>et al.</i> (2006)
Fringe-eared oryx or Gemsbok <i>Oryx gazella callotis</i>	aggression control, bachelor herd, behavioural observation	captive, ♂	A	Patton <i>et al.</i> (2001)
Gerenuk <i>Litocranius walleri walleri</i>	oestrous cycle, seasonality, oestrous synchronization, artificial insemination, pregnancy	captive & wild, ♂ & ♀	A, P	Penfold <i>et al.</i> (2005)
Mhorr gazelle <i>Gazella dama mhorr</i>	oestrous cycle, pregnancy, oestrous synchronization	captive, ♀	P	Pickard <i>et al.</i> (2001)
Rocky mountain goat <i>Ovis canadensis canadensis</i>	faecal parasite loads, seasonality, radiotelemetry	wild, ♂ & ♀	GC	Goldstein <i>et al.</i> (2005)

Rocky mountain goat <i>Oreamnos americanus</i>	reproductive physiology, seasonality, pregnancy	captive, ♀	P	Kallert <i>et al.</i> (2002)
Sable antelope <i>Hippotragus niger</i>	oestrous cycle, seasonality, oestrous synchronization	captive, ♀	E, P	Thompson & Monfort (1999), Thompson <i>et al.</i> (1998)
Scimitar-horned oryx <i>Oryx dammah</i>	oestrous cycle, seasonality, pregnancy, oestrous synchronization, artificial insemination, ultrasonography	captive, ♀	E, P	Morrow <i>et al.</i> (1999, 2000)
CERVIDAE				
Brown brocket deer <i>Mazama gouazoubira</i>	oestrous cycle, pregnancy, oestrous behaviour	captive, ♀	P	Pereira, Polegato <i>et al.</i> (2006)
Chinese water deer <i>Hydropotes inermis</i>	oestrous cycle, seasonality, reproductive behaviour	captive, ♂ & ♀	P, A	Mauget <i>et al.</i> (2007)
Elk <i>Cervus elaphus canadensis</i>	pregnancy, fecundity free-ranging herd, nutrition associated abortion, body condition	wild, ♀	E, P	Stoops <i>et al.</i> (1999), Cook <i>et al.</i> (2002)
Red deer <i>Cervus elaphus</i>	seasonal variation: faecal samples were partly genotyped for individual and sex identification	captive, ♂ & ♀	GC	Huber <i>et al.</i> (2003)
Pampas deer <i>Ozotoceros bezoarticus bezoarticus</i>	seasonality, reproductive behaviour, antler cycle, human disturbance	wild, ♂	A, GC	Pereira <i>et al.</i> (2005), Pereira, Barbanti Duarte & Negrao (2006)
Pere David's deer <i>Elaphurus davidianus</i>	♂ and ♀ reproductive behaviour	wild, ♂ and ♀	E, A, P	Li <i>et al.</i> (2001)
Reindeer <i>Rangifer tarandus</i>	welfare and adrenal activity, comparison of GC in blood, saliva, urine, faeces with IgA	captive, ♀	GC	Rehbinder & Hau (2006)
Sika deer <i>Cervus nippon yesoensis</i>	seasonality, oestrous cycle, conception	captive, ♀	P	Matsuura <i>et al.</i> (2004)
Southern pudu <i>Pudu pudu</i>	oestrous cycle, pregnancy, seasonality	captive, ♀	P	Blanvillain <i>et al.</i> (1997)
White-tailed deer <i>Odocoileus virginianus</i>	oestrous cycle, pregnancy, assay validation through application of oestradiol and progesterone by blow gun	captive, ♀	E, P	Kapke <i>et al.</i> (1999)
	faecal-steroid monitoring of immunococontraception	wild, ♀	P	Walter <i>et al.</i> (2002)



Table 1. Continued

TAXA/SPECIES	APPLICATION (WHAT HAS BEEN STUDIED?)	CAPTIVE/ FREE-RANGING (WILD), ♂/♀	HORMONES DETERMINED IN FAECES	REFERENCES
<b>GIRAFFIDAE</b>				
Giraffe <i>Giraffa camelopardalis</i>	oestrous cycle, post-partum period, pregnancy, behaviour; monitoring contraception (GnRH agonist implant)	captive & semi-wild, ♀	P	Del Castillo <i>et al.</i> (2005), Dumonceaux <i>et al.</i> (2006), Patton <i>et al.</i> (2006)
Okapi <i>Okapia johnstoni</i>	reproductive physiology, oestrous cycle, pregnancy, post partum period, abortion, regumate treatment to prevent abortion	captive, ♀	P	Schwarzenberger <i>et al.</i> (1999)
<b>HIPPOTAMIDAE</b>				
Hippopotamus <i>Hippopotamus amphibius</i>	oestrous cycle, pregnancy, puberty, lactational anoestrous, miscarriage	captive, ♀	P	Graham <i>et al.</i> (2002), Wheaton <i>et al.</i> (2006)
<b>SUIDAE</b>				
Babirusa <i>Babirusa babirusa</i>	oestrous cycle, pregnancy, seasonality, comparative study on reproduction in suidae	captive, ♀	E, A, P	Berger <i>et al.</i> (2006)
Common warthog <i>Phacochoerus africanus</i>				
Red river hog <i>Potamochoerus porcus</i>				
<b>CARNIVORA</b>				
<b>AILURIDAE</b>				
Red panda <i>Ailuurus fulgens fulgens</i>	seasonality, oestrous cycle, induced ovulation, pregnancy diagnosis, pseudopregnancy, captive management (group size)	captive, ♀	E, P, A	Spanner <i>et al.</i> (1997), MacDonald <i>et al.</i> (2005), Wei <i>et al.</i> (2005)
<b>CANIDAE</b>				
African wild dog <i>Lycan pictus</i>	social rank, reproductive suppression, social stress	wild, ♂ & ♀	E, P, A, GC	Creel <i>et al.</i> (1997)
Bush dog <i>Speothos venaticus</i>	oestrous cycle, seasonality, ♂ effect	captive, ♂ & ♀	E, P, A	DeMatteo <i>et al.</i> (2006)
Fennee fox <i>Vulpes zerda</i>	oestrous cycle, vulval swelling, blood and faeces	captive, ♀	E, P	Valdespino <i>et al.</i> (2002)
Gray wolf <i>Canis lupus</i>	oestrous induction; GnRH implant (deslorelin) social rank, reproductive suppression, social stress	captive, ♀ wild, ♂ & ♀	E, P GC	Asa <i>et al.</i> (2006) Sands & Creel (2004)

Maned wolf <i>Chrysocyon brachyurus</i>	seasonality, oestrous cycle, pregnancy, pseudopregnancy, infusion of radiolabelled testosterone	captive, ♂ & ♀	E, P, A	Velloso <i>et al.</i> (1998)
Red wolf <i>Canis rufus</i>	seasonality, oestrous cycle, pregnancy, pseudopregnancy, blood samples; longitudinal study of adrenal activity	captive, ♂ & ♀	E, P, A, GC	Walker <i>et al.</i> (2002), Young, K. M. <i>et al.</i> (2004)
FELIDAE				
Cheetah <i>Acinonyx jubatus</i>	adrenal activity, chronic stress, adrenal gland morphology; social management, movement between facilities	captive & wild, ♂ & ♀	E, A, GC	Wielebnowski, Ziegler <i>et al.</i> (2002), Terio <i>et al.</i> (2004), Wells <i>et al.</i> (2004), Young, K. M. <i>et al.</i> (2004)
Clouded leopard <i>Neofelis nebulosa</i>	oestrous cycle, pregnancy, seasonality, ovulation induction	captive, ♀	E, P	Brown <i>et al.</i> (1995)
	environmental stressor related to captive management; ACTH challenge	captive, ♂ & ♀	GC	Wielebnowski, Fletchall <i>et al.</i> (2002), Young, K. M. <i>et al.</i> (2004)
	ovulation induction, GnRH agonist and gonadotrophin injection, artificial insemination	captive, ♀	E, P	Pelican <i>et al.</i> (2006)
Eurasian lynx <i>Lynx lynx</i>	seasonality, ultrasonography, electroejaculation, spermatogenesis	captive, ♂	A	Göritz <i>et al.</i> (2006)
Jaguar <i>Panthera onca</i>	restraint and electroejaculation	captive, ♂	GC	Morato <i>et al.</i> (2004)
Pallas' cat <i>Otocolobus manul</i>	oestrous cycle, pregnancy, seasonality, ovulation induction, artificial insemination	captive, ♂ & ♀	E, P, A	Brown <i>et al.</i> (2002)
Tiger <i>Panthera tigris</i>	ovulation induction, different doses of gonadotrophins, artificial insemination	captive, ♂	E, P	Graham <i>et al.</i> (2006)
HERPESIIDAE				
Meerkats <i>Suricata suricatta</i>	oestrous cycles, pregnancy, post-partum period; social rank, cooperative breeding, stress, faecal and urine samples	wild, ♂ & ♀	E, P, A, GC	Moss <i>et al.</i> (2001), Young, K. M. <i>et al.</i> (2004), Young, A. J. <i>et al.</i> (2006)
HYAENIDAE				
Spotted hyena <i>Crocuta crocuta</i>	social rank, stress social rank, pregnancy, aggression, rank-related maternal effects	wild, ♂ & ♀ wild, ♀	GC A	Goymann <i>et al.</i> (2001, 2003) Dloniak <i>et al.</i> (2006)

Table 1. Continued

TAXA/SPECIES	APPLICATION (WHAT HAS BEEN STUDIED?)	CAPTIVE/ FREE-RANGING (WILD), ♂/♀	HORMONES DETERMINED IN FAECES	REFERENCES
MUSTELIDAE Black-footed ferret <i>Mustela nigripes</i>	seasonality, oestrous cycles, pregnancy, pseudopregnancy, vaginal lavage, vulva size, litter size correlation with faecal oestradiol; adrenal activity, infusion of radiolabelled cortisol and corticosterone, ACTH challenge	captive, ♂ & ♀	E, P, GC	Young, K. M. <i>et al.</i> (2001, 2004)
European badger <i>Meles meles</i>	behavioural and physiological response to restraint, stress	wild, ♂ & ♀	GC	Schütz <i>et al.</i> (2006)
Eurasian otter <i>Lutra lutra</i>	population structure, faecal DNA and steroids	wild, ♂ & ♀	P, A	Kalz <i>et al.</i> (2006)
Sea otter <i>Enhydra lutris</i>	reproductive physiology, oestrous cycle, pregnancy	captive, ♀	E, P	Larson <i>et al.</i> (2003)
OTARIIDAE Steller sea lion <i>Eumetopias jubatus</i>	wildlife rehabilitation, stress	captive, ♀	GC	Petrauskas <i>et al.</i> (2006)
URSIDAE Brown bear <i>Ursus arctos</i> Giant panda <i>Ailuropoda melanoleuca</i> Spectacled bear <i>Tremarctos ornatus</i>	seasonal reproduction, pheromones, comparative study, faecal and urine samples	captive, ♀	E, P, A	Dehnhard <i>et al.</i> (2006)
Malayan sun bear <i>Helarctos malayanus</i>	oestrous cycle, pregnancy, seasonality, aging, immuno-contraception	captive, ♀	E, P, A	Schwarzenberger <i>et al.</i> (2004)

Table 1. Faecal steroid metabolites have been analysed in a variety of mammalian species; owing to space limitations the enclosed table does not include all available literature. Further tables including references on faecal steroid analysis have been published in previous reviews: for example, studies on faecal steroid monitoring in felids have been summarized by Brown (2006) and studies on rhinoceros have been summarized by Roth (2006). Studies on the application of radioactively labelled steroids and on the validation of faecal glucocorticoid metabolites are only partly included in this table, as these studies have been summarized in recent reviews by Palme *et al.* (2005), Touma & Palme (2005) and Keay *et al.* (2006). Reproductive hormones (E, oestrogens; A, androgens; P, progestagens); adrenal hormones (GC, glucocorticoids). DNA, deoxyribonucleic acid.

and adrenal (glucocorticoid) steroid metabolites. Additional references on glucocorticoids not included in this review can be found in the reviews by Möstl & Palme (2002), Palme *et al.* (2005), Touma & Palme (2005) and Keay *et al.* (2006).

#### METHODOLOGICAL ISSUES IN APPLYING FAECAL STEROID ASSAYS

In comparison with the more traditional analysis of steroid hormones in blood, faecal steroid analysis has several advantages. The most obvious is that the technique is non-invasive to the investigated subjects and thus does not introduce variables that may alter the results. Consequently, in many studies in which either blood sampling on a regular basis was difficult or not possible, such as in research on wildlife, very stress-prone species or animals of small size, faecal samples have become a substitute for analysing hormones in the serum or plasma. Faecal steroid analysis has been applied on species of all sizes, ranging from animals as small as mice to elephants or whales (for references, see Table 1 and Touma & Palme, 2005). Faecal steroid analysis can be applied in longitudinal studies and in conjunction with other parameters, such as behaviour or reproduction, can give an accurate insight into the endocrine physiology of a species. Animals in captivity are ideal research subjects because regular sample collection is possible and, thus, techniques can be validated easily. Once established, non-invasive endocrine analysis can be used as a tool to assist in the husbandry of animals in captivity and to investigate social and ecological effects of animals in the wild (Schwarzenberger *et al.*, 1996, 1997; Whitten *et al.*, 1998; Möstl & Palme, 2002; von der Ohe & Servheen, 2002; Monfort, 2003; Graham, 2004; Millsbaugh & Washburn, 2004; Möstl *et al.*, 2005; Touma & Palme, 2005; Ziegler & Wittwer, 2005; Keay *et al.*, 2006; Lane, 2006).

Studies in free-ranging species are often confronted with difficulties in locating sam-

ples, usually involving observation of known individuals and collecting samples upon defecation. However, non-invasive faecal steroid analysis also offers the opportunity to study free-ranging animals for which direct observation of defecation is difficult or impossible. Difficulties with locating samples in some studies have been recently overcome by training dogs for scat detection (Wasser *et al.*, 2004), and this has even been used in open water for detecting samples from free-ranging North Atlantic right whales *Eubalaena glacialis* (Rolland *et al.*, 2005; Hunt *et al.*, 2006). An even more sophisticated approach would be the combination of faecal steroid analysis with molecular genetic techniques for identifying individual animals from faecal deoxyribonucleic acid (DNA). However, the technical difficulties inherent in the analysis of low quantities of DNA and considerable variability in the results generally tend to limit the efficiency of this approach (Broquet *et al.*, 2007). Therefore, the combination of non-invasive faecal DNA and steroid analysis is not yet common practice. Nonetheless, the potential of this auspicious approach has been demonstrated recently; for example, in a study on seasonal faecal glucocorticoid excretion in Red deer *Cervus elaphus* (Huber *et al.*, 2003) and in a study on the population structure of European otters *Lutra lutra* (Kalz *et al.*, 2006).

Like all laboratory-based methods, assay validation is most important for obtaining useful and accurate results. However, the particularity with faecal steroid analysis is that the parent hormones progesterone, testosterone, cortisol or corticosterone are not (or only barely if at all) present in the faeces. Consequently, it is inaccurate to speak of faecal-progesterone or faecal-cortisol analysis, although this designation is common practice in a considerable proportion of the published literature.

Proper faecal steroid assay validation is all related to steroid metabolism (Palme *et al.*, 1996, 2005; Schwarzenberger *et al.*, 1996, 1997; Möstl *et al.*, 2005; Palme, 2005). Steroids are metabolized by the liver before excretion via urine or bile into the

faeces. During the intestinal passage, steroid metabolites can be re-absorbed into the enterohepatic circulation. The intestinal passage causes a lag-time between the circulation of steroids in plasma and their appearance in the faeces; this delay correlates with the time for the intestinal passage of bile to the rectum (Palme *et al.*, 1996, 2005; Schwarzenberger *et al.*, 1996, 1997; Möstl *et al.*, 2005; Palme, 2005). Owing to the intestinal passage, steroid metabolites in the faeces represent pooled endocrine activity over the previous several hours. Thus, faecal steroid analysis presents a more dampened hormone profile over time with less interference from daily rhythm and acute stress (Schwarzenberger *et al.*, 1996, 1997; Whitten *et al.*, 1998; Möstl & Palme, 2002; von der Ohe & Servheen, 2002; Monfort, 2003; Graham, 2004; Millspough & Washburn, 2004; Möstl *et al.*, 2005; Palme, 2005; Palme *et al.*, 2005; Touma & Palme, 2005; Ziegler & Wittwer, 2005; Keay *et al.*, 2006; Lane, 2006).

A particular disadvantage of faecal steroid analysis is the presence of a vast number of different faecal metabolites existing in even closely related species. For the development of techniques for faecal steroid analysis, experiments on the metabolism of radioactively labelled steroids have provided a valuable insight into the metabolism and the excretion of hormone metabolites via faeces and urine. The route of excretion varies considerably among species, and between steroids within the same species. Radiometabolism studies affirmed that oestrogens in the form of oestradiol and/or oestrone are present in faecal samples and as such can easily be determined by using specific assay or a total oestrogen assay. In contrast, testosterone, progesterone and, especially, cortisol/corticosterone are heavily metabolized and the original hormone is barely present in the faeces. Therefore, specific assays (i.e. those typically used for the analysis of hormones in blood samples) are often of limited value (Palme *et al.*, 1996; Schwarzenberger *et al.*, 1996, 1997; Graham *et al.*, 2001; Möstl *et al.*, 2005; Palme, 2005).

For the development of species-specific assays, the determination of the major faecal hormone metabolites via the analysis of infused radioactively labelled steroids (Palme *et al.*, 1996, 2005; Schwarzenberger *et al.*, 1997; Möstl *et al.*, 2005; Palme, 2005) or the identification of faecal steroid metabolites via gas chromatography/mass spectrometry (Lance *et al.*, 2001) would be analytical possibilities. However, these valuable approaches cannot be applied to all wild-life species and in fact, owing to possible health hazards related to the use of radioactively labelled steroids, radiometabolism studies have mainly been carried out with animals in research facilities (Palme, 2005), rather than animals in zoos. Therefore, the best alternative approach is to test several assays with high cross-reactions against a certain group of steroids; that is, group-specific assays (Palme *et al.*, 1996; Schwarzenberger *et al.*, 1996, 1997; Möstl *et al.*, 2005; Palme, 2005). Because of species-specific differences in hormone metabolite excretion, validation and testing of several such assays should be carried out separately for even closely related species (Palme *et al.*, 1996; Schwarzenberger *et al.*, 1996, 1997; Graham *et al.*, 2001; Young, K. M. *et al.*, 2004; Möstl *et al.*, 2005; Palme, 2005; Berger *et al.*, 2006; Heistermann *et al.*, 2006).

An example of how different the reproductive and endocrine physiology of even closely related species can be, is demonstrated by endocrine studies in the family Rhinocerotidae. Through the use of faecal steroid analysis (additional hormones in urine and saliva have been investigated), endocrine profiles of four of the five extant species have been analysed. None of the four rhinoceros species (White rhinoceros *Ceratotherium simum*, Black rhinoceros *Diceros bicornis*, Indian or Greater one-horned rhinoceros *Rhinoceros unicornis* and Sumatran rhinoceros *Dicerorhinus sumatrensis*) exhibit reproductive cycles of similar length. Furthermore, faecal steroid metabolites excreted in these species vary considerably, underlining the necessity to develop endocrine tests for each species

separately (for references, see Table 1, and Roth, 2006).

Assays that have cross-reactivities with a broad range of pregnanediones and hydroxylated pregnanes have been used successfully to quantify progesterone metabolites in the faeces of a wide range of species including a variety of carnivores (Brown, 2006) and artiodactyl species (Schwarzenberger *et al.*, 1996, 1997; Graham *et al.*, 2001). In contrast, the metabolism of glucocorticoids is more complex and a larger number of assays is used for the analysis of faecal glucocorticoid metabolites (Wasser *et al.*, 2000; Möstl & Palme, 2002; Young, K. M. *et al.*, 2004; Möstl *et al.*, 2005; Palme, 2005; Heistermann *et al.*, 2006; Keay *et al.*, 2006). Currently, as researchers worldwide are applying different techniques, results between studies are only comparable in their physiologic outcome, but usually not in absolute metabolite concentrations.

Although it is strongly suggested that antibodies with known high cross-reactivities for steroid metabolites be used, even commercially available antibodies advertised as hormone specific have been used with success for faecal steroid analysis in several studies. The reason is that these antibodies obviously recognize and bind metabolites, although these cross-reactivities are unknown, because testing is typically carried out only for the better-known hormones and in some instances a few metabolites found in blood. Therefore, biological validation, such as day(s) of mating or parturition, or comparison of stressed versus non-stressed animals, in relation to sustained changes in faecal steroid concentration is of utmost importance. However, while it may not be as critical that the exact steroid metabolites are identified, assays developed for use with faecal steroid analysis usually show a stronger differentiation between pregnant and non-pregnant (Schwarzenberger *et al.*, 1996, 1997), or stressed versus non-stressed animals (Heistermann *et al.*, 2006).

In addition to metabolism, and hence the use of an appropriate immunoassay with adequate cross-reactivities, other concerns in

faecal steroid analysis are extraction techniques, storage and stability of faecal metabolites, determination of faecal immuno-reactive steroid metabolites by high-performance liquid chromatography, gut transit time, diurnal and seasonal variations, as well as gender and diet (Schwarzenberger *et al.*, 1996, 1997; Khan *et al.*, 2002; Möstl & Palme, 2002; von der Ohe & Servheen, 2002; Millspaugh & Washburn, 2004; Möstl *et al.*, 2005; Palme, 2005; Palme *et al.*, 2005; Touma & Palme, 2005; Ziegler & Wittwer, 2005; Keay *et al.*, 2006; Lane, 2006). A general recipe from these studies is to attempt to collect samples at the same time of the day and to freeze samples immediately upon collection. If freezing is not possible, storage in ethanol in order to prevent microbial degradation is the method of choice.

Seasonal variation in faecal steroid analysis is of concern, especially when studying free-ranging wildlife living in a highly seasonal habitat. Of particular importance seems to be the question as to whether food availability, which determines the amount of faeces produced and which is lower during severe winter or drought, would result in higher faecal steroid concentrations. However, experiments on the faecal output of progesterone metabolites in ovariectomized domestic cows supplemented with progesterone-releasing intravaginal devices and with different levels of feed intake (Rabiee *et al.*, 2001) did not demonstrate such a relationship. Although feed intake did influence plasma progesterone concentrations, faecal progesterone metabolites were not affected by the level of feed intake or faecal output, but the daily excretion rate of faecal metabolites was associated with the volume of faeces. Comparable results have been found in an Elk (*C. elaphus*, Cook *et al.*, 2002) on a restricted diet. Derived from this experiment, it is likely that seasonal variation in faecal cortisol metabolites, as described in, for example, Red deer *C. elaphus* (Huber *et al.*, 2003), Chacma baboons *Papio hamadryas* (Weingrill *et al.*, 2004) and Chamois *Rupicapra rupicapra* (Hoby *et al.*, 2006), is influenced by an endogenous rhythm.

## APPLICATION OF FAECAL STEROID ANALYSIS FOR STUDYING WILDLIFE SPECIES

Validation in reproductive studies is rather straightforward as steroid metabolites fluctuate with reproductive events; for example, the oestrous cycle, pregnancy, puberty or seasonality. In recent years, a great many studies using faecal steroid analysis have been conducted on ♀ and ♂ reproductive physiology. Wildlife taxa that are most extensively studied are primates, carnivores, rhinoceros and artiodactyla (Table 1). Traditionally, more focus has been placed on the study of reproductive hormones in ♀♀, but studies on the reproductive physiology in the ♂ are increasing. Studies on the reproductive physiology include, for example, characterization of oestrous-cycle length, spontaneous versus induced ovulation, pregnancy, lactational anovulation, age at the onset of puberty and seasonal patterns of reproduction. Faecal steroid analysis has been used as a tool to enhance the management of wildlife species, both in captivity and in the wild. Important issues are the diagnosis of pregnancy, assessing the presence or absence of ovarian cyclicity, control for treatment of infertility and the assessment of various fertility-control techniques; for example, ultrasonographic monitoring of ovarian events and foetal gestational parameters, contraceptive treatments, oestrous synchronization and endocrine responses to ovulation-induction protocols (Table 1).

The validation of techniques in studying stress responses is more difficult and usually involves questions such as is an animal stressed or non-stressed and does this present as acute or chronic stress, positive or negative stress (distress) (Whitten *et al.*, 1998; Möstl & Palme, 2002; von der Ohe & Servheen, 2002; Wielebnowski, 2003; Millspaugh & Washburn, 2004; Touma & Palme, 2005; Keay *et al.*, 2006; Lane, 2006). Consequently, as compared with reproductive studies interpreting results from glucocorticoid studies is more difficult and more focus should be placed on proper validation of

faecal steroid assays. Besides using known stressors (such as transport or confinement) for physiological validation, several studies (especially in wildlife species kept in captivity) included pharmacological validation experiments. Possibilities include the determination of the excretory fate of radioactively labelled steroids and the application of adrenocorticotrophic hormone challenge and dexamethasone suppression tests (Möstl & Palme, 2002; Palme *et al.*, 2005; Touma & Palme, 2005). In conjunction with proper validation, the analysis of glucocorticoid metabolites is a good indicator of adrenal activity and it has been used to quantify certain stress responses. Results were analysed in relation to behaviour, social status, dominance rank, territoriality, reproduction, environmental factors, population density, disturbance, as an indicator of well-being and welfare, pain, the presence of predators, parasites and as a predictor of mortality (Table 1).

In conclusion, faecal steroid analysis of reproductive and adrenocortical steroid hormones has become an established and widely accepted technique for the analysis of captive and free-ranging wildlife species. Because of species-specific differences in steroid metabolism in even closely related species, careful validation of assay methods is necessary in order to generate meaningful and accurate results. In light of this, captive wildlife species are ideal research subjects, as longitudinal sample collection is possible and studies connecting physiology, endocrinology, reproduction and stress with various social and/or environmental factors can be carried out and used to determine how they impact animal health. For the future management of wildlife populations, these techniques will be important research tools and their importance for studying free-ranging animals within their natural habitat will increase further.

## REFERENCES

- ALTMANN, J., LYNCH, J. W., NGUYEN, N., ALBERTS, S. C. & GESQUIERE, L. R. (2004): Life-history correlates of steroid concentrations in wild peripartum baboons. *American Journal of Primatology* **64**: 95–106.

- ANDRABI, S. M. H. & MAXWELL, W. M. C. (2007): A review on reproductive biotechnologies for conservation of endangered mammalian species. *Animal Reproduction Science* **99**: 223–243.
- ASA, C. S., BAUMAN, J. E., HOUSTON, E. W., FISCHER, M. T., READ, B., BROWNFIELD, C. M. & ROSER, J. F. (2001): Patterns of excretion of fecal estradiol and progesterone and urinary chorionic gonadotropin in Grevy's zebras (*Equus grevyi*): ovulatory cycles and pregnancy. *Zoo Biology* **20**: 185–195.
- ASA, C. S., BAUMAN, K., CALLAHAN, P., BAUMAN, J., VOLKMAN, D. H. & JÖCHLE, W. (2006): GnRH-agonist induction of fertile estrus with either natural mating or artificial insemination, followed by birth of pups in gray wolves (*Canis lupus*). *Theriogenology* **66**: 1778–1782.
- ATKINS, N., JONES, S. M. & EDWARDS, A. (2002): Fecal testosterone concentrations may not be useful for monitoring reproductive status in male blue-tongued lizards (*Tiliqua nigrolutea*: Scincidae). *Journal of Herpetology* **36**: 106–109.
- ATSALIS, S., MARGILIS, S. W., BELLEM, A. & WIELEBNOWSKI, N. (2004): Sexual behavior and hormonal estrus cycles in captive aged lowland gorillas (*Gorilla gorilla*). *American Journal of Primatology* **62**: 123–132.
- BALES, K. L., FRENCH, J. A., MCWILLIAMS, J., LAKE, R. A. & DIETZ, J. M. (2006): Effects of social status, age, and season on androgen and cortisol levels in wild male golden lion tamarins (*Leontopithecus rosalia*). *Hormones and Behavior* **49**: 88–95.
- BALTIC, M., JENNI-EIERMANN, S., ARLETTAZ, R. & PALME, R. (2005): A noninvasive technique to evaluate human-generated stress in the black grouse. *Annals of the New York Academy of Sciences* **1046**: 81–95.
- BARDI, M., SHIMIZU, K., BARRETT, G. M., BORGOGNINI-TARLI, S. M. & HUFFMAN, M. A. (2003): Peripartum sex steroid changes and maternal style in rhesus and Japanese macaques. *General and Comparative Endocrinology* **133**: 323–331.
- BARELLI, C., HEISTERMANN, M., BOESCH, C. & REICHARD, U. H. (2007): Sexual swellings in wild white-handed gibbon females (*Hylobates lar*) indicate the probability of ovulation. *Hormones and Behavior* **51**: 221–230.
- BARRETT, G. M., SHIMIZU, K., BARDI, M., ASABA, S. & MORI, A. (2002): Endocrine correlates of rank, reproduction, and female-directed aggression in male Japanese macaques (*Macaca fuscata*). *Hormones and Behavior* **42**: 85–96.
- BEEHNER, J. C., NGUYEN, N., WANGO, E. O., ALBERTS, S. C. & ALTMANN, J. (2006): The endocrinology of pregnancy and fetal loss in wild baboons. *Hormones and Behavior* **49**: 688–699.
- BERGER, E. M., LEUS, K., VERCAMMEN, P. & SCHWARZENBERGER, F. (2006): Faecal steroid metabolites for non-invasive assessment of reproduction in common warthogs (*Phacochoerus africanus*), red river hogs (*Potamochoerus porcus*) and babirusa (*Babyrousa babyrussa*). *Animal Reproduction Science* **91**: 155–171.
- BERGMAN, T. J., BEEHNER, J. C., CHENEY, D. L., SEYFARTH, R. M. & WHITTEN, P. L. (2005): Correlates of stress in free-ranging male chacma baboons, *Papio hamadryas ursinus*. *Animal Behaviour* **70**: 703–713.
- BERKELEY, E. V., KIRKPATRICK, J. F., SCHAFFER, N. E., BRYANT, W. M. & THRELFALL, W. R. (1997): Serum and fecal steroid analysis of ovulation, pregnancy, and parturition in the black rhinoceros (*Diceros bicornis*). *Zoo Biology* **16**: 121–132.
- BLANVILLAIN, C., BERTHIER, J. L., BOMSEL-DEMONTVOY, M. C., SEMPÈRE, A. J., OLBRIGHT, G. & SCHWARZENBERG, F. (1997): Analysis of reproductive data and measurement of fecal progesterone metabolites to monitor the ovarian function in the pudu, *Pudu puda* (Artiodactyla, Cervidae). *Mammalia* **61**: 589–602.
- BOGDAN, D. & MONFORT, S. L. (2001): Faecal oestrogen and progesterone profiles in breeding and non-breeding female North American porcupine (*Erethizon dorsatum*). *Mammalia* **65**: 73–82.
- BOWERS, S., GANDY, S., PAUL, K., WOODS, L., D'ANGELO, D., HORTON, C., TABAKA, C. & WILLARD, S. (2005): The relationship between vaginal electrical impedance and hormone profiles during pregnancy and parturition of a white rhinoceros (*Ceratotherium simum simum*). *Journal of Zoo and Wildlife Medicine* **36**: 451–456.
- BRADSHAW, F. J., STEAD-RICHARDSON, E. J., REEDER, A. J., OATES, J. E. & BRADSHAW, S. D. (2004): Reproductive activity in captive female honey possums, *Tarsipes rostratus*, assessed by faecal steroid analysis. *General and Comparative Endocrinology* **138**: 20–31.
- BROCKMAN, D. K., WHITTEN, P. L., RICHARD, A. F. & BENANDER, B. (2001): Birth season testosterone levels male Verreaux's sifaka, *Propithecus verreauxi*: insights into socio-demographic factors mediating seasonal testicular function. *Behavioral Ecology and Sociobiology* **49**: 117–127.
- BROQUET, T., MÈNARD, N. & PETIT, E. (2007): Noninvasive population genetics: a review of sample source, diet, fragment length and microsatellite motif effects on amplification success and genotyping error rates. *Conservation Genetics* **8**: 249–260.
- BROWN, J. L. (2006): Comparative endocrinology of domestic and nondomestic felids. *Theriogenology* **66**: 25–36.
- BROWN, J. L., WILDT, D. E., GRAHAM, L. H., BYERS, A. P., COLLINS, L., BARRETT, S. & HOWARD, J. (1995): Natural versus chorionic gonadotropin-induced ovarian responses in the clouded leopard (*Neofelis nebulosa*) assessed by fecal steroid analysis. *Biology of Reproduction* **53**: 93–102.
- BROWN, J. L., BELLEM, A. C., FOURAKER, M., WILDT, D. E. & ROTH, T. L. (2001): Comparative analysis of gonadal and adrenal activity in the black and white rhinoceros in North America by noninvasive endocrine monitoring. *Zoo Biology* **20**: 463–486.
- BROWN, J. L., GRAHAM, L. H., WU, J. M., COLLINS, D. & SWANSON, W. F. (2002): Reproductive endocrine responses to photoperiod and exogenous gonadotropins in the Pallas' cat (*Otocolobus manul*). *Zoo Biology* **21**: 347–364.
- BUCHANAN, K. L. & GOLDSMITH, A. R. (2004): Noninvasive endocrine data for behavioural studies: the importance of validation. *Animal Behaviour* **67**: 183–185.



- BUSSO, J. M., PONZIO, M. F., FIOLE DE CUNEO, M. & RUIZ, R. D. (2007): Noninvasive monitoring of ovarian endocrine activity in the chinchilla (*Chinchilla lanigera*). *General and Comparative Endocrinology* **150**: 288–297.
- CABEZAS, S., BLAS, J., MARCHANT, T. A. & MORENO, S. (2007): Physiological stress levels predict survival probabilities in wild rabbits. *Hormones and Behavior* **51**: 313–320.
- CAMPBELL, C. J., SHIDELER, S. E., TODD, H. E. & LASLEY, B. L. (2001): Fecal analysis of ovarian cycles in female black-handed spider monkeys (*Ateles geoffroyi*). *American Journal of Primatology* **54**: 79–89.
- CARLSTEAD, K. & BROWN, J. L. (2005): Relationships between patterns of fecal corticoid excretion and behavior, reproduction, and environmental factors in captive black (*Diceros bicornis*) and white (*Ceratotherium simum*) rhinoceros. *Zoo Biology* **24**: 215–232.
- CARNEGIE, S. D., FEDIGAN, L. M. & ZIEGLER, T. E. (2005): Behavioral indicators of ovarian phase in white-faced capuchins (*Cebus capucinus*). *American Journal of Primatology* **67**: 51–68.
- CAVIGELLI, S. A., DUBOVICK, T., LEVASH, W., JOLLY, A. & PITTS, A. (2003): Female dominance status and fecal corticoids in a cooperative breeder with low reproductive skew: ring-tailed lemur (*Lemur catta*). *Hormones and Behavior* **43**: 166–179.
- CERDA-MOLINA, A. L., HERNÁNDEZ-LÓPEZ, L., ROJAS-MAYA, S., MURCIA-MEJÍA, C. & MONDRAGÓN-CEBALLOS, R. (2006): Male-induced sociosexual behavior by vaginal secretions in *Macaca arctoides*. *International Journal of Primatology* **27**: 791–807.
- CHAPMAN, C. A., WASSERMAN, M. D., GILLESPIE, T. R., SPEIRS, M. L., LAWES, M. J., SAJ, T. L. & ZIEGLE, T. E. (2006): Do food availability, parasitism, and stress have synergistic effects on red colobus populations living in forest fragments? *American Journal of Physical Anthropology* **131**: 525–534.
- CHELINI, M. O. M., SOUZA, N. L., ROCHA, A. M., FELIPPE, E. C. G. & OLIVEIRA, C. A. (2005): Quantification of fecal estradiol and progesterone metabolites in Syrian hamsters (*Mesocricetus auratus*). *Brazilian Journal of Medical and Biological Research* **38**: 1711–1717.
- COCKREM, J. F. (2005): Conservation and behavioral neuroendocrinology. *Hormones and Behavior* **48**: 492–501.
- COOK, R. C., COOK, J. G., GARROTT, R. A., IRWIN, L. L. & MONFORT, S. L. (2002): Effects of diet and body condition on fecal progesterone excretion in Elk. *Journal of Wildlife Diseases* **38**: 558–565.
- CREEL, S., CREEL, N. M., MILLS, M. G. L. & MONFORT, S. L. (1997): Rank and reproduction in cooperatively breeding African wild dogs: behavioral and endocrine correlates. *Behavioral Ecology* **8**: 298–306.
- CRISTOBAL-AZKARATE, J., CHAVIRA, R., BOECK, L., RODRIGUEZ-LUNA, E. & VEAL, J. J. (2006): Testosterone levels of free-ranging resident mantled howler monkey males in relation to the number and density of solitary males: a test of the challenge hypothesis. *Hormones and Behavior* **49**: 261–267.
- DEHNHARD, M., HILDEBRANDT, T. B., KNAUF, T., JEWGENOW, K., KOLTER, L. & GÖRITZ, F. (2006): Comparative endocrine investigations in three bear species based on urinary steroid metabolites and volatiles. *Theriogenology* **66**: 1755–1761.
- DEL CASTILLO, S. M., BASHAW, M. J., PATTON, M. L., RIECHES, R. R. & BERCOVITCH, F. B. (2005): Fecal steroid analysis of female giraffe (*Giraffa camelopardalis*) reproductive condition and the impact of endocrine status on daily time budgets. *General and Comparative Endocrinology* **141**: 271–281.
- DEMATTEO, K. E., PORTON, I. J., KLEIMAN, D. G. & ASA, C. S. (2006): The effect of the male bush dog (*Speothos venaticus*) on the female reproductive cycle. *Journal of Mammalogy* **87**: 723–732.
- DLONIAK, S. M., FRENCH, J. A. & HOLEKAMP, K. E. (2006): Rank-related maternal effects of androgens on behaviour in wild spotted hyaenas. *Nature* **440**: 1190–1193.
- DUMONCEAUX, G. A., BAUMAN, J. E. & CAMILO, G. R. (2006): Evaluation of progesterone levels in feces of captive reticulated giraffe (*Giraffa camelopardalis reticulata*). *Journal of Zoo and Wildlife Medicine* **37**: 255–261.
- ELLIS, T., JAMES, J. D., STEWART, C. & SCOTT, A. P. (2004): A non-invasive stress assay based upon measurement of free cortisol released into the water by rainbow trout. *Journal of Fish Biology* **65**: 1233–1252.
- EMERY, M. A. & WHITTEN, P. L. (2003): Size of sexual swellings reflects ovarian function in chimpanzees (*Pan troglodytes*). *Behavioral Ecology and Sociobiology* **54**: 340–351.
- ENGELHARDT, A., HODGES, J. K. & HEISTERMANN, M. (2007): Post-conception mating in wild long-tailed macaques (*Macaca fascicularis*): characterization, endocrine correlates and functional significance. *Hormones and Behavior* **51**: 3–10.
- FIESS, M., HEISTERMANN, M. & HODGES, J. K. (1999): Patterns of urinary and fecal steroid excretion during the ovarian cycle and pregnancy in the African elephant (*Loxodonta africana*). *General and Comparative Endocrinology* **115**: 76–89.
- FUJITA, S., MITSUNAGA, F., SUGIURA, H. & SHIMIZU, K. (2001): Measurement of urinary and fecal steroid metabolites during the ovarian cycle in captive and wild Japanese macaques, *Macaca fuscata*. *American Journal of Primatology* **53**: 167–176.
- GANSWINDT, A., HEISTERMANN, M. & HODGES, J. K. (2005): Physical, physiological, and behavioral correlates of musth in captive African elephants (*Loxodonta africana*). *Physiological and Biochemical Zoology* **78**: 505–514.
- GANSWINDT, A., RASMUSSEN, H. B., HEISTERMANN, M. & HODGES, J. K. (2005): The sexually active states of free-ranging male African elephants (*Loxodonta africana*): defining musth and non-musth using endocrinology, physical signals, and behavior. *Hormones and Behavior* **47**: 83–91.
- GARNIER, J. N., HOLT, W. V. & WATSON, P. F. (2002): Non-invasive assessment of oestrous cycles and evaluation of reproductive seasonality in the female wild black rhinoceros (*Diceros bicornis minor*). *Reproduction* **123**: 877–889.

- GESQUIERE, L. R., ALTMANN, J., KHAN, M. Z., COURET, J., YU, J. C., ENDRES, C. S., LYNCH, J. W., OGOLA, P., FOX, E. A., ALBERTS, S. C. & WANGO, E. O. (2005): Coming of age: steroid hormones of wild immature baboons (*Papio cynocephalus*). *American Journal of Primatology* **67**: 83–100.
- GESQUIERE, L. R., WANGO, E. O., ALBERTS, S. C. & ALTMANN, J. (2007): Mechanisms of sexual selection: sexual swellings and estrogen concentrations as fertility indicators and cues for male consort decisions in wild baboons. *Hormones and Behavior* **51**: 114–125.
- GOLDSTEIN, E. J., MILLSAUGH, J. J., WASHBURN, B. E., BRUNDIGE, G. C. & RAEDEKE, K. J. (2005): Relationships among fecal lungworm loads, fecal glucocorticoid metabolites, and lamb recruitment in free-ranging Rocky Mountain bighorn sheep. *Journal of Wildlife Diseases* **41**: 416–425.
- GÖRITZ, F., NEUBAUER, K., NAIDENKO, S. V., FICKEL, J. & JEWGENOW, K. (2006): Investigations on reproductive physiology in the male Eurasian lynx (*Lynx lynx*). *Theriogenology* **66**: 1751–1754.
- GOULD, L., ZIEGLER, T. E. & WITTEWIT, D. J. (2005): Effects of reproductive and social variables on fecal glucocorticoid levels in a sample of adult male ring-tailed lemurs (*Lemur catta*) at the Beza Mahafaly Reserve, Madagascar. *American Journal of Primatology* **67**: 5–23.
- GOYMANN, W. (2005): Noninvasive monitoring of hormones in bird droppings: physiological validation, sampling, extraction, sex differences, and the influence of diet on hormone metabolite levels. *Annals of the New York Academy of Sciences* **1046**: 35–53.
- GOYMANN, W., EAST, M. L., WACHTER, B., HÖNER, O. P., MÖSTL, E., VAN'T HOF, T. J. & HOFER, H. (2001): Social, state-dependent and environmental modulation of faecal corticosteroid levels in free-ranging female spotted hyenas. *Proceedings of the Royal Society – Biological Sciences B* **268**: 2453–2459.
- GOYMANN, W., EAST, M. L., WACHTER, B., HÖNER, O. P., MÖSTL, E. & HOFER, H. (2003): Social status does not predict corticosteroid levels in postdispersal male spotted hyenas. *Hormones and Behavior* **43**: 474–479.
- GRAHAM, L. H. (2004): Non-invasive monitoring of reproduction in zoo and wildlife species. *Annual Review of Biomedical Sciences* **6**: 91–98.
- GRAHAM, L., SCHWARZENBERGER, F., MÖSTL, E., GALAMA, W. & SAVAGE, A. (2001): A versatile enzyme immunoassay for the determination of progesterone in feces and serum. *Zoo Biology* **20**: 227–236.
- GRAHAM, L. H., REID, K., WEBSTER, T., RICHARDS, M. & JOSEPH, S. (2002): Endocrine patterns associated with reproduction in the Nile hippopotamus (*Hippopotamus amphibius*) as assessed by fecal progesterone analysis. *General and Comparative Endocrinology* **128**: 74–81.
- GRAHAM, L. H., BYERS, A. P., ARMSTRONG, D. L., LOSKUTOFF, N. M., SWANSON, W. F., WILDT, D. E. & BROWN, J. L. (2006): Natural and gonadotropin-induced ovarian activity in tigers (*Panthera tigris*) assessed by fecal steroid analyses. *General and Comparative Endocrinology* **147**: 362–370.
- HATTAB, S. A., KADDOOM, A. K., PALME, R. & BAMBERG, E. (2000): Effect of CRESTAR(TM) on estrus synchronization and the relationship between fecal and plasma concentrations of progesterone in buffalo cows. *Theriogenology* **54**: 1007–1017.
- HEISTERMANN, M., ZIEGLER, T., VAN SCHAİK, C. P., LAUNHARDT, K., WINKLER, P. & HODGES, J. K. (2001): Loss of oestrus, concealed ovulation and paternity confusion in free-ranging Hanuman langurs. *Proceedings of the Royal Society – Biological Sciences B* **268**: 2445–2451.
- HEISTERMANN, M., ADEMMEER, C. & KAUMANN, W. (2004): Ovarian cycle and effect of social changes on adrenal and ovarian function in *Pygathrix nemaeus*. *International Journal of Primatology* **25**: 689–708.
- HEISTERMANN, M., PALME, R. & GANSWINDT, A. (2006): Comparison of different enzyme immunoassays for assessment of adrenocortical activity in primates based on fecal analysis. *American Journal of Primatology* **68**: 257–273.
- HERMES, R., HILDEBRANDT, T. B., WALZER, C., GÖRITZ, F., PATTON, M. L., SILINSKI, S., ANDERSON, M. J., REID, C. E., WIBBELT, G., TOMASOVA, K. & SCHWARZENBERGER, F. (2006): The effect of long non-reproductive periods on the genital health in captive female white rhinoceroses (*Ceratotherium simum simum*, *C. s. cottoni*). *Theriogenology* **65**: 1492–1515.
- HERNÁNDEZ-LÓPEZ, L., CERDA-MOLINA, A. L., PÁEZ-PONCE, D. L., ROJAS-MAYA, S. & MONDRAGÓN-CEBALLOS, R. (2007): Artificial insemination in black-handed spider monkey (*Ateles geoffroyi*). *Theriogenology* **67**: 399–406.
- HOBYS, S., SCHWARZENBERGER, F., DOHERR, M. G., ROBERT, N. & WALZER, C. (2006): Steroid hormone related male biased parasitism in chamois, *Rupicapra rupicapra rupicapra*. *Veterinary Parasitology* **138**: 337–348.
- HUBER, S., PALME, R. & ARNOLD, W. (2003): Effects of season, sex, and sample collection on concentrations of fecal cortisol metabolites in red deer (*Cervus elaphus*). *General and Comparative Endocrinology* **130**: 48–54.
- HUNT, K. E., ROLLAND, R. M., KRAUS, S. D. & WASSER, S. K. (2006): Analysis of fecal glucocorticoids in the North Atlantic right whale (*Eubalaena glacialis*). *General and Comparative Endocrinology* **148**: 260–272.
- JOHNSTON, S. D., NICOLSON, V., MADDEN, C., LOGIE, S., PYNE, M., ROSER, A., LISLE, A. T. & D'OCCHIO, M. (2007): Assessment of reproductive status in male echidnas. *Animal Reproduction Science* **97**: 114–127.
- KALLERT, G., GOTTSCHALK, J., NEUMANN, G., EULENBERGER, K. & GRÜN, E. (2002): Fortpflanzungsbiologie der Schneegäümse (*Oreamnos americanus*). Ein Beitrag unter besonderer Berücksichtigung der Gestagenkonzentration der Fäzes im Verlauf der Trächtigkeit. *Tierärztliche Praxis Ausgabe G: Grosstiere – Nutztiere* **30**: 240–243.
- KALZ, B., JEWGENOW, K. & FICKEL, J. (2006): Structure of an otter (*Lutra lutra*) population in Germany: results of DNA and hormone analyses from faecal samples. *Mammalian Biology* **71**: 321–335.
- KAPKE, C. A., ARCESE, P., ZIEGLER, T. E. & SCHEFFLER, G. R. (1999): Estradiol and progesterone metabolite concentration in white-tailed deer (*Odocoileus virginianus*)

- feces. *Journal of Zoo and Wildlife Medicine* **30**: 361–371.
- KEAY, J. M., SINGH, J., GAUNT, M. C. & KAUR, T. (2006): Fecal glucocorticoids and their metabolites as indicators of stress in various mammalian species: a literature review. *Journal of Zoo and Wildlife Medicine* **37**: 234–244.
- KHAN, M. Z., ALTMANN, J., ISANI, S. S. & YU, J. (2002): A matter of time: evaluating the storage of fecal samples for steroid analysis. *General and Comparative Endocrinology* **128**: 57–64.
- KRETZSCHMAR, P., GANSLOEBER, U. & DEHNHARD, M. (2004): Relationship between androgens, environmental factors and reproductive behavior in male white rhinoceros (*Ceratotherium simum simum*). *Hormones and Behavior* **45**: 1–9.
- KUZNETSOV, V. A., TCHABOVSKY, A. V., KOLOSOVA, I. E. & MOSHKIN, M. P. (2004): Effect of habitat type and population density on the stress level of midday gerbils (*Meriones meridianus* Pall.) in free-living populations. *Biology Bulletin* **31**: 628–632.
- LANCE, V. A., PATTON, M. L. & HAGEY, L. R. (2001): Identification of a series of C<sub>21</sub>O<sub>2</sub> pregnanes from fecal extracts of a pregnant black rhinoceros (*Diceros bicornis minor*). *Steroids* **66**: 875–881.
- LANE, J. (2006): Can non-invasive glucocorticoid measures be used as reliable indicators of stress in animals? *Animal Welfare* **15**: 331–342.
- LANYON, J. M., SMITH, K. M. & CARRICK, F. N. (2005): Reproductive steroids are detectable in the faeces of dugongs. *Australian Zoologist* **33**: 247–250.
- LARSON, S., CASSON, C. J. & WASSER, S. (2003): Noninvasive reproductive steroid hormone estimates from fecal samples of captive female sea otters (*Enhydra lutris*). *General and Comparative Endocrinology* **134**: 18–25.
- LI, C., JIANG, Z., JIANG, G. & FANG, J. (2001): Seasonal changes of reproductive behavior and fecal steroid concentrations in Père David's deer. *Hormones and Behavior* **40**: 518–525.
- LYNCH, J. W., ZIEGLER, T. E. & STRIER, K. B. (2002): Individual and seasonal variation in fecal testosterone and cortisol levels of wild male tufted capuchin monkeys, *Cebus apella nigrilus*. *Hormones and Behavior* **41**: 275–287.
- MACDONALD, E. A., NORTHROP, L. E. & CZEKALA, N. M. (2005): Pregnancy detection from fecal progesterin concentrations in the red panda (*Ailures fulgens fulgens*). *Zoo Biology* **24**: 419–429.
- MATSUURA, Y., SASAMOTO, Y., SATO, K., TAKAHASHI, Y., SUZUKI, M. & OHTAISHI, N. (2004): Monitoring ovarian cycle and conception by fecal progesterone analysis in sika deer. *Ecological Research* **19**: 397–404.
- MAUGET, R., MAUGET, C., DUBOST, G., CHARRON, F., COURCOUL, A. & RODIER, A. (2007): Non-invasive assessment of reproductive status in Chinese water deer (*Hydropotes inermis*): correlation with sexual behaviour. *Mammalian Biology* **72**: 14–26.
- MCKENZIE, S. & DEANE, E. M. (2005): Faecal corticosteroid levels as an indicator of well-being in the tammar wallaby, *Macropus eugenii*. *Comparative Biochemistry and Physiology – A Molecular and Integrative Physiology* **140**: 81–87.
- MILLSPAUGH, J. J. & WASHBURN, B. E. (2004): Use of fecal glucocorticoid metabolite measures in conservation biology research: considerations for application and interpretation. *General and Comparative Endocrinology* **138**: 189–199.
- MONCLUS, R., RÖDEL, H. G., PALME, R., VON HOLST, D. & DEMIGUEL, J. (2006): Non-invasive measurement of the physiological stress response of wild rabbits to the odour of a predator. *Chemoecology* **16**: 25–29.
- MONFORT, S. L. (2003): Non-invasive endocrine measures of reproduction and stress in wild populations. In *Reproductive science and integrated conservation*: 147–165.
- Holt, W. V., Pickard, A. R., Rodger, J. & Wildt, D. E. (Eds). Cambridge: Cambridge University Press.
- MOORING, M. S., PATTON, M. L., LANCE, V. A., HALL, B. M., SCHAAD, E. W., FORTIN, S. S., JELLA, J. E. & McPEAK, K. M. (2004): Fecal androgens of bison bulls during the rut. *Hormones and Behavior* **46**: 392–398.
- MOORING, M. S., PATTON, M. L., LANCE, V. A., HALL, B. M., SCHAAD, E. W., FETTER, G. A., FORTIN, S. S. & McPEAK, K. M. (2006): Glucocorticoids of bison bulls in relation to social status. *Hormones and Behavior* **49**: 369–375.
- MORATO, R. G., BUENO, M. G., MALMHEISTER, P., VERRESCHI, I. T. N. & BARNABE, R. C. (2004): Changes in the fecal concentrations of cortisol and androgen metabolites in captive male jaguars (*Panthera onca*) in response to stress. *Brazilian Journal of Medical and Biological Research* **37**: 1903–1907.
- MORROW, C. J., WILDT, D. E. & MONFORT, S. L. (1999): Reproductive seasonality in the female scimitar-horned oryx (*Oryx dammah*). *Animal Conservation* **2**: 261–268.
- MORROW, C. J., WOLFE, B. A., ROTH, T. L., WILDT, D. E., BUSH, M., BLUMER, E. S., ATKINSON, M. W. & MONFORT, S. L. (2000): Comparing ovulation synchronization protocols for artificial insemination in the scimitar-horned oryx (*Oryx dammah*). *Animal Reproduction Science* **59**: 71–86.
- MOSS, A. M., CLUTTON-BROCK, T. H. & MONFORT, S. L. (2001): Longitudinal gonadal steroid excretion in free-living male and female meerkats (*Suricata suricatta*). *General and Comparative Endocrinology* **122**: 158–171.
- MOSTL, E. & PALME, R. (2002): Hormones as indicators of stress. *Domestic Animal Endocrinology* **23**: 67–74.
- MOSTL, E., RETTENBACHER, S. & PALME, R. (2005): Measurement of corticosterone metabolites in birds' droppings: an analytical approach. *Annals of the New York Academy of Sciences* **1046**: 17–34.
- MUEHLENBEIN, M. P. (2006): Intestinal parasite infections and fecal steroid levels in wild chimpanzees. *American Journal of Physical Anthropology* **130**: 546–550.
- MUEHLENBEIN, M. P., WATTS, D. P. & WHITTEN, P. L. (2004): Dominance rank and fecal testosterone levels in adult male chimpanzees (*Pan troglodytes schweinfurthii*) at Ngogo, Kibale National Park, Uganda. *American Journal of Primatology* **64**: 71–82.
- MÜHLBAUER, M., DUARTE, D. P. F., GILMORE, D. P. & DACOSTA, C. P. (2006): Fecal estradiol and progesterone metabolite levels in the three-toed sloth (*Bradypus*

- variegatus*). *Brazilian Journal of Medical and Biological Research* **39**: 289–295.
- OATES, J. E., BRADSHAW, F. J., BRADSHAW, S. D. & LONSDALE, R. A. (2002): Sex identification and evidence of gonadal activity in the short-beaked echidna (*Tachyglossus aculeatus*) (Monotremata: Tachyglossidae): non-invasive analysis of faecal sex steroids. *Australian Journal of Zoology* **50**: 395–406.
- OATES, J. E., BRADSHAW, F. J. & BRADSHAW, S. D. (2004): The influence of photoperiod on the reproductive activity of female honey possums, *Tarsipes rostratus* (Marsupialia: *Tarsipedidae*): assessed by faecal progesteragens and oestradiol-17 $\beta$ . *General and Comparative Endocrinology* **139**: 103–112.
- OATES, J. E., BRADSHAW, F. J., BRADSHAW, S. D., STEAD-RICHARDSON, E. J. & PHILIPPE, D. L. (2007): Reproduction and embryonic diapause in a marsupial: insights from captive female honey possums, *Tarsipes rostratus* (*Tarsipedidae*). *General and Comparative Endocrinology* **150**: 445–461.
- VON DER OHE, C. G. & SERVHEEN, C. (2002): Measuring stress in mammals using fecal glucocorticoids: opportunities and challenges. *Wildlife Society Bulletin* **30**: 1215–1225.
- O'NEILL, A. C., FEDIGAN, L. M. & ZIEGLER, T. E. (2004): Relationship between ovarian cycle phase and sexual behavior in female Japanese macaques (*Macaca fuscata*). *American Journal of Physical Anthropology* **125**: 352–362.
- OSTNER, J. & HEISTERMANN, M. (2003): Endocrine characterization of female reproductive status in wild red-fronted lemurs (*Eulemur fulvus rufus*). *General and Comparative Endocrinology* **131**: 274–283.
- OSTNER, J., KAPPELER, P. M. & HEISTERMANN, M. (2002): Seasonal variation and social correlates of androgen excretion in male redfronted lemurs (*Eulemur fulvus rufus*). *Behavioral Ecology and Sociobiology* **52**: 485–495.
- OSTROWSKI, S., BLANVILLAIN, C., MESOCHINA, P., ISMAIL, K. & SCHWARZENBERGER, F. (2005): Monitoring reproductive steroids in feces of Arabian oryx: toward a non-invasive method to predict reproductive status in the wild. *Wildlife Society Bulletin* **33**: 965–973.
- PALME, R. (2005): Measuring fecal steroids: guidelines for practical application. *Annals of the New York Academy of Sciences* **1046**: 75–80.
- PALME, R., FISCHER, P., SCHILDORFER, H. & ISMAIL, M. N. (1996): Excretion of infused 14C-steroid hormones via faeces and urine in domestic livestock. *Animal Reproduction Science* **43**: 43–63.
- PALME, R., RETTENBACHER, S., TOUMA, C., EL-BAHR, S. M. & MÖSTL, E. (2005): Stress hormones in mammals and birds: comparative aspects regarding metabolism, excretion, and noninvasive measurement in fecal samples. *Annals of the New York Academy of Sciences* **1040**: 162–171.
- PARIS, M. C. J., WHITE, A., REISS, A., WEST, M. & SCHWARZENBERGER, F. (2002): Faecal progesterone metabolites and behavioural observations for the non-invasive assessment of oestrous cycles in the common wombat (*Vombatus ursinus*) and the southern hairy-nosed wombat (*Lasiorhinus latifrons*). *Animal Reproduction Science* **72**: 245–257.
- PATTON, M. L., SWAISGOOD, R. R., CZEKALA, N. M., WHITE, A. M., FETTER, G. A., MONTAGNE, J. P., RIECHES, R. G. & LANCE, V. A. (1999): Reproductive cycle length and pregnancy in the southern white rhinoceros (*Ceratotherium simum simum*) as determined by fecal pregnane analysis and observations of mating behavior. *Zoo Biology* **18**: 111–127.
- PATTON, M. L., WHITE, A. M., SWAISGOOD, R. R., SPROUL, R. L., FETTER, G. A., KENNEDY, J., EDWARDS, M. S., RIECHES, R. G. & LANCE, V. A. (2001): Aggression control in a bachelor herd of fringe-eared oryx (*Oryx gazella callotis*), with melengestrol acetate: behavioral and endocrine observations. *Zoo Biology* **20**: 375–388.
- PATTON, M. L., BASHAW, M. J., DEL CASTILLO, S. M., JÖCHLE, W., LAMBERSKI, N., RIECHES, R. & BERCOVITZ, F. B. (2006): Long-term suppression of fertility in female giraffe using the GnRH agonist deslorelin as a long-acting implant. *Theriogenology* **66**: 431–438.
- PATZL, M., SCHWARZENBERGER, F., OSMANN, C., BAMBERG, E. & BARTMANN, W. (1998): Monitoring ovarian cycle and pregnancy in the giant anteater (*Myrmecophaga tridactyla*) by faecal progesteragen and oestrogen analysis. *Animal Reproduction Science* **53**: 209–219.
- PELICAN, K. M., WILDT, D. E. & HOWARD, J. G. (2006): GnRH agonist Lupron<sup>®</sup> (leuprolide acetate) pre-treatments prevent ovulation in response to gonadotropin stimulation in the clouded leopard (*Neofelis nebulosa*). *Theriogenology* **66**: 1768–1777.
- PELLETIER, F., BAUMAN, J. & FESTA-BIANCHET, M. (2003): Fecal testosterone in bighorn sheep (*Ovis canadensis*): behavioural and endocrine correlates. *Canadian Journal of Zoology* **81**: 1678–1684.
- PENFOLD, L. M., MONFORT, S. L., WOLFE, B. A., CITINO, S. B. & WILDT, D. E. (2005): Reproductive physiology and artificial insemination studies in wild and captive gerenuk (*Litocranius walleri walleri*). *Reproduction, Fertility and Development* **17**: 707–714.
- PEREIRA, R. J. G., DUARTE, J. M. B. & NEGRAO, J. A. (2005): Seasonal changes in fecal testosterone concentrations and their relationship to the reproductive behavior, antler cycle and grouping patterns in free-ranging male pampas deer (*Ozotoceros bezoarticus bezoarticus*). *Theriogenology* **63**: 2113–2125.
- PEREIRA, R. J. G., BARBANTI DUARTE, J. M. & NEGRAO, J. A. (2006): Effects of environmental conditions, human activity, reproduction, antler cycle and grouping on fecal glucocorticoids of free-ranging Pampas deer stags (*Ozotoceros bezoarticus bezoarticus*). *Hormones and Behavior* **49**: 114–122.
- PEREIRA, R. J. G., POLEGATO, B. F., DE SOUZA, S., NEGRAO, J. A. & DUARTE, J. M. B. (2006): Monitoring ovarian cycles and pregnancy in brown brocket deer (*Mazama gouazoubira*) by measurement of fecal progesterone metabolites. *Theriogenology* **65**: 387–399.
- PETRAUSKAS, L., TUOMI, P. & ATKINSON, S. (2006): Non-invasive monitoring of stress hormone levels in a female Steller sea lion (*Eumetopias jubatus*) pup undergoing rehabilitation. *Journal of Zoo and Wildlife Medicine* **37**: 75–78.

- PETTITT, B. A., WHEATON, C. J. & WATERMAN, J. M. (2007): Effects of storage treatment on fecal steroid hormone concentrations of a rodent, the cape ground squirrel (*Xerus inauris*). *General and Comparative Endocrinology* **150**: 1–11.
- PICKARD, A. R., ABAIGAR, T., GREEN, D. I., HOLT, W. V. & CANO, M. (2001): Hormonal characterization of the reproductive cycle and pregnancy in the female Mohor gazelle (*Gazella dama mhorr*). *Reproduction* **122**: 571–580.
- PRIDE, R. E. (2005): High faecal glucocorticoid levels predict mortality in ring-tailed lemurs (*Lemur catta*). *Biology Letters* **1**: 60–63.
- PUKAZHENTHI, B. S. & WILDT, D. E. (2004): Which reproductive technologies are most relevant to studying, managing and conserving wildlife? *Reproduction, Fertility and Development* **16**: 33–46.
- QUEYRAS, A. & CAROSI, M. (2004): Non-invasive techniques for analysing hormonal indicators of stress. *Annali dell'Istituto Superiore di Sanita* **40**: 211–221.
- RABIEE, A. R., MACMILLAN, K. L. & SCHWARZENBERGER, F. (2001): The effect of level of feed intake on progesterone clearance rate by measuring faecal progesterone metabolites in grazing dairy cows. *Animal Reproduction Science* **67**: 205–214.
- RACHLOW, J. L., BERKELEY, E. V. & BERGER, J. (1998): Correlates of male mating strategies in white rhinos (*Ceratotherium simum*). *Journal of Mammalogy* **79**: 1317–1324.
- RADCLIFFE, R. W., CZEKALA, N. M. & OSOFSKY, S. A. (1997): Combined serial ultrasonography and fecal progesterin analysis for reproductive evaluation of the female white rhinoceros (*Ceratotherium simum simum*): preliminary results. *Zoo Biology* **16**: 445–456.
- RADCLIFFE, R. W., EYRES, A. I., PATTON, M. L., CZEKALA, N. M. & EMSLIE, R. H. (2001): Ultrasonographic characterization of ovarian events and fetal gestational parameters in two southern black rhinoceros (*Diceros bicornis minor*) and correlation to fecal progesterone. *Theriogenology* **55**: 1033–1049.
- REHBINDER, C. & HAU, J. (2006): Quantification of cortisol, cortisol immunoreactive metabolites, and immunoglobulin A in serum, saliva, urine, and feces for noninvasive assessment of stress in reindeer. *Canadian Journal of Veterinary Research* **70**: 151–154.
- REIMERS, M., SCHWARZENBERGER, F. & PREUSCHOF, S. (2007): Rehabilitation of research chimpanzees: stress and coping after long-term isolation. *Hormones and Behavior* **51**: 428–435.
- RITTENHOUSE, C. D., MILLSPAUGH, J. J., WASHBURN, B. E. & HUBBARD, M. W. (2005): Effects of radiotransmitters on fecal glucocorticoid metabolite levels of three-toed box turtles in captivity. *Wildlife Society Bulletin* **33**: 706–713.
- ROLLAND, R. M., HUNT, K. E., KRAUS, S. D. & WASSER, S. K. (2005): Assessing reproductive status of right whales (*Eubalaena glacialis*) using fecal hormone metabolites. *General and Comparative Endocrinology* **142**: 308–317.
- ROTH, T. L. (2006): A review of the reproductive physiology of rhinoceros species in captivity. *International Zoo Yearbook* **40**: 130–143.
- ROTH, T. L., O'BRIEN, J. K., McRAE, M. A., BELLEM, A. C., ROMO, S. J., KROLL, J. L. & BROWN, J. L. (2001): Ultrasound and endocrine evaluation of the ovarian cycle and early pregnancy in the Sumatran rhinoceros, *Dicerorhinus sumatrensis*. *Reproduction* **121**: 139–149.
- ROTH, T. L., BATEMAN, H. L., KROLL, J. L., STEINETZ, B. G. & REINHART, P. R. (2004): Endocrine and ultrasonographic characterization of a successful pregnancy in a Sumatran rhinoceros (*Dicerorhinus sumatrensis*) supplemented with a synthetic progestin. *Zoo Biology* **23**: 219–238.
- SANDS, J. & CREEL, S. (2004): Social dominance, aggression and faecal glucocorticoid levels in a wild population of wolves, *Canis lupus*. *Animal Behaviour* **67**: 387–396.
- SCHEIBE, K. M., DEHNHARD, M., MEYER, H. H. D. & SCHEIBE, A. (1999): Noninvasive monitoring of reproductive function by determination of faecal progestagens and sexual behaviour in a herd of Przewalski mares in a semireserve. *Acta Theriologica* **44**: 451–463.
- SCHOENECKER, K. A., LYDA, R. O. & KIRKPATRICK, J. (2004): Comparison of three fecal steroid metabolites for pregnancy detection used with single sampling in bighorn sheep (*Ovis canadensis*). *Journal of Wildlife Diseases* **40**: 273–281.
- SCHÜTZ, K. E., AGREN, E., AMUNDIN, M., RÖKEN, B., PALME, R. & MÖRNER, T. (2006): Behavioral and physiological response of trap-induced stress in European badgers. *Journal of Wildlife Management* **70**: 884–891.
- SCHWARZENBERGER, F., MÖSTL, E., PALME, R. & BAMBERG, E. (1996): Faecal steroid analysis for non-invasive monitoring of reproductive status in farm, wild and zoo animals. *Animal Reproduction Science* **42**: 515–526.
- SCHWARZENBERGER, F., PALME, R., BAMBERG, E. & MÖSTL, E. (1997): A review of faecal progesterone metabolite analysis for non-invasive monitoring of reproductive function in mammals. *Journal of Mammalian Biology* **62**(suppl. 2): 214–221.
- SCHWARZENBERGER, F., WALZER, C., TOMASOVA, K., VAHALA, J., MEISTER, J., GOODROWE, K. L., ZIMA, J., STRAUB, G. & LYNCH, M. (1998): Faecal progesterone metabolite analysis for non-invasive monitoring of reproductive function in the white rhinoceros (*Ceratotherium simum*). *Animal Reproduction Science* **53**: 173–190.
- SCHWARZENBERGER, F., RIETSCHEL, W., MATERN, B., SCHAFTENAAR, W., BIRCHER, P., VAN PUJENBROECK, B. & LEUS, K. (1999): Noninvasive reproductive monitoring in the okapi (*Okapia johnstoni*). *Journal of Zoo and Wildlife Medicine* **30**: 497–503.
- SCHWARZENBERGER, F., RIETSCHEL, W., VAHALA, J., HOLECKOVA, D., THOMAS, P., MALTZAN, J., BAUMGARTNER, K. & SCHAFTENAAR, W. (2000): Fecal progesterone, estrogen, and androgen metabolites for noninvasive monitoring of reproductive function in the female Indian rhinoceros, *Rhinoceros unicornis*. *General and Comparative Endocrinology* **119**: 300–307.
- SCHWARZENBERGER, F., FREDRIKSSON, G., SCHALLER, K. & KOLTER, L. (2004): Fecal steroid analysis for monitoring reproduction in the sun bear (*Helarctos malayanus*). *Theriogenology* **62**: 1677–1692.
- SHIMIZU, K. (2005): Studies on reproductive endocrinology in non-human primates: application of non-invasive

- methods. *Journal of Reproduction and Development* **51**: 1–13.
- SPANNER, A., STONE, G. M. & SCHULTZ, D. (1997): Excretion profiles of some reproductive steroids in the faeces of captive Nepalese red panda (*Ailurus fulgens fulgens*). *Reproduction, Fertility and Development* **9**: 565–570.
- STEAD-RICHARDSON, E. J., BRADSHAW, S. D., BRADSHAW, F. J. & GAIKHORST, G. (2001): Monitoring the oestrous cycle of the chuditch (*Dasyurus geoffroii*) (Marsupialia: Dasyuridae): non-invasive analysis of faecal oestradiol-17 $\beta$ . *Australian Journal of Zoology* **49**: 183–193.
- STOOPS, M. A., ANDERSON, G. B., LASLEY, B. L. & SHIDELER, S. E. (1999): Use of fecal steroid metabolites to estimate the pregnancy rate of a free-ranging herd of tule Elk. *Journal of Wildlife Management* **63**: 561–569.
- STRIER, K. B., LYNCH, J. W. & ZIEGLER, T. E. (2003): Hormonal changes during the mating and conception seasons of wild northern muriquis (*Brachyteles arachnoides hypoxanthus*). *American Journal of Primatology* **61**: 85–99.
- SZOKÉ, Z., VARADI, E., KELEMEN, K., BICZO, A. & PECZELY, P. (2006): Effect of testosterone loading on the kinetic of faecal testosterone excretion in mallards. *Acta Biologica Hungarica* **57**: 13–21.
- SZYMANSKI, D. C., GIST, D. H. & ROTH, T. L. (2006): Anuran gender identification by fecal steroid analysis. *Zoo Biology* **25**: 35–46.
- TERIO, K. A., MARKER, L. & MUNSON, L. (2004): Evidence for chronic stress in captive but not free-ranging cheetahs (*Acinonyx jubatus*) based on adrenal morphology and function. *Journal of Wildlife Diseases* **40**: 259–266.
- THOMPSON, K. V. & MONFORT, S. L. (1999): Synchronization of oestrous cycles in sable antelope. *Animal Reproduction Science* **57**: 185–197.
- THOMPSON, K. V., MASHBURN, K. L. & MONFORT, S. L. (1998): Characterization of estrous cyclicity in the sable antelope (*Hippotragus niger*) through fecal progesterone monitoring. *General and Comparative Endocrinology* **112**: 129–137.
- TOUMA, C. & PALME, R. (2005): Measuring fecal glucocorticoid metabolites in mammals and birds: the importance of validation. *Annals of the New York Academy of Sciences* **1046**: 54–74.
- TURNER JR, J. W., TOLSON, P. & HAMAD, N. (2002): Remote assessment of stress in white rhinoceros (*Ceratotherium simum*) and black rhinoceros (*Diceros bicornis*) by measurement of adrenal steroids in feces. *Journal of Zoo and Wildlife Medicine* **33**: 214–221.
- TURNER JR, J. W., NEMETH, R. & ROGERS, C. (2003): Measurement of fecal glucocorticoids in parrotfishes to assess stress. *General and Comparative Endocrinology* **133**: 341–352.
- VALDESPINO, C., ASA, C. S. & BAUMAN, J. E. (2002): Estrous cycles, copulation, and pregnancy in the fennec fox (*Vulpes zerda*). *Journal of Mammalogy* **83**: 99–109.
- VELLOSO, A. L., WASSER, S. K., MONFORT, S. L. & DIETZ, J. M. (1998): Longitudinal fecal steroid excretion in maned wolves (*Chrysocyon brachyurus*). *General and Comparative Endocrinology* **112**: 96–107.
- VERVAECKE, H. & SCHWARZENBERGER, F. (2006): Endocrine and behavioral observations during transition of non-breeding into breeding season in female American bison (*Bison bison*). *Theriogenology* **66**: 1107–1114.
- WALKER, S. L., WADDELL, W. T. & GOODROWE, K. L. (2002): Reproductive endocrine patterns in captive female and male red wolves (*Canis rufus*) assessed by fecal and serum hormone analysis. *Zoo Biology* **21**: 321–335.
- WALTER, W. D., PEKINS, P. J., RUTBERG, A. T. & KILPATRICK, H. J. (2002): Evaluation of immunocontraceptive adjuvants, titers, and fecal pregnancy indicators in free-ranging white-tailed deer. *Wildlife Society Bulletin* **30**: 908–914.
- WASSER, S. K. & HUNT, K. E. (2005): Noninvasive measures of reproductive function and disturbance in the barred owl, great horned owl, and northern spotted owl. *Annals of the New York Academy of Sciences* **1046**: 109–137.
- WASSER, S. K., HUNT, K. E., BROWN, J. L., COOPER, K., CROCKETT, C. M., BECHERT, U., MILLSAUGH, J. J., LARSON, S. & MONFORT, S. L. (2000): A generalized fecal glucocorticoid assay for use in a diverse array of nondomestic mammalian and avian species. *General and Comparative Endocrinology* **120**: 260–275.
- WASSER, S. K., DAVENPORT, B., RAMAGE, E. R., HUNT, K. E., PARKER, M., CLARKE, C. & STENHOUSE, G. (2004): Scat detection dogs in wildlife research and management: application to grizzly and black bears in the Yellowhead Ecosystem, Alberta, Canada. *Canadian Journal of Zoology* **82**: 475–492.
- WEI, F., LÜ, X., LI, C., LI, M., REN, B. & HU, J. (2005): Influences of mating groups on the reproductive success of the southern Sichuan red panda (*Ailurus fulgens styani*). *Zoo Biology* **24**: 169–176.
- WEINGRILL, T., GRAY, D. A., BARRETT, L. & HENZI, S. P. (2004): Fecal cortisol levels in free-ranging female chacma baboons: relationship to dominance, reproductive state and environmental factors. *Hormones and Behavior* **45**: 259–269.
- WELLS, A., TERIO, K. A., ZICCARDI, M. H. & MUNSON, L. (2004): The stress response to environmental change in captive cheetahs (*Acinonyx jubatus*). *Journal of Zoo and Wildlife Medicine* **35**: 8–14.
- WHEATON, C. J., JOSEPH, S., REID, K., WEBSTER, T., RICHARDS, M. & SAVAGE, A. (2006): Body weight as an effective tool for determination of onset of puberty in captive female Nile hippopotami (*Hippopotamus amphibius*). *Zoo Biology* **25**: 59–71.
- WHITTEN, P. L., BROCKMAN, D. K. & STAVISKY, R. C. (1998): Recent advances in noninvasive techniques to monitor hormone-behavior interactions. *Yearbook of Physical Anthropology* **41**: 1–23.
- WIELEBNOWSKI, N. (2003): Stress and distress: evaluating their impact for the well-being of zoo animals. *Journal of the American Veterinary Medical Association* **223**: 973–977.
- WIELEBNOWSKI, N. C., FLETCHALL, N., CARLSTEAD, K., BUSSO, J. M. & BROWN, J. L. (2002): Noninvasive assessment of adrenal activity associated with husbandry and behavioral factors in the North American clouded leopard population. *Zoo Biology* **21**: 77–98.
- WIELEBNOWSKI, N. C., ZIEGLER, K., WILDT, D. E., LUKAS, J. & BROWN, J. L. (2002): Impact of social management

- on reproductive, adrenal and behavioural activity in the cheetah (*Acinonyx jubatus*). *Animal Conservation* **5**: 291–301.
- WITTEMYER, G., GANSWINDT, A. & HODGES, K. (2007): The impact of ecological variability on the reproductive endocrinology of wild female African elephants. *Hormones and Behavior* **51**: 346–354.
- WOODD, C., CZARNY, N. A., GUNN, I. M. G. & STURROCK, W. (2006): Faecal steroid analysis and urinary cytology of the squirrel glider (*Petaurus norfolcensis*). *Australian Journal of Zoology* **54**: 277–285.
- WYSOCKI, L. E., DITTAMI, J. P. & LADICH, F. (2006): Ship noise and cortisol secretion in European freshwater fishes. *Biological Conservation* **128**: 501–508.
- YOUNG, K. M., BROWN, J. L. & GOODROWE, K. L. (2001): Characterization of reproductive cycles and adrenal activity in the black-footed ferret (*Mustela nigripes*) by fecal hormone analysis. *Zoo Biology* **20**: 517–536.
- YOUNG, K. M., WALKER, S. L., LANTHIER, C., WADDELL, W. T., MONFORT, S. L. & BROWN, J. L. (2004): Noninvasive monitoring of adrenocortical activity in carnivores by fecal glucocorticoid analyses. *General and Comparative Endocrinology* **137**: 148–165.
- YOUNG, A. J., CARLSON, A. A., MONFORT, S. L., RUSSELL, A. F., BENNETT, N. C. & CLUTTON-BROCK, T. (2006): Stress and the suppression of subordinate reproduction in cooperatively breeding meerkats. *Proceedings of the National Academy of Sciences* **103**: 12005–12010.
- ZIEGLER, T. E. & SOUSA, M. B. C. (2002): Parent–daughter relationships and social controls on fertility in female common marmosets, *Callithrix jacchus*. *Hormones and Behavior* **42**: 356–367.
- ZIEGLER, T. E. & WITTWER, D. J. (2005): Fecal steroid research in the field and laboratory: improved methods for storage, transport, processing, and analysis. *American Journal of Primatology* **67**: 159–174.

Manuscript submitted 2 February 2007;  
revised 27 March 2007; accepted 12 April  
2007